Productivity Losses of Chronic Diseases Among Canadian Labour Force in 1994 & 2005: Estimates from a Nationally Representative Samples

Farah Farahati, Ph.D.

Conferences for Health Over the Life Course University of Western Ontario, October 15, 2009



Objective: To Estimate the Productivity Labour-Related Losses from:

- 22 Chronic Disorders including:
 - Diabetes Mellitus (DM),
 - DM-Related Comorbidities (DRCOM),
 - Other non-DM related Chronic disorders (e.g., arthritis, cancer),
 - Some risk factors: BMI, smoking, # of regular drinkers (> 3 drinks/week), and physical exercise.

Motivations: Burden of DM Projected to Grow

- WHO: worldwide prevalence of DM is increasing from 4.0% in 1995 to 5.4% in 2025
- ADA: Economic cost of DM is increasing from \$132B in 2002 to \$192B in 2020, in the US
- Reason:
 - Growing prevalence, decreasing age of onset, aging population, changes in diabetes-related service utilization, and a greater understanding of the wide range of diabetes-related comorbidities (DRCOM)

Previous Studies: Economic Burden of DM

- ...no standard method has been established.."Ettaro et al, 2004
- Economic cost of DM in US. In 2002: \$132B (ADA)
 - Direct medical costs: US\$91.8B
 - Indirect costs: \$US40B (lost workdays, restricted activity & early mortality)
- Decreasing age of onset & increasing rate of DRCOM among teenagers suggests even greater burden of DM on productivity in future
 - Accurately estimating the social costs hampered by the unavailability of good data.
 - Non-representative samples or impute key data, which can introduce bias (e.g., Kraut et al. 2001 (Manitoba), ADA))

Hypothesis

DM, DRCOM increase the probability of having disability days, the number of disability days, and earnings, ceteris paribus.

The multivariate labor supply model:

 $D = f(X, DM, DRCOM, other chronic disorders \xii)$

 H_0 : $\partial D / \partial DM$ related disorders ≥ 0

 H_1 : $\partial D / \partial DM$ related disorders< 0



Data: NPHS 1994 & CCHS 2005 National Population Health Survey Canadian Community Health Survey

- Ages between 20-65, excluding students & retired
- NPHS: 5,627 women and 4,867 men
- CCHS: 32,637 women and 30,119 men
 - If Disability days > 0 = 1, else = 1(0,1), #Disability days/year, annual earnings
 - Socio-demographics (age, sex, education, marital status, #kids, regional economic conditions)
 - DM, DRCOM, Other chronic disorders, #drinking, smoking



- A two-part model is used to estimate the impact of DM on labour market outcomes:
 - Part I: logistic regression for probability of having disability day
 - part II: log-transformed OLS regression to estimate
 # of disability days and annual earnings, for workers
- Part I x Part II: to estimate #disability days & earnings losses for all sample



Empirical Model: Annual Productivity Loss (PL), per Diabetic

 $PL_i = (DisD_{i,NDM}E_{i,NDM}) - (DisD_{i,DM}E_{i,DM})$

where:

i = individual i with DM

DisD_{i,DM}=predicted disability days/yr for person i if has DM

DisD_{i, NDM} = predicted disability days/yr for i if didn't have DM

E_{i,DM}= predicted annual earnings of person i if has DM

E_{i,NDM} = predicted annual earnings of person i if didn't have

 These individual-level costs are then inflated to national costs using the survey population weights.




Table 1: Sample Descriptive Statistics, Means (T-Test between women & men)									
Data / Variable	NPHS 1994			CCHS 2005					
	Full Sample	Women	Men	Full Sample	Women	Men			
	(N = 10494) *	(N = 5627)	(N = 4867))	$(N = 62756)^*$	(N = 32637)	(N =30119)			
Dependent variable									
Disability days	0.15(0.003)	.165a(0.005)	0.12(0.005)	0.185a(0.002)	0.216(0.002)	0.15 (0.002)			
status (0,1)									
Disability days	0.85(0.03)	0.961a(.039)	0.727 (0.04)	0.968a(0.011)	1.12(0.017)	0.80(0.015)			
Annual income	44,206(264)	42302 a(359)	46410 (388)	54,099a(148)	50,944(206)	57,518(210)			
Income(C\$93&\$04)		21,674a(404)	34088 (462)	46,575a(159)	40988(220)	5,230 (225)			
	Independent Variables								
DM	0.020 (0.001)		.019 (0.002)	0.037a(0.001)	0.035(0.001)	0.04 (0.001)			
DRCOM	0.091 (0.003)	.096 (0.004)	.08 (0.004)	0.150b(0.001)	0.147(0.002)	0.15 (0.002)			
Other Chronic †	0.40 (0.005)	.451 (0.007)	.33 (0.007)	0.596a(0.002)	0.665(0.003)	.522 (0.003)			
BMI	25.45 (0.041)	24.83 (0.06)	26.16(0.05)	26.28a(0.021)	25.73(0.031)	26.88 (0.03)			
Age	6.44 (0.022)	6.45 (0.030)	6.42 (0.032)	38.2a (3.80)	39.0 (4.10)	37.8 (4.2)			
Married	0.64 (0.310)	0.63 a (0.18)	0.64 (0.28)	0.620a(0.002)	0.638(0.003)	0.60 (0.003)			
< Secondary	0.219 (0.004)	0.212 (0.005)	.227 (0.006)	0.07b (0.001)	0.073 (0.001)	0.08 (0002)			
Secondary	0.167 (0.004)	0.174 (0.005)	.159 (0.01)	0.12 b (0 .001)	0.12 (0.002)	0.12 (0.002)			
Post-Secondary	0.378 (0.005)	0.369 (0.006)	.388 (0.007)	0.80 a (0.002)	0.81 (0.002)	0.79 (0.002)			
Kids < 11 years	0.194 (0.004)	0.22 (0.005)	.167 (0.005)	0.28 a (0.002)	0.309 (0.003)	0.26 (0.003)			
Smokers	0.368 (0.005)	0.35 (0.006)	.387 (0.007)	0.30 a (0.002)	0.283 (0.002)	0.32 (0.003)			
Drinkers(>3/week)	0.308 (0.005)	0.39 (0.007)	.21 (0.006)	0.11 a (0.001)	0.07 (0.001)	0.16 (0.002)			
Physical Exercise	0.40a (.01)	0.41 (.01)	0.38 (0.004)	0.50 (0.002)	0.51 (0.003)	0.50 (0.003)			
Unemployment	11.77 (0.028)	11 (0.036)	12.4 (0.041)	7.50 a (0.009)	7.18 (0.010)	7.80 (0.014)			
Rate									



Table 2. Impact of Diabetes on Labour Market Outcomes in Canada, in 1994 & 2005, women (SE)

	NPHS 1994			CCHS 2005			
	DISABILITY			DISABILITY			
	STATUS	LN DISABILITY	LN Annual	STATUS	LN DISABILITY	LN Annual	
	: Logistic	DAY AMONG	INCOME	: Logistic	DAY AMONG	INCOME	
	ANALYSIS,	THOSE WITH	AMONG	Analysis,	THOSE WITH	AMONG	
	Odds Ratio	DISABILITY	THOSE WITH	ODDS RATIO	DISABILITY	THOSE WITH	
		DAYS > 0	DISABILITY		DAYS > 0	DISABILITY	
DM	2.92 ***(0.848)	0.23 (0.18)	0.03 (0.09)	1.5 *** (0.15)	0.11 (0.069)	002 (0.19)	
DRCOM	1.69 ***(0.281)	0.35*** (0.11)	0.04 (0.09)	1.37 ***(0.088)	0.17 ***(0.05)	0.02 (0.12)	
Depression	2.20 ***(0.29)	0.226** (0.10)	-0.03 (0.08)	2.17 *** (0.15)	0.29*** (0.05)	-0.22 (0.17)	
Other	3.29***(0.73)	0.202 (0.13)	0.055 (0.07)	2.85 *** (0.15)	0.12 *** (0.04)	0.07 (0.17)	
Chronics							
Smoking	1.15 (0.12)	0.16* (0.08)	-0.03 (0.07)	1.35 ***	0.06* (0.03)	0.03 (0.1)	
Daily/occasio				(0.064)			
Regular	1.09 (0.11)	-0.09 (0.08)	-0.11 (0.08)	0.79 ***	-0.01 (0.07)	0.34 ***(0.13)	
Drinker > 3 /	, ,	' '	, ,	(0.07)	, , ,		
week				, ,			
Sample size	5627	931	358	32634	7065	7065	
***Panracente P.Value < 0.01 ** rapracente P.value < 0.05 * rapracente P.value < 0.10							

^{***}Represents P-Value < 0.01, ** represents P-value < 0.05, * represents P-value < 0.10
Reference Categories are single, divorced, less than high school education, and healthy women.




Table 2 A- (continued): Impact of Diabetes on Labour Market Outcomes, in Canada, for Women (standard error using bootstrap weights)

NPHS1994 **CCHS2005** DISABILITY DAYS LN DISABILITY DAY LN ANNUAL WORK STATUS LN DISABILITY DAY **STATUS** : LOGISTIC AMONG THOSE WITH INCOME AMONG THOSE WITH LN ANNUAL INCOME : LOGISTIC DISABILITY DAYS > 0 AMONG THOSE ANALYSIS. DISABILITY DAYS > AMONG THOSE ANALYSIS, WITH DISABILITY ODDS RATIO WITH DISABILITY ODDS RATIO DAYS > 0DAYS > 01.33** (0.189) 0.156 (0.183) 0.118(0.111)0.945 (0.075) 0.039 (0.056) 0.456** (0.187) Age Age^2 0.979 **(0.010) 0.004 (0.008) 0.006 (0.008) 1.00 (0.005) 0.0002 (0.004) -0.028** (0.012) Married 0.833* (0.088) 0.403*** (0.067) 0.822*** (0.039) 0.551 ***(0.098) -0.030 (0.080) -0.031 (0.034) Less Than High 0.957 (0.131) 0.408*** (0.098) -0.453*** (.089) 0.930 (0.100) 0.0003 (0.072) -0.181 (0.509) school College 1.08 (0.120) 0.093 (0.091) -0.168** (0.084) 1.146 * (0.081) -0.177*** (0.049) 1.005** (0.375) Any Kids < 11 years 0.835 (0.110) -0.059 (0.102) 0.008 (0067) -0.046 (0.038) -0.017 (0.129) 1.01 (0.054) in House-Hold Unemployment Rate 0.901*** (0.023) -0.010 (0.019) 0.009 (0.016) 0.959 *** (0.012) 0.001 (0.008) -0.018 (0.037) 1.139*** (0.390) 10.09*** (0.382) 0.897*** (0.212) 7.52*** (0.836) Intercept Observations 5627 931 358 32634 138.79 0.115 695.46 7065 4292 Wald Chi2 (17) Log-pseudo--15773 -2349 likelihood Pseudo R2 0.0463 0.0539 R-Squared 0.0504 0.0479 Root MSE .87265 .47539 .86904 2.0916



	NPHS 1994			CCHS 2005			
	DISABILITY STATUS : LOGISTIC ANALYSIS, ODDS RATIO	LN DISABILITY DAY AMONG THOSE WITH DISABILITY DAYS > 0	LN ANNUAL INCOME AMONG THOSE WITH DISABILITY DAYS > 0	DISABILITY STATUS : LOGISTIC ANALYSIS, ODDS RATIO	LN DISABILITY DAY AMONG THOSE WITH DISABILITY DAYS > 0	LN ANNUAL INCOME AMONG THOSE WITH DISABILITY DAYS > 0	
DM	0.68 (0.27)	. 0.03 (0.31)	0.10 (0.23)	1.15 (0.14)	0.13 (0.10)	32 (0.27)	
DRCOM	2.0 *** (0.38)	0.22 (0.17)	0.07 (0.10)	1.55*** (0.11)	0.15 ***(0.06)	0.02 (0.11)	
Greater than 2 weeks Depression	2.10 *** (0.43)	0.57 ***(0.17)	05 (0.10)	2.77 ***(0.26)	0.36 (0.07)	.56 *** (0.21)	
Other Chronics	2.47*** (0.74)	0.18 (0.20)	02 (0.12)	2.38*** (0.13)	.17 (0.04)	-0.13 (0.10)	
Smoking Daily/occasionally	1.22 (0.16)	.01 (0.11)	05 (0.08)	1.23 ***(0.07)	.12*** (0.04)	-0.25 ***(0.10)	
Regular Drinker > 3 / week	1.11 (0.15)	0.06 (0.11)	16* (0.08)	0.97 (0.07)	06 (0.06)	.14 (0.12)	
Sample size	4867	600	340	30116	4525	3244	

^{***}Represents P-Value < 0.01, ** represents P-value < 0.05, * represents P-value < 0.10
Reference Categories are single, divorced, less than high school education, and healthy men.



Table 2. B- (continued): Impact of Diabetes on Labour Market Outcomes, in Canada, for Men (standard error)

NPHS1994 CCHS2005 DISABILITY DAYS LN DISABILITY DAY **STATUS** WORK STATUS LN DISABILITY DAY LN WEEKLY WAGE AMONG THOSE WITH LN WEEKLY WAGE AMONG THOSE : LOGISTIC AMONG THOSE : LOGISTIC DISABILITY DAYS > 0 AMONG THOSE WITH ANALYSIS, DISABILITY DAYS > 0 WITH DISABILITY ANALYSIS, WITH DISABILITY ODDS RATIO ODDS RATIO DAYS > 0DAYS > 01.085 (0.181) 0.066 (0.138) 0.118 (0.111) 0.945 (0.075) 0.039 (0.056) 0.456** (0.187) Age 0.990 (0.012) 0.006 (0.008) -0.028** (0.012) Age^2 0.001 (0.010) 1.00 (0.005) 0.0002 (0.004) Married 1.193 (0.171) -0.068 (0.114) 0.403*** (0.067) 0.822*** (0.039) -0.031 (0.034) 0.551 ***(0.098) Less Than High 0.903 (0.151) 0.376*** (0.144) -0.453*** (.089) -0.181 (0.509) 0.930 (0.100) 0.0003 (0.072) school -0.177*** (0.049) College 1.130 (0.165) 0.233* (0.123) -0.168** (0.084) 1.146 * (0.081) 1.005** (0.375) Any Kids < 11 years 1.041 (0.191) -0.177 (0.162) 0.008 (0067) 1.01 (0.054) -0.046 (0.038) -0.017 (0.129) in House-Hold Unemployment Rate 0.949 (.029) 0.003 (0.019) 0.009 (0.016) 0.959 *** (0.012) 0.001 (0.008) -0.018 (0.037) Intercept 0.496 (0.487) 10.09*** (0.382) 0.897*** (0.212) 7.52*** (0.836) Observations 4867 600 358 32634 7065 4292 Wald Chi2 (17) 58..79 695.46 Log-pseudo-0.126 -15773 likelihood -1749 0.929 Pseudo R2 0.0223 .87265 0.0539 R-Squared 0.0504 0.0479 Root MSE .47539 .86904 2.0916



Limitations of Study

- In self reported survey many are unaware of having DM
- We assumed DM is determined exogenously of work conditions
- We assumed DRCOM are consequences of DM not preceded DM

Results

- The average # of disability days increased from 0.85 to 0.96 (during the past 2 weeks of the survey) for both men and women.
 - While the average prevalence of DM, DRCOM, 16 non-DM related chronic disorders increased, depression decreased for men and women.
 - Among risk factors, # of smokers dropped, # of regular drinkers and those who had physical exercise increased, however, BMI increased for both men and women.




Results

The productivity losses (probability of having disability days, number of disability days, and earnings) due to DM & DRCOM are significant in both 1994 and 2005 for women and men.

While the productivity losses associated with DM, DRCOM, depression, and sixteen other non-DM related disorders decreased in 1994 & 2005 for women, they increased for men during these years only for DM, but decreased for other chronic disorders.

Conclusion



Few studies suggest that people are behaving healthier. The number of deaths due to heart disease decreased during the past decade due to better risk factors, such as, decreased number of smokers, increased number of regular drinkers and those who exercise regularly.

This is the first study in our knowledge estimating the productivity losses of chronic disorders in 2 points of time by controlling for risk factors and socio-demographic characteristics using nationally representative sample. Our study results suggest that prevention through risk factors may decrease disability days beside rate of mortality and morbidity. This study especially could be replicated by longitudinal version of NPHS 1994 and 2009.



Thank You!

Questions? Suggestions?