

2012

Durand et al 2012 Supplemental figures

M. Durand
Université de Sherbrooke

Svetlana V. Komarova
McGill University


Ajay Bhargava

Keying Li

Cara Fiorino

See next page for additional authors

Follow this and additional works at: <http://ir.lib.uwo.ca/physpharmpub>

 Part of the [Diagnosis Commons](#), [Medical Physiology Commons](#), [Musculoskeletal, Neural, and Ocular Physiology Commons](#), and the [Therapeutics Commons](#)

Citation of this paper:


Durand, M.; Komarova, Svetlana V.; Bhargava, Ajay; Li, Keying; Fiorino, Cara; Maria, Osama; Nabavi, Noushin; Manolson, Morris; Harrison, Rene; Dixon, S. Jeffrey; Sims, Stephen; Mizianty, Marcin; Kurgan, Lukasz; Haroun, Sonia; Boire, Gilles; Lucena-Fernandes, Maria; and de Brum-Fernandes, Artur, "Durand et al 2012 Supplemental figures" (2012). *Physiology and Pharmacology Publications*. Paper 71.
<http://ir.lib.uwo.ca/physpharmpub/71>


Authors

M. Durand, Svetlana V. Komarova, Ajay Bhargava, Keying Li, Cara Fiorino, Osama Maria, Noushin Nabavi, Morris Manolson, Rene Harrison, S. Jeffrey Dixon, Stephen Sims, Marcin Mizianty, Lukasz Kurgan, Sonia Haroun, Gilles Boire, Maria Lucena-Fernandes, and Artur de Brum-Fernandes

Supplementary Figure 1

Durand et al. "Monocytes from Patients with Osteoarthritis Display Increased Osteoclastogenesis and Bone Resorption: The *In Vitro* Osteoclast Differentiation in Arthritis (IODA) Study"






Supplementary Figure 1. The gene expression of RANK, c-fms, cathepsin K, MMP9, NFATc1 and CCR1 in controls and OA patients. PBMCs were differentiated for 21 days, mRNA was isolated and the expression of RANK (A), c-fms (B), cathepsin K (C), MMP9 (D), NFATc1 (E) and CCR1 (F) was assessed by quantitative RT-PCR with β -actin used as an endogenous control. Means \pm standard error of the mean are shown. There were no statistically significant differences in the expression of these genes in osteoclasts derived from patients with OA and controls. Within the group of OA patients, treatment with bisphosphonates, the presence of osteoporosis or BMI had no significant impact on the expression of studied genes.

Supplementary Figure 2

Durand et al. "Monocytes from Patients with Osteoarthritis Display Increased Osteoclastogenesis and Bone Resorption: The *In Vitro* Osteoclast Differentiation in Arthritis (IODA) Study"



Supplementary Figure 2. Six representative immunoblots used for the analyses illustrated in Figure 4. Proteins were isolated from *in vitro* differentiated OCs from control (C) and OA (OA) patients. Blots were probed with antibodies directed against IL-1R1, IL-1R2, RANK and GAPDH. For quantification of IL1-R1, doublets were treated as a single band. Samples were excluded from subsequent analysis if bands were distorted or the signal was not within the linear range of the chemiluminescence detector (e.g. lanes marked with *).