

Spring 4-2018

Maternal Nicotine Exposure Leads to Augmented Expression of the Antioxidant Adipose Tissue Triglyceride Lipase Long-Term in the White Adipose of Female Rat Offspring.

Nicole Barra
barrang@mcmaster.ca

Taylor VanDuzer

Alison C. Holloway
hollow@mcmaster.ca

Daniel B. Hardy
Physiology and Pharmacology, daniel.hardy@schulich.uwo.ca

Follow this and additional works at: <https://ir.lib.uwo.ca/physpharmpub>

Part of the [Medical Physiology Commons](#), [Pharmacology Commons](#), and the [Pharmacy and Pharmaceutical Sciences Commons](#)

Citation of this paper:

Barra, Nicole; VanDuzer, Taylor; Holloway, Alison C.; and Hardy, Daniel B., "Maternal Nicotine Exposure Leads to Augmented Expression of the Antioxidant Adipose Tissue Triglyceride Lipase Long-Term in the White Adipose of Female Rat Offspring." (2018). *Physiology and Pharmacology Publications*. 108.
<https://ir.lib.uwo.ca/physpharmpub/108>

Maternal nicotine exposure (MNE) leads to augmented expression of the anti-oxidant adipose tissue triglyceride lipase (ATGL) long-term in the white adipose of female rat offspring

Journal:	<i>Toxicological Sciences</i>
Manuscript ID	TOXSCI-17-0684.R1
Manuscript Type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Barra, Nicole ; University of Western Ontario, Physiology and Pharmacology VanDuzer, Taylor; McMaster University Faculty of Health Sciences, Obstetrics and Gynaecology Holloway, Alison; McMaster University, Department of Obstetrics and Gynecology; Hardy, Daniel; University of Western Ontario, Physiology and Pharmacology
Key Words:	white adipose tissue, maternal nicotine exposure, adipocytes, antioxidants < Agents, inflammation < Immunotoxicology

1
2
3 Maternal Nicotine Exposure (MNE) leads to augmented expression of the anti-oxidant adipose
4
5 tissue triglyceride lipase (ATGL) long-term in the white adipose of female rat offspring
6
7
8
9

10 Nicole G. Barra*, Taylor VanDuzer †, Alison C Holloway† and Daniel B Hardy*‡

11
12 *Department of Physiology and Pharmacology, Western University, London, Ontario, Canada
13
14 N6A 5C1
15

16
17 †Department of Obstetrics and Gynecology, McMaster University, Hamilton, Ontario, Canada
18
19 L8S 4K1
20

21 ‡Departments of Obstetrics and Gynecology, Children's Health Research Institute, Lawson,
22
23 Health Research Institute, Western University, London, Ontario, Canada N6C 2V5
24
25
26
27

28 *Corresponding author:

29
30 Dr. Daniel Hardy

31
32 Department of Physiology and Pharmacology

33
34 Dental Sciences Building Room 2023

35
36 University of Western Ontario

37
38 London, Ontario, Canada

39
40
41 N6A 5C1

42
43
44 Email: daniel.hardy@schulich.uwo.ca

45
46
47 Tel: 519-661-2111 ext. 84238

48
49
50 Fax: 519-661-3827

51
52
53
54 **RUNNING TITLE:** The effect of MNE on white adipose tissue
55
56
57
58
59
60

ABSTRACT

Globally, approximately 10-25% of women smoke during pregnancy. Since nicotine is highly addictive, women may use nicotine containing products like nicotine replacement therapies for smoking cessation, but the long-term consequences of early life exposure to nicotine remain poorly defined. Our laboratory has previously demonstrated that maternal nicotine exposed (MNE) rat offspring exhibit hypertriglyceridemia due to increased hepatic *de novo lipogenesis*. Hypertriglyceridemia may also be attributed to impaired white adipose tissue (WAT) lipid storage; however, the effects of MNE on WAT are not completely understood. We hypothesize that nicotine-induced alterations in adipose function (*e.g.* lipid storage) underlie dyslipidemia in MNE adults. Female 6 month old rats exposed to nicotine during gestation and lactation exhibited significantly decreased visceral adipocyte cell area by 40%, attributed, in part, to a 3-fold increase in adipose triglyceride lipase (ATGL) protein expression compared to vehicle. Given ATGL has antioxidant properties and *in utero* nicotine exposure promotes oxidative stress in various tissues, we next investigated if there was evidence of increased oxidative stress in MNE WAT. At both 3 weeks and 6 months, MNE offspring expressed 37-48% higher protein levels of SOD1 and SOD2 in WAT. Since oxidative stress can induce inflammation, we examined the inflammatory profile of WAT and found increased expression of cytokines (IL-1 β , TNF α , and IL-6) by 44-61% at 6 months. Collectively, this suggests that the expression of WAT ATGL may be induced to counter MNE-induced oxidative stress and inflammation. However, higher levels of ATGL would further promote lipolysis in WAT, culminating in impaired lipid storage and long-term dyslipidemia.

KEY WORDS: White adipose tissue, maternal nicotine exposure, oxidative stress, adipocytes

INTRODUCTION

Evidence now demonstrates that cigarette smoke plays a critical role in the development of dyslipidemia and obesity in children exposed during perinatal life (Wen *et al.*, 2010; Weng *et al.*, 2012). This is especially concerning given that approximately 10-25% of women still smoke during pregnancy, with rates as high as 59% in certain Indigenous communities (Cui *et al.*, 2014; Roman-Galvez *et al.*, 2017; Tappin *et al.*, 2010; Tong *et al.*, 2013). To date, nicotine replacement therapies (NRT) for smoking cessation (*i.e.* transdermal patches, gums, e-cigarettes) are thought to benefit pregnant women who are highly addicted and unable to quit smoking by other means (Oncken and Kranzler, 2003). However, the consequences of fetal and neonatal exposure to nicotine alone on the long-term metabolic health of the offspring have yet to be fully defined.

Maternal nicotine use during gestation and lactation leads to significant exposure to the fetus and neonate since it is present in fetal blood and transferred through amniotic fluid, placental tissue, and breast milk (Luck and Nau, 1987; Luck *et al.*, 1985). Numerous animal studies demonstrate that nicotine use during pregnancy leads to adverse neurobehavioral, pulmonary, cardiovascular, and metabolic outcomes in the offspring (Barra *et al.*, 2017; Chou and Chen, 2014; Dasgupta *et al.*, 2012; Ma *et al.*, 2014; Pauly and Slotkin, 2008). Specifically, we and others have shown in rats that perinatal exposure to nicotine alone leads to increased blood pressure, adiposity, decreased glucose tolerance, and impaired pancreatic beta cell development (Fox *et al.*, 2012; Gao *et al.*, 2008; Gao *et al.*, 2005; Holloway *et al.*, 2008). Moreover, MNE during gestation and lactation leads to increased hepatic *de novo* lipogenesis and increased hepatic and circulating triglycerides in postnatal life (Ma *et al.*, 2014). This reciprocates human studies which have demonstrated that children exposed to maternal smoke *in*

1
2
3 *utero* have higher triglyceride levels later in life (Cupul-Uicab *et al.*, 2012). Since increased
4 circulating plasma triglycerides is strongly associated with cardiovascular disease (CVD) (Bansal
5 *et al.*, 2007), elucidating the underlying molecular mechanisms promoting dyslipidemia will
6 undoubtedly uncover therapeutic targets to reduce unwarranted CVD risk to these nicotine-
7 exposed individuals.
8
9

10
11
12
13
14
15 Aside from the liver, the augmented circulating and hepatic triglyceride accumulation
16 observed in these MNE offspring could also be attributed to impairments in other metabolic
17 tissues, such as white adipose tissue (WAT). The major role of WAT is to store excess
18 circulating triglycerides to prevent glucose and lipid toxicity, and moreover, ectopic fat
19 deposition (Abate, 2012). Excess circulating triglycerides promote adipocytes to undergo
20 differentiation, hypertrophy, and/or hyperplasia to process and store lipids. Once the maximum
21 triglyceride storage capacity is reached, fatty acids can spillover in the plasma, increasing
22 substrate availability for hepatic triglyceride synthesis (Abate, 2012). Ultimately,
23 hypertriglyceridemia contributes to various systemic abnormalities like dyslipidemia, insulin
24 resistance, and CVD (Abate, 2012). Currently, the effects of MNE on white adipose tissue
25 function are not completely understood. Studies by Somm *et al.* indicated that MNE during
26 gestation resulted in adipocyte hypertrophy and increased expression of adipogenic transcription
27 factors in 3 week male offspring, but the long-term effects were not examined (Somm *et al.*,
28 2008). A more recent study by Fan *et al.* (2016) demonstrated that maternal exposure of nicotine
29 (from gestational day 9 to weaning) led to decreased adipocyte size in 26 week old male
30 offspring, suggesting an impaired capacity for lipid storage. Furthermore, this study also found
31 an increase in steady-state mRNA levels of adipocyte differentiation and lipogenic markers in
32 MNE WAT compared to vehicle (Fan *et al.*, 2016). However, given most women addicted to
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 smoking would be exposed to nicotine (*i.e.* NRT or cigarettes) in both prenatal and perinatal life,
4
5 the goal of this study was to determine if this longer, more relevant, window of nicotine exposure
6
7 impacts adipose function in *both* male and female offspring. Moreover, this window covers the
8
9 entire period of rat adipose tissue differentiation (Greenwood and Hirsch, 1974). We hypothesize
10
11 that *in utero* nicotine exposure will adversely impact WAT in the offspring, leading to
12
13 dyslipidemia in adulthood. To examine this further, we will investigate if WAT function is
14
15 affected due to altered adipocyte size, proliferation (*i.e.* Akt-1), differentiation (*i.e.* C/ebp α / β ,
16
17 Srebp-1c), lipogenesis (*i.e.* Lpl, Acc α , Fas, Acsl1), fatty acid transport (*i.e.* Fatp1, Fatp4), and/or
18
19 lipolysis (Atgl) in our well-established rat model of maternal nicotine exposure.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

MATERIALS AND METHODS

Maternal Nicotine Exposure (MNE) Rat Model

Nulliparous 200-250g female Wistar rats (Harlan, Indianapolis, IN, USA) were injected daily subcutaneously with either saline (vehicle) or nicotine bitartrate at 1 mg/kg/day (Sigma-Aldrich, St. Louis, MO, USA) two weeks prior to mating, during gestation, and until weaning (PND21). This nicotine dose results in maternal serum cotinine concentrations of 135.9 ± 7.86 ng/mL, which is comparable to “moderate” female smokers (80 ng/mL) or NRT users (169.9 ng/mL) (Holloway *et al.*, 2006; Shahab *et al.*, 2016). Litters were culled to eight at birth and following weaning, rats were housed as sibling pairs until 6 weeks of age, then subsequently housed individually. Male and female offspring were sacrificed via carbon dioxide inhalation at 3 weeks and 6 months of age. Gonadal white adipose tissues were extracted, either fixed in formalin or frozen in liquid nitrogen and stored at -80°C for histological and molecular analyses, respectively. All rats were conventionally housed in polycarbonate microisolator cages with ad libitum access to water and standard chow diet (Teklad 22/5 rodent diet; Envigo) under controlled lighting (12:12 L:D), humidity (40-50%), and temperature (22°C). In accordance with the Canadian Council for Animal Care guidelines, animal experiments were approved by the Animal Research Ethics Board at McMaster University.

Gonadal Adipocyte Cell Area

A portion of gonadal WAT was fixed in 10% (v/v) neutral buffered formalin overnight, washed in water, and embedded in paraffin. Cross sections were stained with hematoxylin and eosin (H&E) and photographed using the Olympus BX50 microscope under a 10x objective. Cell areas were quantified from a minimum of 160 adipocytes from two cross sections/rat using Northern

1
2
3 Eclipse software (Empix Imaging Inc.).
4
5
6
7

8 **RNA Extraction and Real Time-Polymerase Chain Reaction (RT-PCR)**

9

10 Total RNA was extracted from homogenized 3 week and 6 month WAT samples using TRIzol
11 reagent according to the manufacturer's instructions (Invitrogen). Two micrograms of RNA were
12 reversed-transcribed to cDNA (high-capacity cDNA Reverse Transcription Kit, Applied
13 Biosystems). Forward and reverse primer sets used for RT-PCR, listed in Table 1, were designed
14 with the National Center for Biotechnology Information's primer designing tool. Relative
15 transcript abundance was determined using SensiFAST No-ROX SYBR Green Supermix
16 (FroggaBio) and the Bio-Rad CFX384 Real Time System. Samples were assayed in triplicate
17 and relative fold change was calculated using comparative cycle times (Ct) method normalized
18 to β -actin. The relative abundance was calculated using the formula $2^{\Delta\Delta Ct}$, where $\Delta\Delta Ct$ was the
19 normalized value.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

35 **Protein extraction and Western blot**

36

37 WAT was homogenized in RIPA buffer (50 mM Tris-HCL, pH 7.4, 150 mM NaCl, 1 mM
38 EDTA, 1% Nonidet P40, 0.25% $C_{24}H_{39}NaO_4$) supplemented with a protease inhibitor cocktail
39 (Roche) and phosphatase inhibitors (20 mM NaF, 40mM Na-pyrophosphate, 40mM Na_3VO_4 ,
40 200mM β -glycerophosphate disodium salt hydrate). The solution was sonicated, mixed in a
41 rotator for 1hr at 4 °C, and centrifuged at 300g for 15 min at 4 °C. The total cellular protein
42 extract in the collected supernatant was quantified by colorimetric DC protein assay (BioRad).
43
44 Loading samples were heated at 50 °C for 10 min to denature the proteins. Proteins (20 μ g/well)
45 were separated by size via gel electrophoresis and transferred onto polyvinylidene difluoride
46 membrane (Millipore). Membranes were blocked in 1x Tris-buffered saline-Tween 20 buffer
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 with 5% non-fat milk or 5% BSA (blocking solutions), and then probed using primary antibodies
4
5 diluted in the blocking solution (Table 2). A mouse or rabbit secondary antibody was used to
6
7 detect primary antibody diluted in the blocking solution at 1:5,000 or 1:10,000 dilution,
8
9 respectively (Table 2). Immuno-reactive bands were visualized, and relative band intensity was
10
11 calculated using ImageLab software (BioRad) and normalized to β -actin, as previously
12
13 performed (Ma *et al.*, 2014).
14
15
16
17
18

19 **Statistical Analysis**

20
21 Statistical analyses were performed using Graphpad Prism 6 software. Results were
22
23 presented as the mean of arbitrary values \pm SEM. Grubbs' test was used to determine significant
24
25 outliers and data were tested for normality and equal variance. Comparisons between vehicle and
26
27 nicotine exposed offspring were assessed using an unpaired Student's t-test, with a p-value of
28
29 less than 0.05 deemed as significant.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RESULTS

Maternal nicotine exposure leads to decreased visceral adipocyte size in 6 month old offspring.

Using our well-established rat MNE model leading to dyslipidemia (Gao *et al.*, 2005; Ma *et al.*, 2014), we wanted to first confirm whether prenatal nicotine exposure has long term effects on adipocyte size in the offspring. Histological analyses revealed both MNE male and female offspring exhibited smaller gonadal adipocytes compared to vehicle controls at 6 months of age (Figure 1A, B). Given the more pronounced effects of perinatal nicotine exposure on female WAT area (13.2% male vs. 39.7% female decrease in adipocyte size), we continued to examine only female offspring to explore the underlying mechanisms involved. Gonadal adipose tissue weight did not differ between MNE and vehicle offspring (data not shown), suggesting a possible increase in adipocyte number due to prenatal nicotine exposure. With decreased adipocyte cell area and subsequent increase in cell number, transcript and protein levels of the proliferation marker AKT-1 were subsequently measured. Despite a significant increase in mRNA levels, there were no corresponding changes in AKT-1 protein levels in 6 month female MNE WAT (Figure 1C, D).

***In utero* nicotine exposure increases the expression of the differentiation marker C/EBP α in WAT of 6 month old female offspring.**

Given impairments in adipocyte differentiation can also decrease adipocyte size, we next measured the expression of key targets involved in adipogenesis (Abate, 2012), including CCAAT/enhancer-binding protein (C/EBP)- α/β and sterol regulatory element-binding

1
2
3 transcription factor (SREBP)-1c. Protein analyses revealed a significant increase in the 42kDa
4 band of C/EBP- α in female MNE offspring compared to vehicle at 6 months independent of
5 changes in transcript levels (Figure 2A, B). Quantitative real time PCR analyses revealed that
6
7
8 WAT from MNE exposed offspring had a significant increase in steady-state transcript levels of
9
10 C/EBP β and SREBP-1c compared to controls (Figure 2C, E). However, immunoblot analyses
11
12 revealed no significant differences at the protein level for either target (Figure 2D, F). To
13
14 examine if perinatal nicotine exposure had direct effects on C/EBP α , we measured the transcript
15
16 and protein levels at 3 weeks age but found no difference in expression from vehicle (Figure 2G,
17
18 H).

19
20
21
22
23
24
25
26 **Maternal nicotine exposure does not increase the expression of markers involved in**
27
28 **lipogenesis and fatty acid transporters in 6 month old female offspring.**

29
30 Adipocyte size is influenced by its ability to breakdown triglyceride (TG) molecules from
31
32 circulation, transport its constitutive components intracellularly, and re-assemble TGs for storage
33
34 (Kersten, 2014). Therefore, we wanted to determine whether perinatal nicotine exposure
35
36 impaired triglyceride transport and esterification by measuring the steady state transcript levels
37
38 of key targets involved in lipogenesis and fatty acid transport. At six months of age we found no
39
40 significant difference between groups in the expression of lipogenic markers including
41
42 lipoprotein lipase (LPL), acetyl-coA carboxlyase (ACC α), and fatty acid synthase (FAS), along
43
44 with fatty acid transporters FATP1/4, and long-chain acyl CoA synthetase (ASCL)-1 in female
45
46 MNE WAT (Figure 3A-F).
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 ***In utero* nicotine exposed female offspring exhibit increased expression of ATGL at 6**
4 **months, but not at 3 weeks.**
5

6
7 Since enhanced lipolysis decreases adipocyte size (Zhang *et al.*, 2017), we then examined
8 whether MNE WAT had elevated expression of adipose triglyceride lipase (ATGL), the
9 predominate enzyme involved in intracellular degradation (Zechner *et al.*, 2009). At six months
10 of age, we found a significant increase in both ATGL mRNA expression and protein levels in
11 nicotine exposed female WAT compared to vehicle controls (Figure 4A, B), however this was
12 not observed at 3 weeks (Figure 4C-D). Altogether, these data suggest that augmented ATGL-
13 induced lipolysis may lead to reduced adipocyte cell area in MNE WAT at 6 months.
14
15
16
17
18
19
20
21
22
23
24
25

26 **Maternal nicotine exposure exhibit enhanced anti-oxidant expression at 6 months and 3**
27 **weeks.**
28
29

30 Evidence demonstrates that ATGL deficiency in mice is associated with accumulation of reactive
31 oxygen species (ROS) and decreased SOD2 expression, suggesting this enzyme may act as an
32 antioxidant and suppress oxidative stress (Aquilano *et al.*, 2016; Chen *et al.*, 2017a; Chen *et al.*,
33 2017b). Moreover, we and others have previously shown that maternal nicotine exposure
34 promotes oxidative stress in the heart, pancreas, and placenta of adult offspring (Barra *et al.*,
35 2017; Bruin *et al.*, 2008; Sbrana *et al.*, 2011). Therefore, we wanted to determine whether an
36 imbalance between antioxidant defenses occurs in WAT of 6 month female MNE offspring by
37 measuring the transcript and protein levels of the antioxidants superoxide dismutase (SOD)-1
38 and SOD-2. Both SOD 1 and 2 steady state mRNA and protein levels were significantly elevated
39 in prenatal nicotine exposed WAT in 6 month female offspring compared to vehicle (Figure 5A-
40 D). At 3 weeks of age, there was also a significant increase in SOD-1 protein levels, and a
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 trending increase in SOD-2, in the nicotine exposed WAT group compared to controls (Figure
4 5E-H). To determine if there was oxidative stress, we measured protein levels of 4-HNE and
5 6 found a non-significant upward trend in nicotine-exposed WAT at both 3 week and 6 month
7 8 timepoints (Figure 5I, 5J). Overall, these data suggest that prenatal nicotine exposure directly
9 10 increase the expression of the antioxidant SOD-1 and SOD-2 in 6 month WAT which was
11 12 enough to prevent lipid oxidative damage.
13 14
15 16
17 18

19 ***In utero* nicotine exposure exhibit inflammation at 6 months, but not at 3 weeks.**

20
21 Enhanced oxidative stress can activate nuclear factor- κ B (NF- κ B), leading the expression of pro-
22 23 inflammatory cytokines and inflammation (Li and Karin, 1999; Morgan and Liu, 2011). As well,
24 25 activation of this pathway in differentiated 3T3-L1 cells leads to increased lipase expression and
26 27 lipolysis (Chi *et al.*, 2014). Since oxidative stress, inflammation, and lipolysis are integrated
28 29 pathways, we wanted to determine whether inflammation was present in 6 month *in utero*
30 31 nicotine exposed WAT. At 6 months, we found a significant increase in the steady-state mRNA
32 33 expression profile of pro-inflammatory cytokines tumor necrosis factor (TNF)- α , interleukin
34 35 (IL)-1 β , IL-6, and the macrophage marker CD68 (Figure 6A-D). In contrast at 3 weeks of age,
36 37 there was no significant difference in steady state mRNA levels of inflammatory markers
38 39 between groups (Figure 6E-H).
40 41
42 43
44 45
46 47
48 49
50 51
52 53
54 55
56 57
58 59
60

DISCUSSION

Under normal physiological conditions, WAT serves as the principle lipid storage site to prevent ectopic lipid deposition in peripheral organs. The balance between lipolysis and lipogenesis dictates the overall amount of triglyceride molecules stored as a single large lipid droplet in the cytoplasm of adipocytes, ultimately influencing its size (Bolsoni-Lopes and Alonso-Vale, 2015). Adipocytes have a limited triglyceride storage capacity. Once exceeded, excess lipids will spill-over into circulation resulting in dyslipidemia and increased lipid content at ectopic sites like the liver (Abate, 2012). In our model, we have previously shown that MNE offspring have dyslipidemia with increased levels of circulating triglycerides (Gao *et al.*, 2005; Ma *et al.*, 2014). While we have shown that hepatic de novo lipogenesis is increased in nicotine-exposed offspring, the effect of nicotine exposure on WAT function were not examined. Previous studies show maternal nicotine exposure during gestation or from gestation until weaning altered adipocyte size and increased the expression of adipocyte differentiation markers, but this was only examined in male offspring (Fan *et al.*, 2016; Somm *et al.*, 2008). Moreover, this exposure to nicotine in these studies (gestation only or mid-gestation to weaning only) does not mimic women's smoking habits and/or nicotine use prior to pregnancy. Therefore, our goal was to determine if WAT plays a role in promoting dyslipidemia in both male and female offspring exposed to nicotine in prenatal and perinatal life.

In this study, we demonstrated that both male and female offspring exposed to nicotine during prenatal and perinatal life decreased adipocyte cell area compared to vehicle controls at 6 months of age. Somm *et al.* reported that rats exposed to nicotine during pregnancy alone (from gestational day 4 for 14 days at 3mg/kg/day) had adipocyte hypertrophy compared to vehicle controls in 3 week old male offspring, but the effect on WAT size into adulthood, after nicotine

1
2
3 exposure, was not examined (Somm *et al.*, 2008). Similar to our findings, Fan *et al.*
4 demonstrated that *in utero* nicotine exposure from mid-pregnancy (*e.g.* gestational day 9) until
5 weaning decreased adipocyte size in male WAT at 26 weeks of age (Fan *et al.*, 2016). The dose
6 of nicotine (1 mg/kg) administered in this study is relevant as it corresponds to moderate female
7 smokers or NRT users (Holloway, Kellenberger and Petrik, 2006; Shahab *et al.*, 2016), while
8 doses used in other studies (2-3 mg/kg/day) correspond to heavy smokers (Fan *et al.*, 2016;
9 Somm *et al.*, 2008). Examining the effects of MNE on WAT for a longer duration is important
10 given adipose development occurs both pre- and postnatally in rats. In rodent adipose tissue,
11 growth and differentiation occurs from late gestation to 4 weeks in postnatal life, while in
12 humans this occurs from 5 to 29 weeks gestation (Greenwood and Hirsch, 1974; Poissonnet *et*
13 *al.*, 1984). In both species, adipose tissue expansion happens throughout life (Greenwood and
14 Hirsch, 1974; Spalding *et al.*, 2008). Given the extensive differentiation of adipose tissue during
15 perinatal life, it is vulnerable to alterations by environmental cues (*i.e.* drugs) during this
16 developmental window. Furthermore, both mature adipocytes and adipocyte precursors like
17 mesenchymal stem cells express nicotinic acetyl-choline receptor subunits, suggesting that
18 nicotine may directly alter adipogenesis and adipocyte formation (Gochberg-Sarver *et al.*, 2012;
19 Hoogduijn *et al.*, 2009).

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42 Since the effect on adipocyte size was more pronounced in female offspring, and
43 previous studies focused on males (Fan *et al.*, 2016; Somm *et al.*, 2008), we decided to examine
44 the effects of MNE on female adipose tissue function. One limitation of our study is that we did
45 not evaluate the estrous cycle of the female rats. This is an important consideration since prenatal
46 nicotine exposure increased progesterone levels in offspring compared to vehicle controls, which
47 is reflective of estrous cycle perturbations (Holloway, Kellenberger and Petrik, 2006).
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Furthermore, others have reported that oscillations in metabolic related genes in the liver occur
4 with estrous cycle (Villa *et al.*, 2012). Since small adipocyte size results from aberrant
5 adipogenesis, we examined whether MNE impaired adipocyte differentiation, lipid synthesis and
6 transport, and/or lipolysis. We found that female MNE offspring exhibited increased mRNA and
7 protein expression of the adipocyte differentiation factor C/EBP α and the lipolytic enzyme
8 ATGL, without altering the expression of targets involved in lipogenesis and lipid transport.
9
10 Similar to our results, Fan *et al.* found an increased in steady-state mRNA levels of adipocyte
11 differentiation markers in MNE male offspring, including C/EBP α and SREBP-1c compared to
12 control WAT in adulthood (Fan *et al.*, 2016). Since these studies did not perform protein
13 analyses, our study is the first to demonstrate that MNE increased C/EBP α protein levels in 6
14 month female WAT. Overall, the increased C/EBP α (42 kDa) expression in WAT suggests that
15 MNE promote adipocyte differentiation and adipogenesis. C/EBP α contains two isoforms (30/42
16 kDa) alternatively translated from one mRNA transcript (Lin *et al.*, 1993; Ossipow *et al.*, 1993).
17
18 Constitutive expression of the 42 kDa isoform can induce adipocyte differentiation in 3T3-L1
19 preadipocytes, suggesting that enhanced C/EBP α expression of this isoform can enhance
20 adipocyte formation (Lin and Lane, 1994). While the steady-state levels of AKT1, C/EBP- α ,
21 SREBP 1c mRNA were augmented in 6 month nicotine-exposed offspring, this did not translate
22 to an alteration in protein levels. There are several cases in the literature whereby changes in the
23 mRNA levels do not correlate to an alteration in protein, highlighting the importance of looking
24 at both mRNA and protein (Vogel and Marcotte, 2012). Moreover, it is possible that the mRNA
25 and protein levels of these key proliferation and differentiation markers may be altered earlier in
26 development (i.e. gestation and/or lactation) due to the direct effects of nicotine on their
27 expression. For example, we have demonstrated that nicotine directly enhances the expression of
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 the proliferation markers GADD45 and Rb1 in *differentiating* 3T3-L1 pre-adipocyte cells (data
4 not shown). Therefore, in the absence of nicotine exposure after lactation, the levels of these
5 proliferation and differentiation may have become normalized.
6
7
8
9

10 This study was the first to demonstrate that both prenatal and perinatal nicotine exposure
11 leads to increased ATGL mRNA and protein expression in adult female MNE offspring. This
12 suggests that MNE may impair the lipid storage capacity of WAT via increased ATGL
13 expression. Studies demonstrate that augmented ATGL leads to enhanced basal and stimulated
14 WAT lipolysis and reduced adipocyte size (Ahmadian *et al.*, 2009; Bezaire *et al.*, 2009;
15 Kershaw *et al.*, 2006). The primary function of ATGL is to initiate lipolysis, by hydrolyzing the
16 *sn*-2 ester bond of a triglyceride molecule, resulting in the formation of diacylglycerol and fatty
17 acid (Zechner *et al.*, 2009). Therefore, decreased adipocyte cell size in gonadal WAT may be
18 attributed to enhanced lipolysis in MNE offspring.
19
20
21
22
23
24
25
26
27
28
29

30 Recently, evidence supports an additional role for ATGL - as an antioxidant and
31 suppressor of oxidative stress. Oxidative stress is defined as an imbalance between antioxidant
32 defenses and ROS production (free radicals) (Sena and Chandel, 2012). Prolonged oxidative
33 stress can result in free radical damage, ultimately leading to mitochondrial-mediated apoptosis
34 (Sena and Chandel, 2012). *In vitro* and *in vivo* analyses demonstrate that ATGL deficiency is
35 associated with ROS accumulation and cellular apoptosis in renal podocytes and proximal
36 tubules (Chen *et al.*, 2017a; Chen *et al.*, 2017b). Similarly, ATGL inhibition in C2C12
37 myoblasts and in murine ATGL deficient skeletal muscle promotes oxidative damage and a
38 defective cellular antioxidant response, characterized by a significant reduction in antioxidant
39 SOD-2 expression and glutathione levels (Aquilano *et al.*, 2016). Altogether, these findings
40 suggest that ATGL may protect against oxidative stress. We and others have previously shown
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 that maternal nicotine exposure can cause oxidative stress in the offspring in various tissues
4 including the heart, pancreas, and placenta (Barra *et al.*, 2017; Bruin *et al.*, 2008; Sbrana *et al.*,
5
6 2011). Therefore, we examined the expression of antioxidants SOD-1/2 in WAT of 6 month
7
8 female MNE offspring to assess whether an imbalance in anti-oxidant expression is present at the
9
10 same time ATGL levels are elevated. Oxygen can oxidize other molecules to generate ROS,
11
12 whereby majority of intracellular ROS are derived from superoxide (O₂⁻). Superoxide is formed
13
14 in the mitochondrial respiratory chain, emitted to the matrix and intermembrane space, and
15
16 subsequently converted to hydrogen peroxide (H₂O₂) by superoxide dismutases (SODs) (Sena
17
18 and Chandel, 2012). This study is the first to show that MNE WAT had increased transcript
19
20 expression and protein levels of both SOD-1 and -2 at 6 months with evidence for increased
21
22 SOD-1 at 3 weeks of age. Although non-significant, we also observed an upward trend for the
23
24 oxidative marker 4-HNE in nicotine exposed WAT compared to vehicle at both time points. This
25
26 suggests that perinatal nicotine exposure may instigate WAT oxidative stress, triggering a rapid
27
28 increase in anti-oxidant expression by weaning which persists into adulthood and may prevent
29
30 long-term oxidative damage.
31
32
33
34
35
36

37
38 Oxidative stress can contribute to altered adipocyte differentiation and lipolysis.
39
40 Depending on environmental cues, mesenchymal stem cells can differentiate into various cell
41
42 types derived from the mesodermal lineage including myocytes, adipocytes, osteocytes, and
43
44 chondrocytes. Evidence suggests that elevated ROS levels, produced during oxidative stress, can
45
46 stimulate adipogenesis while suppressing osteogenesis (Atashi *et al.*, 2015). During conversion
47
48 of stem cells to adipocytes, mitochondrial complexes I/III and the NADPH oxidase isoform
49
50 NOX4 produce ROS to initiate adipocyte signaling cascades, leading to terminal differentiation
51
52 (Atashi, Modarressi and Pepper, 2015). Therefore, increased SOD-1 expression in fetal and
53
54
55
56
57
58
59
60

1
2
3 neonatal nicotine exposed WAT may suggest that augmented ROS production may directly
4 promote the expression of adipogenic transcription factors like C/EBP α (Hu *et al.*, 2005). *In vitro*
5 analyses using both differentiated 3T3-L1 and human primary adipocyte cultures demonstrate
6 that exposure to exogenous 4-HNE elevated lipolysis in basal and stimulated conditions
7 characterized by a dose-dependent significant increase in glycerol and fatty acid release (Zhang
8 *et al.*, 2013). Altogether, these findings suggest that MNE may augment oxidative stress in WAT
9 characterized by increased expression of antioxidants SOD-1/2 and ATGL, leading to altered
10 adipocyte differentiation and lipolysis.
11
12
13
14
15
16
17
18
19
20

21
22 Enhanced oxidative stress can promote pathways, like inflammation, that also influence
23 adipose function. Exposure to hydrogen peroxide or ROS accumulation activates the
24 transcription factor NF- κ B, leading the expression of pro-inflammatory cytokines such as TNF- α
25 or IL-1 (Lawrence, 2009; Li and Karin, 1999; Morgan and Liu, 2011). Intriguingly, treatment
26 with either TNF- α or IL-6 in differentiated adipocytes can increase the expression of ATGL and
27 enhance glycerol production (a marker of lipolysis). Moreover, *in vitro* studies reveal that TNF- α
28 induces lipolysis in differentiated human and murine adipocytes through multiple distinct
29 pathways (i.e. mitogen-activated protein kinase kinase, (MAPKK); extracellular signal-related
30 kinase 1/2 (ERK); and elevation of cyclic adenosine monophosphate (cAMP); NH₂-terminal
31 kinase (JNK)) (Ryden *et al.*, 2002; Yang *et al.*, 2011; Zhang *et al.*, 2002). Similarly, treatment
32 with IL-6 induces lipolysis through increased expression of ATGL in soleus muscles *ex vivo*
33 (Macdonald *et al.*, 2013) and enhanced glycerol production in 3T3-L1 cells (Ji *et al.*, 2011). Our
34 results reveal that fetal and neonatal nicotine exposure culminates in increased steady-state
35 mRNA levels of proinflammatory mediators, including TNF- α and IL-6 by 6 months of age.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 resulted in increased circulating pro-inflammatory cytokines throughout development and in 11
4 week old offspring (Mohsenzadeh *et al.*, 2014; Orellana *et al.*, 2014). Since inflammation was
5 present in 6 month female WAT and not at 3 weeks, this suggests that nicotine does not have a
6 direct effect on inflammation but that augmented oxidative stress may mediate the expression of
7 pro-inflammatory cytokines. One research group demonstrated that CD-1 mice fed a nicotine-
8 containing diet for 14 weeks postnatally decreased adipocyte cell size without affecting white
9 adipose inflammation (Liu *et al.*, 2018). In fact, treatment with 4-HNE in 3T3-L1 cells induced
10 inflammation by activating p38, subsequently increasing the expression of cyclooxygenase-2
11 (COX) (Zarrouki *et al.*, 2007). Further studies are warranted to determine whether inflammation
12 may in part promote lipolysis in prenatal nicotine exposed WAT long term.
13
14
15
16
17
18
19
20
21
22
23
24
25

26 In summary, our findings demonstrate that maternal nicotine exposure during pregnancy
27 and lactation leads to decreased WAT adipocyte size and impaired lipid storage in postnatal life
28 characterized by augmented lipolysis and inflammation triggered possibly by a pro-oxidative
29 stress environment. This impairment in WAT lipid storage can contribute to increased circulating
30 triglycerides, leading to enhanced hepatic *de novo* lipogenesis, high blood pressure, and
31 increased CVD risk as we have previously shown (Gao *et al.* 2008; Ma *et al.*, 2014). These
32 findings provide insight into the consequence of prolonged oxidative stress driving WAT
33 dysfunction long-term. Postnatal intervention strategies targeting oxidative stress may benefit
34 children exposed to nicotine *in utero* through cigarette smoke and/or NRT. In fact, studies
35 demonstrate that the use of antioxidants (i.e. anthocyanin, quercetin, bottle gourd) can
36 significantly reduce dyslipidemia (Li *et al.*, 2015; Talirevic and Jelena, 2012). Overall, results
37 from this study suggests that therapeutic strategies targeting oxidative stress may effectively
38 protect nicotine exposed offspring from WAT dysfunction and dyslipidemia.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FUNDING INFORMATION

This work was supported by the Canadian Institutes of Health Research [grant numbers MOP86474 to ACH, MOP111011 to DBH] and a research grant from the Women's Development Council to DBH. NGB is a recipient of the Whaley Fellowship and Molly Towell Perinatal Research Foundation Fellowship.

ACKNOWLEDGEMENTS

The authors would like to acknowledge Dr. Lin Zhao for his technical support.

REFERENCES

- 1
2
3
4
5 Abate, N. (2012). Adipocyte maturation arrest: a determinant of systemic insulin resistance to
6 glucose disposal. *J Clin Endocrinol Metab* **97**(3), 760-3.
7
8
9 Ahmadian, M., Duncan, R. E., Varady, K. A., Frasson, D., Hellerstein, M. K., Birkenfeld, A. L.,
10 Samuel, V. T., Shulman, G. I., Wang, Y., Kang, C., *et al.* (2009). Adipose overexpression of
11 desnutrin promotes fatty acid use and attenuates diet-induced obesity. *Diabetes* **58**(4), 855-66.
12
13 Aquilano, K., Baldelli, S., La Barbera, L., Lettieri Barbato, D., Tatulli, G., and Ciriolo, M. R.
14 (2016). Adipose triglyceride lipase decrement affects skeletal muscle homeostasis during aging
15 through FAs-PPARalpha-PGC-1alpha antioxidant response. *Oncotarget* **7**(17), 23019-32.
16
17
18 Atashi, F., Modarressi, A., and Pepper, M. S. (2015). The role of reactive oxygen species in
19 mesenchymal stem cell adipogenic and osteogenic differentiation: a review. *Stem Cells Dev*
20 **24**(10), 1150-63.
21
22
23 Bansal, S., Buring, J. E., Rifai, N., Mora, S., Sacks, F. M., and Ridker, P. M. (2007). Fasting
24 compared with nonfasting triglycerides and risk of cardiovascular events in women. *JAMA*
25 **298**(3), 309-16.
26
27 Barra, N. G., Lisyansky, M., Vanduzer, T. A., Raha, S., Holloway, A. C., and Hardy, D. B.
28 (2017). Maternal nicotine exposure leads to decreased cardiac protein disulfide isomerase and
29 impaired mitochondrial function in male rat offspring. *J Appl Toxicol* doi: 10.1002/jat.3503.
30
31
32 Bezaire, V., Mairal, A., Ribet, C., Lefort, C., Girusse, A., Jocken, J., Laurencikiene, J., Anesia,
33 R., Rodriguez, A. M., Ryden, M., *et al.* (2009). Contribution of adipose triglyceride lipase and
34 hormone-sensitive lipase to lipolysis in hMADS adipocytes. *J Biol Chem* **284**(27), 18282-91.
35
36
37 Bolsoni-Lopes, A., and Alonso-Vale, M. I. (2015). Lipolysis and lipases in white adipose tissue -
38 An update. *Arch Endocrinol Metab* **59**(4), 335-42.
39
40 Bruin, J. E., Petre, M. A., Lehman, M. A., Raha, S., Gerstein, H. C., Morrison, K. M., and
41 Holloway, A. C. (2008). Maternal nicotine exposure increases oxidative stress in the offspring.
42 *Free Radic Biol Med* **44**(11), 1919-25.
43
44
45 Chen, W., Jiang, Y., Han, J., Hu, J., He, T., Yan, T., Huang, N., Zhang, Q., Mei, H., Liao, Y., *et*
46 *al.* (2017a). Atgl deficiency induces podocyte apoptosis and leads to glomerular filtration barrier
47 damage. *FEBS J* **284**(7), 1070-1081.
48
49
50 Chen, W., Zhang, Q., Cheng, S., Huang, J., Diao, G., and Han, J. (2017b). Atgl gene deletion
51 predisposes to proximal tubule damage by impairing the fatty acid metabolism. *Biochem Biophys*
52 *Res Commun* **487**(1), 160-166.
53
54
55
56
57
58
59
60

1
2
3 Chi, W., Dao, D., Lau, T. C., Henriksbo, B. D., Cavallari, J. F., Foley, K. P., and Schertzer, J. D.
4 (2014). Bacterial peptidoglycan stimulates adipocyte lipolysis via NOD1. *PLoS One* **9**(5),
5 e97675.

6
7
8 Chou, H. C., and Chen, C. M. (2014). Maternal nicotine exposure during gestation and lactation
9 induces cardiac remodeling in rat offspring. *Reprod Toxicol* **50**, 4-10.

10
11 Cui, Y., Shooshtari, S., Forget, E. L., Clara, I., and Cheung, K. F. (2014). Smoking during
12 pregnancy: findings from the 2009-2010 Canadian Community Health Survey. *PLoS One* **9**(1),
13 e84640.

14
15
16 Cupul-Uicab, L. A., Skjaerven, R., Haug, K., Travlos, G. S., Wilson, R. E., Eggesbo, M.,
17 Hoppin, J. A., Whitworth, K. W., and Longnecker, M. P. (2012). Exposure to tobacco smoke in
18 utero and subsequent plasma lipids, ApoB, and CRP among adult women in the MoBa cohort.
19 *Environ Health Perspect* **120**(11), 1532-7.

20
21
22 Dasgupta, C., Xiao, D., Xu, Z., Yang, S., and Zhang, L. (2012). Developmental nicotine
23 exposure results in programming of alveolar simplification and interstitial pulmonary fibrosis in
24 adult male rats. *Reprod Toxicol* **34**(3), 370-7.

25
26
27 Fan, J., Zhang, W. X., Rao, Y. S., Xue, J. L., Wang, F. F., Zhang, L., and Yan, Y. E. (2016).
28 Perinatal Nicotine Exposure Increases Obesity Susceptibility in Adult Male Rat Offspring by
29 Altering Early Adipogenesis. *Endocrinology* **157**(11), 4276-4286.

30
31
32 Fox, K. A., Longo, M., Tamayo, E., Gamble, P., Makhlof, M., Mateus, J. F., and Saade, G. R.
33 (2012). Sex-specific effects of nicotine exposure on developmental programming of blood
34 pressure and vascular reactivity in the C57Bl/6J mouse. *Am J Obstet Gynecol* **207**(3), 208 e1-9.

35
36
37 Gao, Y. J., Holloway, A. C., Su, L. Y., Takemori, K., Lu, C., and Lee, R. M. (2008). Effects of
38 fetal and neonatal exposure to nicotine on blood pressure and perivascular adipose tissue
39 function in adult life. *Eur J Pharmacol* **590**(1-3), 264-8.

40
41
42 Gao, Y. J., Holloway, A. C., Zeng, Z. H., Lim, G. E., Petrik, J. J., Foster, W. G., and Lee, R. M.
43 (2005). Prenatal exposure to nicotine causes postnatal obesity and altered perivascular adipose
44 tissue function. *Obes Res* **13**(4), 687-92.

45
46
47 Gochberg-Sarver, A., Kedmi, M., Gana-Weisz, M., Bar-Shira, A., and Orr-Urtreger, A. (2012).
48 Tnfalpha, Cox2 and AdipoQ adipokine gene expression levels are modulated in murine adipose
49 tissues by both nicotine and nACh receptors containing the beta2 subunit. *Mol Genet Metab*
50 **107**(3), 561-70.

51
52
53 Greenwood, M. R., and Hirsch, J. (1974). Postnatal development of adipocyte cellularity in the
54 normal rat. *J Lipid Res* **15**(5), 474-83.

55
56
57 Holloway, A. C., Kellenberger, L. D., and Petrik, J. J. (2006). Fetal and neonatal exposure to
58 nicotine disrupts ovarian function and fertility in adult female rats. *Endocrine* **30**(2), 213-6.
59
60

1
2
3 Holloway, A. C., Petrik, J. J., Bruin, J. E., and Gerstein, H. C. (2008). Rosiglitazone prevents
4 diabetes by increasing beta-cell mass in an animal model of type 2 diabetes characterized by
5 reduced beta-cell mass at birth. *Diabetes Obes Metab* **10**(9), 763-71.
6

7
8 Hoogduijn, M. J., Cheng, A., and Genever, P. G. (2009). Functional nicotinic and muscarinic
9 receptors on mesenchymal stem cells. *Stem Cells Dev* **18**(1), 103-12.
10

11 Hu, Y., Rosen, D. G., Zhou, Y., Feng, L., Yang, G., Liu, J., and Huang, P. (2005). Mitochondrial
12 manganese-superoxide dismutase expression in ovarian cancer: role in cell proliferation and
13 response to oxidative stress. *J Biol Chem* **280**(47), 39485-92.
14

15
16 Ji, C., Chen, X., Gao, C., Jiao, L., Wang, J., Xu, G., Fu, H., Guo, X., and Zhao, Y. (2011). IL-6
17 induces lipolysis and mitochondrial dysfunction, but does not affect insulin-mediated glucose
18 transport in 3T3-L1 adipocytes. *J Bioenerg Biomembr* **43**(4), 367-75.
19

20
21 Kershaw, E. E., Hamm, J. K., Verhagen, L. A., Peroni, O., Katic, M., and Flier, J. S. (2006).
22 Adipose triglyceride lipase: function, regulation by insulin, and comparison with adiponutrin.
23 *Diabetes* **55**(1), 148-57.
24

25
26 Kersten, S. (2014). Physiological regulation of lipoprotein lipase. *Biochim Biophys Acta* **1841**(7),
27 919-33.
28

29
30 Lawrence, T. (2009). The nuclear factor NF-kappaB pathway in inflammation. *Cold Spring Harb*
Perspect Biol **1**(6), a001651.
31

32
33 Li, D., Zhang, Y., Liu, Y., Sun, R., and Xia, M. (2015). Purified anthocyanin supplementation
34 reduces dyslipidemia, enhances antioxidant capacity, and prevents insulin resistance in diabetic
35 patients. *J Nutr* **145**(4), 742-8.
36

37
38 Li, N., and Karin, M. (1999). Is NF-kappaB the sensor of oxidative stress? *FASEB J* **13**(10),
39 1137-43.
40

41
42 Lin, F. T., and Lane, M. D. (1994). CCAAT/enhancer binding protein alpha is sufficient to
43 initiate the 3T3-L1 adipocyte differentiation program. *Proc Natl Acad Sci U S A* **91**(19), 8757-
44 61.
45

46
47 Lin, F. T., MacDougald, O. A., Diehl, A. M., and Lane, M. D. (1993). A 30-kDa alternative
48 translation product of the CCAAT/enhancer binding protein alpha message: transcriptional
49 activator lacking antimitotic activity. *Proc Natl Acad Sci U S A* **90**(20), 9606-10.
50

51
52 Liu, M., Chuang Key, C. C., Weckerle, A., Boudyguina, E., Sawyer, J. K., Gebre, A. K., Spoo,
53 W., Makwana, O., and Parks, J. S. (2018). Feeding of tobacco blend or nicotine induced weight
54 loss associated with decreased adipocyte size and increased physical activity in male mice. *Food*
55 *Chem Toxicol* **113**, 287-295.
56
57
58
59
60

1
2
3 Luck, W., and Nau, H. (1987). Nicotine and cotinine concentrations in the milk of smoking
4 mothers: influence of cigarette consumption and diurnal variation. *Eur J Pediatr* **146**(1), 21-6.
5

6 Luck, W., Nau, H., Hansen, R., and Steldinger, R. (1985). Extent of nicotine and cotinine
7 transfer to the human fetus, placenta and amniotic fluid of smoking mothers. *Dev Pharmacol*
8 *Ther* **8**(6), 384-95.
9

10
11 Ma, N., Nicholson, C. J., Wong, M., Holloway, A. C., and Hardy, D. B. (2014). Fetal and
12 neonatal exposure to nicotine leads to augmented hepatic and circulating triglycerides in adult
13 male offspring due to increased expression of fatty acid synthase. *Toxicol Appl Pharmacol*
14 **275**(1), 1-11.
15

16
17 Macdonald, T. L., Wan, Z., Frenedo-Cumbo, S., Dyck, D. J., and Wright, D. C. (2013). IL-6 and
18 epinephrine have divergent fiber type effects on intramuscular lipolysis. *J Appl Physiol (1985)*
19 **115**(10), 1457-63.
20

21
22 Mohsenzadeh, Y., Rahmani, A., Cheraghi, J., Pyrani, M., and Asadollahi, K. (2014). Prenatal
23 exposure to nicotine in pregnant rat increased inflammatory marker in newborn rat. *Mediators*
24 *Inflamm* **2014**, 274048.
25

26
27 Morgan, M. J., and Liu, Z. G. (2011). Crosstalk of reactive oxygen species and NF-kappaB
28 signaling. *Cell Res* **21**(1), 103-15.
29

30
31 Oncken, C. A., and Kranzler, H. R. (2003). Pharmacotherapies to enhance smoking cessation
32 during pregnancy. *Drug Alcohol Rev* **22**(2), 191-202.

33
34 Orellana, J. A., Busso, D., Ramirez, G., Campos, M., Rigotti, A., Eugenin, J., and von Bernhardt,
35 R. (2014). Prenatal nicotine exposure enhances Cx43 and Panx1 unopposed channel activity in
36 brain cells of adult offspring mice fed a high-fat/cholesterol diet. *Front Cell Neurosci* **8**, 403.
37

38
39 Ossipow, V., Descombes, P., and Schibler, U. (1993). CCAAT/enhancer-binding protein mRNA
40 is translated into multiple proteins with different transcription activation potentials. *Proc Natl*
41 *Acad Sci U S A* **90**(17), 8219-23.
42

43
44 Pauly, J. R., and Slotkin, T. A. (2008). Maternal tobacco smoking, nicotine replacement and
45 neurobehavioural development. *Acta Paediatr* **97**(10), 1331-7.

46
47 Poissonnet, C. M., Burdi, A. R., and Garn, S. M. (1984). The chronology of adipose tissue
48 appearance and distribution in the human fetus. *Early Hum Dev* **10**(1-2), 1-11.
49

50
51 Roman-Galvez, R. M., Amezcua-Prieto, C., Olmedo-Requena, R., Lewis-Mikhael Saad, A. M.,
52 Martinez-Galiano, J. M., and Bueno-Cavanillas, A. (2017). Partner smoking influences whether
53 mothers quit smoking during pregnancy: A prospective cohort study. *BJOG* doi: 10.1111/1471-
54 0528.14986.
55
56
57
58
59
60

- 1
2
3 Ryden, M., Dicker, A., van Harmelen, V., Hauner, H., Brunnberg, M., Perbeck, L., Lonnqvist,
4 F., and Arner, P. (2002). Mapping of early signaling events in tumor necrosis factor-alpha -
5 mediated lipolysis in human fat cells. *J Biol Chem* **277**(2), 1085-91.
6
7
8 Sbrana, E., Suter, M. A., Abramovici, A. R., Hawkins, H. K., Moss, J. E., Patterson, L., Shope,
9 C., and Aagaard-Tillery, K. (2011). Maternal tobacco use is associated with increased markers of
10 oxidative stress in the placenta. *Am J Obstet Gynecol* **205**(3), 246 e1-7.
11
12 Sena, L. A., and Chandel, N. S. (2012). Physiological roles of mitochondrial reactive oxygen
13 species. *Mol Cell* **48**(2), 158-67.
14
15
16 Shahab, L., Dobbie, F., Hiscock, R., McNeill, A., and Bauld, L. (2016). Prevalence and Impact
17 of Long-term Use of Nicotine Replacement Therapy in UK Stop-Smoking Services: Findings
18 From the ELONS Study. *Nicotine Tob Res* doi: 10.1093/ntr/ntw258.
19
20
21 Somm, E., Schwitzgebel, V. M., Vauthay, D. M., Camm, E. J., Chen, C. Y., Giacobino, J. P.,
22 Sizonenko, S. V., Aubert, M. L., and Huppi, P. S. (2008). Prenatal nicotine exposure alters early
23 pancreatic islet and adipose tissue development with consequences on the control of body weight
24 and glucose metabolism later in life. *Endocrinology* **149**(12), 6289-99.
25
26
27 Spalding, K. L., Arner, E., Westermark, P. O., Bernard, S., Buchholz, B. A., Bergmann, O.,
28 Blomqvist, L., Hoffstedt, J., Naslund, E., Britton, T., *et al.* (2008). Dynamics of fat cell turnover
29 in humans. *Nature* **453**(7196), 783-7.
30
31
32 Talirevic, E., and Jelena, S. (2012). Quercetin in the treatment of dyslipidemia. *Med Arh* **66**(2),
33 87-8.
34
35
36 Tappin, D. M., MacAskill, S., Bauld, L., Eadie, D., Shipton, D., and Galbraith, L. (2010).
37 Smoking prevalence and smoking cessation services for pregnant women in Scotland. *Subst*
38 *Abuse Treat Prev Policy* **5**, 1.
39
40
41 Tong, V. T., Dietz, P. M., Morrow, B., D'Angelo, D. V., Farr, S. L., Rockhill, K. M., England, L.
42 J., Centers for Disease, C., and Prevention (2013). Trends in smoking before, during, and after
43 pregnancy--Pregnancy Risk Assessment Monitoring System, United States, 40 sites, 2000-2010.
44 *MMWR Surveill Summ* **62**(6), 1-19.
45
46
47 Villa, A., Della Torre, S., Stell, A., Cook, J., Brown, M., and Maggi, A. (2012). Tetradian
48 oscillation of estrogen receptor alpha is necessary to prevent liver lipid deposition. *Proc Natl*
49 *Acad Sci U S A* **109**(29), 11806-11.
50
51
52 Vogel, C., and Marcotte, E. M. (2012). Insights into the regulation of protein abundance from
53 proteomic and transcriptomic analyses. *Nat Rev Genet* **13**(4), 227-32.
54
55
56 Wen, X., Triche, E. W., Hogan, J. W., Shenassa, E. D., and Buka, S. L. (2010). Birth weight and
57 adult hypercholesterolemia: subgroups of small-for-gestational-age based on maternal smoking
58 status during pregnancy. *Epidemiology* **21**(6), 786-90.
59
60

1
2
3 Weng, S. F., Redsell, S. A., Swift, J. A., Yang, M., and Glazebrook, C. P. (2012). Systematic
4 review and meta-analyses of risk factors for childhood overweight identifiable during infancy.
5 *Arch Dis Child* **97**(12), 1019-26.
6

7
8 Yang, X., Zhang, X., Heckmann, B. L., Lu, X., and Liu, J. (2011). Relative contribution of
9 adipose triglyceride lipase and hormone-sensitive lipase to tumor necrosis factor-alpha (TNF-
10 alpha)-induced lipolysis in adipocytes. *J Biol Chem* **286**(47), 40477-85.
11

12 Zarrouki, B., Soares, A. F., Guichardant, M., Lagarde, M., and Geloën, A. (2007). The lipid
13 peroxidation end-product 4-HNE induces COX-2 expression through p38MAPK activation in
14 3T3-L1 adipose cell. *FEBS Lett* **581**(13), 2394-400.
15

16
17 Zechner, R., Kienesberger, P. C., Haemmerle, G., Zimmermann, R., and Lass, A. (2009).
18 Adipose triglyceride lipase and the lipolytic catabolism of cellular fat stores. *J Lipid Res* **50**(1),
19 3-21.
20

21 Zhang, H. H., Halbleib, M., Ahmad, F., Manganiello, V. C., and Greenberg, A. S. (2002). Tumor
22 necrosis factor-alpha stimulates lipolysis in differentiated human adipocytes through activation
23 of extracellular signal-related kinase and elevation of intracellular cAMP. *Diabetes* **51**(10),
24 2929-35.
25

26
27 Zhang, X., Wang, Z., Li, J., Gu, D., Li, S., Shen, C., and Song, Z. (2013). Increased 4-
28 hydroxynonenal formation contributes to obesity-related lipolytic activation in adipocytes. *PLoS*
29 *One* **8**(8), e70663.
30

31
32 Zhang, Y., Li, J., and Wen, X. (2017). Jueming prescription and its ingredients, semen cassiae
33 and Rhizoma Curcumae Longae, stimulate lipolysis and enhance the phosphorylation of
34 hormonesensitive lipase in cultured rat white adipose tissue. *Mol Med Rep* **16**(5), 6200-6207.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FIGURE LEGENDS

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1: Maternal nicotine exposure leads to decreased visceral adipocyte cell area in 6 month old offspring. Representative cross-sections of gonadal WAT excised from (A) vehicle and *in utero* nicotine exposed 6 month old male and female offspring were stained with hematoxylin and eosin. (B) Bar graph representing calculated cell areas analyzed using Northern Eclipse software (n=4/group/sex). (C) Transcript and (D) protein levels of AKT-1 in six month vehicle and perinatal nicotine exposed WAT were determined via real time PCR and Western blot, respectively. mRNA levels were expressed as means normalized to β -actin \pm SEM (n=15/vehicle; n=11/nicotine). Protein levels were expressed as means normalized to β -actin \pm SEM (n=8/vehicle; n=7/nicotine). **,*** Significant difference (**p<0.01, ***p < 0.001).

Figure 2: *In utero* nicotine exposure increases the expression of the differentiation marker C/EBP α in WAT of 6 month old female offspring. Transcript and protein levels of targets of interest in 3 week and six month vehicle and perinatal nicotine exposed WAT were determined via real time PCR and Western blot, respectively. Six month (A) C/EBP α mRNA and (B) protein levels. (C) C/EBP β mRNA and (D) protein levels. (E) SREBP1-c mRNA and (F) protein levels. Three week (G) C/EBP α mRNA and (H) protein levels. mRNA levels were expressed as means normalized to β -actin \pm SEM (n=15/vehicle; n=11/nicotine). Protein levels were expressed as means normalized to β -actin \pm SEM (n=8/vehicle; n=7/nicotine). *, Significant difference (*p<0.05).

Figure 3: Maternal nicotine exposure does not increase the expression of markers involved in lipogenesis and fatty acid transporters in WAT of 6 month old female offspring.

1
2
3 Transcript levels of targets of interest in six month vehicle and perinatal nicotine exposed WAT
4 were determined via real time PCR. Six month (A) LPL, (B) ACC α , (C) FAS, (D) ASCL1, (E)
5 FATP1, and (F) FATP4 mRNA levels. mRNA levels were expressed as means normalized to β -
6 actin \pm SEM (n=15/vehicle; n=11/nicotine).
7
8
9
10
11
12
13

14 **Figure 4: *In utero* nicotine exposed female offspring exhibit increased expression of ATGL**
15 **at 6 months, but not at 3 weeks.** Transcript and protein levels of ATGL in 3 week and six
16 month vehicle and perinatal nicotine exposed WAT were determined via real time PCR and
17 Western blot, respectively. Six month ATGL (A) mRNA and (B) protein levels. Three week
18 ATGL (C) mRNA and (D) protein levels. mRNA levels were expressed as means normalized to
19 β -actin \pm SEM (n=15/vehicle; n=11/nicotine). Protein levels were expressed as means
20 normalized to β -actin \pm SEM (n=8/vehicle; n=8/nicotine). *,** Significant difference (*p<0.05,
21 **p<0.01).
22
23
24
25
26
27
28
29
30
31
32
33
34

35 **Figure 5: *In utero* nicotine exposed female offspring exhibit increased anti-oxidant**
36 **expression in WAT at 3 weeks and 6 months.** Transcript and protein levels of SOD-1 and
37 SOD-2 in 3 week and six month vehicle and perinatal nicotine exposed WAT were determined
38 via real time PCR and Western blot, respectively. Six month SOD-1 (A) mRNA and (B) protein
39 levels and SOD-2 (C) mRNA and (D) protein levels. Three week SOD-1 (E) mRNA and (F)
40 protein levels and SOD-2 (G) mRNA and (H) protein levels. Protein levels of 4-HNE at (I) 6
41 months and (J) 3 weeks of age. mRNA levels were expressed as means normalized to β -actin \pm
42 SEM (n=15/vehicle; n=11/nicotine). Protein levels were expressed as means normalized to β -
43 actin \pm SEM (n=8/vehicle; n=8/nicotine). *,** Significant difference (*p<0.05, **p<0.01).
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8 **Figure 6: Maternal nicotine exposed female offspring exhibit increases in WAT**
9 **inflammation at 6 months, but not at 3 weeks.** Transcript levels of targets of interest in six
10 month vehicle and perinatal nicotine exposed WAT were determined via real time PCR. Six
11 month vehicle and perinatal nicotine exposed WAT were determined via real time PCR. Six
12 month vehicle and perinatal nicotine exposed WAT were determined via real time PCR. Six
13 month vehicle and perinatal nicotine exposed WAT were determined via real time PCR. Six
14 month (A) IL-1 β , (B) TNF α , (C) IL-6, and (D) CD68 mRNA levels. Three week (E) IL-1 β , (F)
15 TNF α , (G) IL-6, and (H) CD68 mRNA levels. mRNA levels were expressed as means
16 normalized to β -actin \pm SEM (n=15/vehicle; n=11/nicotine). *,**,*** Significant difference
17 (*p<0.05, **p<0.01, ***p<0.001).
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TABLES

Table 1. Forward and reverse primer sequences used for quantitative Real-Time PCR

GENE	Forward	Reverse	GenBank/Reference
AKT-1	ATGTGTATGAGAAGAAGC TGAGCC	GTTCACTGTCCACACACT CCA	NM_033230.2
C/EBP α	GCCGGGAGAACTCTAACT CC	TCGATGTAGGCGCTGATG TC	NM_00128757 7.1
C/EBP β	ACCACGACTTCCTTTCCG AC	TAACCGTAGTCGGACGGC TT	NM_024125.5
SREBP1 c	CATGGACGAGCTACCCTT CG	TCTTCGATGTCGGTCAAG AGC	NM_00127670 7.1
LPL	GGATGCAACATTGGAGAA CCC	GCTGGGGTTTTCTTCATTC AGC	NM_012598.2
ACC α	TCCGTATGTGACCAAAGA CC	TACGTTGTTCCCAAGGAC TG	NM_022193.1
FAS	GGACATGGTCACAGACGA TGAC	CGTCGAACTTGGACAGAT CCTT	NM_017332.1
ACSL1	CTACAGGCAACCCCAAAG GA	AATGCACTCTCCGTCGCT T	NM_012820.1
FATP1	CAGCCTCTGTGGCCTCAT T	ACCCACGTACACACCGAA C	NM_053580.2
FATP4	CTTGGGCAACTTTGACAG CC	AGGACAGGATGCGGCTAT TG	NM_00110070 6.1
ATGL	AACGCCACTCACATCTAC GG	TACCAGGTTGAAGGAGGG GT	NM_00110850 9.2
EGR-1	CGAGCGAACAACCCTACG A	CGATGTCAGAAAAGGACT CTGTG	NM_012551.2
FOXO-1	TGCAGCAGACACCTTGCT AT	TTGGGGCTGGGGGAATTT AG	NM_00119184 6.2
IL-1 β	CAGCTTTCGACAGTGAGG AGA	GTCGAGATGCTGCTGTGA GA	NM_031512.2
TNF α	CCGGGCAGGTCTACTTTG GA	AGGCCACTACTTCAGCGT CTCG	NM_012675.3
IL-6	CTTCCAGCCAGTTGCCTT CTTG	TGGTCTGTTGTGGGTGGT ATCC	NM_012589.2
CD68	ACTGGGGCTCTTGGAAC TACAC	CCTTGGTTTTGTTTCGGGTT CA	NM_00103163 8.1
SOD-1	ATTGGCCGTACTATGGTG GTC	GCAATCCCAATCACACCA CA	NM_017050.1
SOD-2	ATTGCCGCCTGCTCTAAT CA	TCCACACATCAATCCCC AG	NM_017051.2

Table 2. Western Blot antibodies, dilutions used in experiments, and company and catalogue information.

<i>Antibody name</i>	<i>Source</i>	<i>Dilution</i>	<i>Company (#Catalogue)</i>
4-HNE	Mouse monoclonal	1:500	R&D Systems Minneapolis, MN, USA (#MAB3249)
AKT-1	Rabbit polyclonal	1:500	Abcam Inc., Cambridge, MA, USA (#abcam 5919)
ATGL (H-144)	Rabbit polyclonal	1:1000	Santa Cruz Biotechnology Inc., Santa Cruz, CA, USA (#sc-67355)
C/EBP α (14AA)	Rabbit polyclonal	1:500	Santa Cruz Biotechnology Inc., Santa Cruz, CA, USA (#sc-61)
C/EBP β (C-19)	Rabbit polyclonal	1:300	Santa Cruz Biotechnology Inc., Santa Cruz, CA, USA (#sc-150)
SOD-1 (FL-154)	Rabbit polyclonal	1:1000	Santa Cruz Biotechnology Inc., Santa Cruz, CA, USA (#sc-11407)
SOD-2 (FL-222)	Rabbit polyclonal	1:1000	Santa Cruz Biotechnology Inc., Santa Cruz, CA, USA (#sc-30080)
SREBP 1c (H-160)	Rabbit polyclonal	1:500	Santa Cruz Biotechnology Inc., Santa Cruz, CA, USA (#sc-8984)
Mouse IgG (H+L) Secondary	Sheep	1:5000	GE Healthcare UK, Pittsburgh, PA, USA (#NA931)
Rabbit IgG (H+L) Secondary	Donkey	1:10000	Jackson ImmunoResearch Laboratories, West Grove, PA, USA (#711-001-003)




Figure 1

124x188mm (300 x 300 DPI)




Figure 2

124x201mm (300 x 300 DPI)




Figure 3

124x167mm (300 x 300 DPI)




Figure 4

124x141mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 5

112x202mm (300 x 300 DPI)




Figure 6

124x184mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60