

Western University

Scholarship@Western

Physiology and Pharmacology Publications

Physiology and Pharmacology Department

4-20-2020

Maternal exposure to Δ^9 -tetrahydrocannabinol impairs female offspring glucose homeostasis and endocrine pancreatic development in the rat.

Ryan Gillies

Kendrick Lee

Sebastian Vanin

Steven R Laviolette

Alison C Holloway

See next page for additional authors

Follow this and additional works at: <https://ir.lib.uwo.ca/physpharmpub>



Part of the [Medical Physiology Commons](#), and the [Pharmacy and Pharmaceutical Sciences Commons](#)

Citation of this paper:

Gillies, Ryan; Lee, Kendrick; Vanin, Sebastian; Laviolette, Steven R; Holloway, Alison C; Arany, Edith; and Hardy, Daniel B, "Maternal exposure to Δ^9 -tetrahydrocannabinol impairs female offspring glucose homeostasis and endocrine pancreatic development in the rat." (2020). *Physiology and Pharmacology Publications*. 234.

<https://ir.lib.uwo.ca/physpharmpub/234>

Authors

Ryan Gillies, Kendrick Lee, Sebastian Vanin, Steven R Laviolette, Alison C Holloway, Edith Arany, and Daniel B Hardy



Maternal exposure to $\Delta 9$ -tetrahydrocannabinol impairs female offspring glucose homeostasis and endocrine pancreatic development in the rat

Ryan Gillies^{a,e}, Kendrick Lee^{c,e}, Sebastian Vanin^{c,e}, Steven R. Laviolette^{d,e}, Alison C. Holloway^g, Edith Arany^{a,b,e,f}, Daniel B. Hardy^{c,def,*}

^a Department of Pathology and Laboratory Medicine, London, Ontario, Canada

^b Department of Medicine, London, Ontario, Canada

^c Department of Physiology and Pharmacology, London, Ontario, Canada

^d Department of Anatomy and Cell Biology, London, Ontario, Canada

^e Western University, London, Ontario, Canada

^f Children's Health Research Institute, Lawson Health Research Institute, St. Joseph's Health Care, London, Ontario, Canada

^g Department of Obstetrics and Gynecology, McMaster University, Hamilton, Ontario, Canada

ARTICLE INFO

Keywords:

Endocannabinoid system
Fetal growth restriction
Sexual dimorphism
Glucose intolerance
Islet
 β -cell

ABSTRACT

Recent reports indicate that 7% of pregnant mothers in North America use *cannabis*. This is concerning given that *in utero* exposure to $\Delta 9$ -tetrahydrocannabinol ($\Delta 9$ -THC), the main psychoactive component in *cannabis*, causes fetal growth restriction and may alter replication and survival of pancreatic β -cells in the offspring. Accordingly, we hypothesized that maternal exposure to $\Delta 9$ -THC during pregnancy will impair postnatal glucometabolic health of offspring. To test this hypothesis, pregnant Wistar rats were treated with daily intraperitoneal injections of either 3 mg/kg $\Delta 9$ -THC or vehicle from gestational day 6 to birth. Offspring were subsequently challenged with glucose and insulin at 5 months of age to assess glucose tolerance and peripheral muscle insulin sensitivity. Female offspring exposed to $\Delta 9$ -THC *in utero* were glucose intolerant, associated with blunted insulin response in muscle and increased serum insulin concentration 15 min after glucose challenge. Additionally, pancreata from male and female offspring were harvested at postnatal day 21 and 5 months of age for assessment of endocrine pancreas morphometry by immunostaining. This analysis revealed that gestational exposure to $\Delta 9$ -THC reduced the density of islets in female, but not male, offspring at postnatal day 21 and 5 months, culminating in reduced β -cell mass at 5 months. These results demonstrate that fetal exposure to $\Delta 9$ -THC causes female-specific impairments in glucose homeostasis, raising concern regarding the metabolic health of offspring, particularly females, exposed to *cannabis in utero*.

1. Introduction

Recent reports indicate that 7% of pregnant mothers in North America, and 19 % of those aged 18–24, use *cannabis* [1]. Much of this use may be motivated by the belief that *cannabis* eases nausea, vomiting, and lost appetite during pregnancy [2], in conjunction with media portrayals of *cannabis* as a natural alternative to pharmaceuticals [3,4]. A recent integrative review also concluded that women who used *cannabis* during pregnancy did so given the perception that there were no significant risks to the mother or fetus [5]. However, little is known regarding the effects of *cannabis* and its constituent compounds on postnatal outcomes.

We and others have recently demonstrated in rodents that $\Delta 9$ -tetrahydrocannabinol ($\Delta 9$ -THC), the main psychoactive component of *cannabis*, leads to fetal growth restriction, in part, due to placental insufficiency [6,7]. This is pertinent because since 1995, selective breeding has increased $\Delta 9$ -THC content in *cannabis* from 4% to 12 % [8]. Given the strong association between im-

paired fetal growth and the development of type 2 diabetes [9], this raises concern regarding the metabolic health of offspring exposed to $\Delta 9$ -THC *in utero*. In addition, $\Delta 9$ -THC may directly influence development of the fetal pancreas, through its interaction with the cannabinoid 1 receptor (CB1R) [10–12].

CB1R is expressed by both insulin-producing β -cells and glucagon-producing α -cells [13], and contributes to islet formation and organization during fetal development [10]. Moreover, CB1R activation suppresses mitogenic signaling in β -cells, while also promoting apoptosis [11,12]. In a healthy fetus, these effects of CB1R are modulated by temporally-controlled endogenous endocannabinoid levels in fetal tissue and circulation [10,14,15]. However, studies indicate that $\Delta 9$ -THC readily traverses the placental barrier and 10–28 % of maternal concentrations are detected in the fetal plasma, with 2–5X higher concentrations in fetal tissues [16–18]. This would allow maternal $\Delta 9$ -THC to potentially dysregulate physiologic control of CB1R and its downstream processes within the developing fetal pancreas. Given that the long-term effects of $\Delta 9$ -THC on metabolic organs outside of the brain remain elusive, we inves-

Abbreviations: $\Delta 9$ -THC, $\Delta 9$ -tetrahydrocannabinol; CB1R, Cannabinoid 1 receptor; GD, gestational day; PND, postnatal day; IP-GTT, intraperitoneal glucose tolerance test.

* Corresponding author at: The Department of Physiology and Pharmacology, The University of Western Ontario, London, Ontario N6A 5C1, Canada.

E-mail address: Daniel.Hardy@schulich.uwo.ca (D.B. Hardy)

tigated the impact of *in utero* exposure to Δ^9 -THC on the development of the endocrine pancreas, including its impact on whole-body glucose metabolism in the offspring.

2. Research design and methods

2.1. Animals

All procedures were performed in accordance with the guidelines of the Canadian Council of Animal Care. Female Wistar rats were purchased from Charles River (La Salle) and throughout the experimental procedure were maintained at 22 °C on a 12:12-h light-dark cycle with access to food and water *ad libitum*. Dams were randomly assigned to receive daily intraperitoneal (*i.p.*) injections of either 3 mg/kg Δ^9 -THC (Sigma-Aldrich, Saint Louis) or vehicle (18:1 saline:cremophor) from gestational day (GD) 6 to GD22. This dosage of Δ^9 -THC *i.p.* yields maternal blood concentrations of Δ^9 -THC (8.6–12.4 ng/mL) comparable to those seen after moderate recreational *cannabis* smoking in human adults (13–63 ng/mL), as well as in aborted fetal tissue (4–287 ng/mL) after maternal *cannabis* use [19–21]. Administration of Δ^9 -THC was delayed until GD6 since administration of the drug earlier in pregnancy can induce spontaneous abortions [22]. We previously demonstrated that 3 mg/kg Δ^9 -THC given from GD6 to GD22 does not cause fetal demise, alterations in litter size, gestational length, or maternal weight gain [7]. Dams were allowed to deliver normally, and at birth (post-natal day 1; PND1), pups were weighed, and litters randomly culled to 8 offspring. Measurements at PND1 represent mixed males and females. Offspring were killed at either PND21 or 5 months of age (5 m) by *i.p.* pentobarbital overdose (100 mg/kg). 5 m was chosen as the age for sacrifice, as this is beyond the point (130 days-of-age) when rats have been demonstrated to exhibit sexual dimorphism in glucose intolerance and β -cell mass [23]. All offspring were fasted for ~14 h prior to sacrifice, then weighed, the pancreas dissected, fixed in 10 % formalin, and embedded in paraffin. Prior to sacrifice at 5 m, glycaemia was measured with a OneTouch Ultra2 hand-held glucometer (LifeScan, Zug, Switzerland) from blood obtained by tail vein snip. Blood was also drawn by cardiac puncture for quantification of serum insulin by ELISA (ALPCO, Salem, NH, USA), as well as serum estrogen (Biovision Milpitas, CA, USA), and free testosterone (R&D Systems, Minneapolis, MN, USA). Fasting glucose (FG) and insulin (FI) levels were used to calculate HOMA1-IR by the formula $\text{HOMA1-IR} = \text{FG} \times \text{FI} / 22.5$ as a measure of peripheral insulin resistance [24].

2.2. Intraperitoneal glucose tolerance test (IP-GTT)

Glucose homeostasis was investigated at 5 m using an intraperitoneal glucose tolerance test (IP-GTT). 5–6 offspring per sex per treatment were fasted overnight (~14 h) then allowed to acclimate in a dark, quiet room for 2 h. Baseline blood glucose levels were obtained as above, and 2 g/kg glucose administered by *i.p.* injection. Glucose levels were taken 5, 15, 30, 60, 90, and 120 min post injection. Total glucose response to the challenge was calculated using the trapezoidal area under the curve (AUC) for each animal. Blood was collected from the tail vein at 15 and 30 min after glucose bolus for quantification of serum insulin by ELISA (Crystal Chem, Elk Grove Village, IL, USA).

2.3. Intraperitoneal insulin challenge

To evaluate insulin sensitivity in muscle, phosphorylation of Akt was assessed in 11 female animals (6 vehicle and 5 Δ^9 -THC), each from a distinct dam, aged 5 m [25]. Rats were anaesthetized with isoflurane, the abdominal wall dissected, and the portal vein exposed and ligated distally. Saline was infused over 10 s and a portion of soleus muscle harvested. Insulin, 2 U/kg Humulin (Lilly, Indianapolis, IN, USA), was then infused over 10 s and a portion of soleus muscle was further harvested after 1 min. Tissue samples were snap frozen in liquid nitrogen and kept at -80 °C.

2.4. Protein extraction and western blot

Proteins were extracted from the muscle of rats subjected to intra-portal insulin challenge as described previously [23]. Proteins (50 μ g) were separated on an 8% SDS-PAGE, transferred to nitro-cellulose membranes, blocked for 60 min at room temperature in Tris-buffered saline containing 0.05 % Tween 20 (TTBS) and 5% non-fat dry milk or 5% BSA, then probed overnight at 48 °C with 1:1000 anti-phospho Akt (Ser 473) [pAkt (Ser 473)] mouse antibody or 1:1 000 anti-Akt2 rabbit antibody (Cell Signalling Technology, New England Biolabs, Beverly, MA, USA). Proteins were detected by enhanced chemiluminescence (Pierce, IL, USA) with horseradish peroxidase-labelled secondary antibodies (Sigma, St. Louis, MO, USA). The optical density of the bands was quantified with a Bio Imaging Gel System (Chemi Genius II, Syngene) with Gene Tools software. Each membrane was probed with an antibody against pAkt [Ser 473], then incubated in stripping buffer (Pierce, IL, USA) and re-probed with an antibody against Akt2 to normalize the results. Results are expressed as a ratio of these optical density values.

2.5. Immunohistochemistry and endocrine pancreas morphometry

Sections of 5 μ m were cut from paraffin-embedded pancreata of offspring harvested at PND21 and 5 m and mounted on SuperFrost Plus glass slides (Fischer Scientific, Toronto). Two or three sections per animal, separated by at least 50 μ m, were deparaffinized in xylene, rehydrated in descending ethanol series (100 %, 90 %, 70 %), then washed in tap water prior to application of rabbit anti-glucagon IgG (Novus Biologicals, Centennial, CO, USA) and mouse anti-insulin IgG (Sigma-Aldrich, Saint Louis, MO, USA) primary antibodies. ImmPRESS Duet Double Staining Kit (Vector Laboratories, Burlingame) was used to detect insulin- and glucagon-positive cells.

Analysis of pancreatic sections was performed as described previously [23]. Briefly, a composite image of the whole section was obtained using Microsoft Image Composite Editor and microphotographs taken with a 2.5x objective lens. Under 200x magnification, all individual islets were microphotographed and FIJI (<http://fiji.sc>) used to quantify the area occupied by β - and α -cells. β - and α -cell fractional area were calculated as the total insulin- and glucagon-positive area, respectively, divided by total pancreatic tissue area. The density of islets was calculated by counting the total number of islets present on each slide and dividing by total pancreatic tissue area. Mean islet size and mean β -cell area per islet were calculated by dividing total islet area (insulin- and glucagon-positive) or total β -cell area, respectively, by the total number of islets present on each slide. β - and α -cell mass were calculated as the product of pancreas weight and β - and α -cell fractional area, respectively. The density of extra-islet clusters (less than 500 μ m² or ~5 endocrine cells) was also assessed as a marker of endocrine cell neogenesis [26,27]. All calculations were averaged across 2–3 slides per animal.

To assess replication in β - and α -cells, 2–3 slides per animal, separated by at least 50 μ m, were incubated with mouse anti-Ki67 (Sigma-Aldrich, Saint Louis, MO, USA), guinea pig anti-insulin (Abcam, Toronto, ON, Canada), and rabbit anti-glucagon (Novus Biologicals, Centennial, CO, USA) primary antibodies. Donkey anti-mouse 488, anti-guinea pig 555, and anti-rabbit 647 fluorescent secondary antibodies (ThermoFisher, Toronto, ON, Canada) were then applied, and DAPI (ThermoFisher, Toronto, ON, Canada) used to counterstain nuclei. Apoptosis in β - and α -cells was examined similarly using ApopTag Fluorescein In Situ Apoptosis Detection Kit (Sigma-Aldrich, Saint Louis). At least 20 randomly-selected islets per section were microphotographed using a Nikon Eclipse Ts2R Epifluorescence Microscope. Manual tracing was used to outline insulin- and glucagon-positive areas, and particle analysis and water shedding filters applied to find and count DAPI⁺ nuclei within these regions [28]. Ki67⁺ or TUNEL⁺ nuclei were counted manually within each region and used to determine the percentage of Ki67⁺ or TUNEL⁺ β - and α -cells. An average of 1656 β -cells and 1201 α -cells were counted for every animal.

2.6. Statistical analysis

A sample size of 5–7 animals per sex per age group per treatment was chosen based on achieving a statistically significant difference with an expected standard deviation of 15 % or less, based on previous studies [27]. Values depicted are mean \pm SEM and considered significant if $p < 0.05$. Analysis of the data was performed using Excel v16.16 (Microsoft) and GraphPad Prism v7 (GraphPad Software, Inc). Unless indicated otherwise, statistical analysis was by two-way ANOVA followed by Holm-Sidak-corrected multiple comparisons between $\Delta 9$ -THC-exposed offspring and their sex-matched controls [23].

3. Results

3.1. Fetal growth restriction and postnatal catch-up growth

Offspring bodyweight and pancreatic weight were measured at birth, at weaning on PND21, and in young adulthood at 5 m. Gestational exposure to $\Delta 9$ -THC significantly reduced both birthweights (Fig. 1 A; 6.62 ± 0.11 g for $\Delta 9$ -THC and 7.02 ± 0.11 g for vehicle, $p = 0.0135$) and pancreatic weights (Fig. 1 D; 12.18 ± 1.54 mg for $\Delta 9$ -THC vs 17.33 ± 1.62 mg for vehicle, $p = 0.029$) in male and female offspring. By PND21, $\Delta 9$ -THC-exposed offspring recovered in terms of both bodyweight and pancreatic weight (Fig. 1 B, E), and thereafter maintained comparable weights to that of their sex-

matched controls (Fig. 1 C, F). Male did not differ from female offspring in bodyweight or pancreatic weight at PND21, therefore results are pooled.

3.2. Impaired glucose homeostasis

To determine the impact of *in utero* $\Delta 9$ -THC exposure on glucose metabolism in postnatal life, blood glucose levels were examined after ~14 h of fasting in adulthood, and offspring underwent an IP-GTT. Exposure to $\Delta 9$ -THC was associated with a decrease in fasting blood glucose at 5 m ($p = 0.0393$ for effect of $\Delta 9$ -THC by two-way ANOVA), though differences did not achieve statistical significance for either males or females when evaluated independently (Table 1). Fasting insulin levels, however, were not affected by gestational exposure to $\Delta 9$ -THC (Table 1), and HOMA1-IR scores were also not different between vehicle and $\Delta 9$ -THC-exposed offspring in males or females (data not shown). Following glucose challenge, female offspring exposed to $\Delta 9$ -THC exhibited elevated blood glucose at 5 min (18.37 ± 1.74 mmol/L for $\Delta 9$ -THC vs 11.48 ± 0.84 mmol/L for vehicle, adjusted p value = 0.0496), as well as an overall increased area under the curve for blood glucose (Fig. 2 A). This was associated with significantly augmented serum insulin concentration compared to vehicle offspring 15 min after glucose challenge (0.47 ± 0.05 ng/mL for $\Delta 9$ -THC vs 0.29 ± 0.07 for vehicle, adjusted p value = 0.0264), but which subsided to normal levels by 30 min (Fig. 2 B). No similar alterations were present in males (Fig. 2 C, D).




Fig. 1. Gestational exposure to $\Delta 9$ -THC reduces birthweight and pancreatic weight which is followed by catch-up growth. Bodyweight (A, B, C) and pancreatic weight (D, E, F) at PND1, PND21, and 5 m. *, $p < 0.05$ for vehicle vs $\Delta 9$ -THC assessed by Student's t test. Values are mean \pm SEM. n = mean of 4–5 litters per treatment (PND1), 16–23 individuals per treatment (PND21), 7–8 individuals per treatment per sex (5 m).

Table 1

Fasting blood glucose, insulin, and sex steroid levels at 5 m in male and female offspring. T, $p < 0.05$ for effect of gestational $\Delta 9$ -THC exposure; S, $p < 0.05$ for effect of sex by two-way ANOVA. Values are mean \pm SEM.

	Female		Male		Two-way ANOVA
	Vehicle	$\Delta 9$ -THC	Vehicle	$\Delta 9$ -THC	
Fasting Glucose (mmol/L)	6.038 \pm 0.2419	5.49 \pm 0.2157	6.064 \pm 0.155	5.714 \pm 0.1079	T
n =	13	10	14	7	
Fasting Insulin (ng/mL)	1.335 \pm 0.2164	1.473 \pm 0.2115	2.208 \pm 0.4498	2.253 \pm 0.5146	S
n =	8	10	12	6	
Fasting Estrogen (pg/mL)	57.34 \pm 5	57.75 \pm 4	N/A	N/A	N/A
n =	11	10			
Fasting Testosterone (ng/mL)	3.4 \pm 5	3.1 \pm 0.7	N/A	N/A	N/A
n =	9	7			




Fig. 2. Gestational exposure to $\Delta 9$ -THC causes glucose intolerance exclusively in female offspring at 5 months of age associated with increased serum insulin 15 min after glucose challenge. Blood glucose levels and integrated area under the curve for blood glucose tracked for 120 min following 2 g/kg glucose challenge in female and male offspring aged 5 months (A, C). Serum insulin concentrations 15 and 30 min after glucose challenge (B, D). *, $p < 0.05$ for vehicle vs $\Delta 9$ -THC assessed by repeated measures two-way ANOVA and Holm-Sidak-corrected post-hoc tests, or Student's t test (AUC). Values are mean \pm SEM. $n = 4$ –6 per treatment per sex.

Given the alterations in glucose tolerance and serum insulin concentration in $\Delta 9$ -THC-exposed female offspring, peripheral insulin sensitivity in the soleus muscle was examined after insulin challenge in these animals via phosphorylated Akt [Ser 473], a hallmark of insulin receptor activation and sensitivity [25]. Female $\Delta 9$ -THC-exposed offspring exhibited a significantly reduced ratio of pAkt [Ser 473] relative to total Akt2 protein expression after insulin challenge (Fig. 3; 0.58 ± 0.22 for $\Delta 9$ -THC vs 2.56 ± 0.64 for vehicle, adjusted p value = 0.0031).

Given the importance of estrogen and testosterone for glucose tolerance and preservation of β -cells [29–31], we also quantified fasting estrogen and testosterone levels at 5 m. However, analysis revealed that estrogen and testosterone were unaltered in females and males (Table 1), and both levels obtained were similar to those previously reported in young adult Wistar rats [23].

3.3. Altered endocrine pancreatic development

In female offspring, gestational $\Delta 9$ -THC exposure reduced the total density of islets at both PND21 (2.23 ± 0.17 islets/mm² for $\Delta 9$ -THC vs 2.78 ± 0.19 islets/mm² for vehicle, adjusted p value = 0.0247) and 5 m (1.32 ± 0.10 islets/mm² for $\Delta 9$ -THC vs 1.94 ± 0.22 islets/mm² for vehicle, adjusted p value = 0.0002), attributable to a specific decrease in the density of small islets (Fig. 4 A, G). By 5 m, this decrease in total islet density culminated in a reduction in β -cell mass in female $\Delta 9$ -THC-exposed offspring (Fig. 4 H; 6.68 ± 0.71 mg for $\Delta 9$ -THC vs 11.32 ± 1.76 mg for vehicle, adjusted p value = 0.0332). No equivalent alterations were apparent in male offspring (Fig. 4). This reduction in β -cell mass in female $\Delta 9$ -THC-exposed off-

spring was attributable exclusively to a reduction in the total number of islets, as mean islet size and mean β -cell area per islet were found to be unchanged at both weaning and in adulthood (Table 2).

To gain insight into possible mechanisms involved in the observed reductions in islet density and β -cell mass in $\Delta 9$ -THC-exposed females, endocrine cell neogenesis, as well as β - and α -cell replication and apoptosis were examined. Neogenesis was assessed via the density of extra-islet clusters ($< 500 \mu\text{m}^2$) [26,27], and was found to be unaltered at either PND21 or 5 m (Table 2). Pancreatic islet replication and apoptosis were examined in PND21 female offspring by immunohistochemistry with Ki67 antibody and TUNEL assay, respectively (Fig. 5 A (Ki67) and B (TUNEL)). PND21 was chosen given the reduction in islet density at this age and that replication and apoptosis in α - and β -cells declines markedly after this age [32,33]. Neither the percentage of β -cells (Ins⁺) or α -cells (Glucagon⁺) positive for Ki67 (Fig. 5 C, D), nor the percentage of β -cells (Ins⁺) positive for TUNEL (Fig. 5 E) were altered following $\Delta 9$ -THC exposure. No TUNEL-positive α -cells were observed in either group.

4. Discussion

Considering the increased popularity of *cannabis* in pregnancy [1], and the rising concentration of $\Delta 9$ -THC in *cannabis* over the last two decades [8], it is essential to understand the long-term metabolic effects of gestational exposure to this specific cannabinoid. In the current study, we and others utilized a physiologically-relevant dosage of $\Delta 9$ -THC (3 mg/kg) which yields maternal blood concentrations of $\Delta 9$ -THC (8.6–12.4 ng/mL), which are comparable to those seen after moderate recreational *cannabis* smoking in human adults (13–63 ng/




Fig. 3. Gestational exposure to $\Delta 9$ -THC impairs insulin response in muscle in female offspring aged 5 months. Ratio of phosphorylated Akt [Ser 473] relative to total Akt2 protein levels. **, $p < 0.01$; ns, not significant for vehicle vs $\Delta 9$ -THC assessed by two-way ANOVA and Holm-Sidak-corrected post-hoc tests. Values are mean \pm SEM. $n = 5-6$ per treatment, where each n value represents one offspring per dam.

mL), as well as in the tissue of aborted fetuses after maternal *cannabis* use (4–287 ng/mL) [19–21,34–36]. Previous studies in humans as well as animals have demonstrated that $\Delta 9$ -THC readily crosses the placenta, and that 10–28 % of maternal concentrations are detected in the fetal plasma, with 2–5X higher concentrations in fetal tissues [16–18]. However, given evidence that fetal plasma concentrations of $\Delta 9$ -THC may depend on the route of maternal exposure [16,37–39], quantification of $\Delta 9$ -THC in fetal circulation after maternal *i.p.* $\Delta 9$ -THC exposure would add insight to the results obtained in the present study. Nevertheless, this lower dosage of $\Delta 9$ -THC (3 mg/kg) led to fetal growth restriction, and unlike other studies employing 5 mg/kg [6], it did so *without* fetal demise [7]. Additionally, it should be noted that in same cohort of vehicle and $\Delta 9$ -THC offspring, we have published that $\Delta 9$ -THC exposure in pregnancy did not lead to changes in maternal food intake, maternal weight gain, or gestational length [7]. We demonstrated alterations in endocrine pancreatic development culminating in decreased β -cell mass, impaired glucose tolerance, and aberrant insulin response in adult female $\Delta 9$ -THC-exposed offspring.

At birth, both male and female offspring exposed to $\Delta 9$ -THC had reduced bodyweight and pancreatic weight. We have previously demonstrated that $\Delta 9$ -THC-exposed offspring also exhibit significant decreases in brain to bodyweight ratio, and liver to bodyweight ratio, indicative of symmetrical intrauterine growth restriction [7]. Both male and female $\Delta 9$ -THC-exposed offspring underwent catch-up growth by PND21, but only females were glucose intolerant at 5 m. Specifically, 5-month-old females exposed to $\Delta 9$ -THC exhibited elevated blood glucose 5 min after glucose challenge and overall elevated area under the curve for blood glucose. Interestingly, this occurred in association with increased serum insulin 15 min after glucose challenge, suggesting that peripheral insulin resistance, leading to impaired glucose disposal, contributes to the observed glucose intolerance. To explore this further, we challenged vehicle and $\Delta 9$ -THC-exposed female offspring with insulin and found that $\Delta 9$ -THC-exposed offspring demonstrated blunted pAkt [Ser473] activation, a hallmark of insulin receptor activation and sensitivity [25], in the soleus muscle.

In addition to the observed impairments in downstream insulin signaling, $\Delta 9$ -THC-exposed females also exhibited altered development of the endocrine pancreas. Specifically, these $\Delta 9$ -THC female offspring had decreased total and small islet density at both PND21 and 5 m, along with a significant reduction (41 %) in β -cell mass at 5 m. Despite this decrease in β -cell mass, fasting serum insulin levels were maintained, suggesting some degree of β -cell compensation [40]. Other studies have shown that this type of β -cell compensation is implicated in eventual β -cell exhaustion [41], which along with chronic glucose in-

tolerance, culminates in β -cell death [42], or dedifferentiation [43] seen in the progression from prediabetes to overt type 2 diabetes. It will therefore be interesting to explore the future metabolic health of these offspring in later life (*i.e.* > 6 months of age).

The sexual dimorphism observed in this study is noteworthy as it has previously been demonstrated that female offspring exposed to $\Delta 9$ -THC *in utero* exhibit neurodevelopmental abnormalities pertaining to addiction, while their male counterparts do not, suggesting that female offspring may be more sensitive to the effects of $\Delta 9$ -THC *in utero* [44]. Based on the collective results of this current study, the mechanisms responsible for the reductions in islet density and β -cell mass in female $\Delta 9$ -THC-exposed offspring remain elusive. While decreases in estrogen is associated with β -cell apoptosis [30], no changes were apparent in either estrogen or testosterone levels, and no alterations were apparent in postnatal (PND21) β - or α -cell replication, neogenesis, or apoptosis. This contrasts with what is observed in offspring exposed to another psychoactive compound, nicotine. When administered during gestation, nicotine-exposed offspring exhibit reduced β -cell mass at four weeks of age, but not 26 weeks of age, due to compensatory increased islet cell proliferation at four weeks [45]. However, this compensatory mechanism is not seen when nicotine exposure extends through lactation, and islet cell apoptosis is now increased at 4 weeks of age, resulting in reduced β -cell mass from 4 to 26 weeks [45]. Despite lacking any such alterations in islet cell proliferation, apoptosis, or neogenesis, we still observed reduced islet density in both PND21 and 5 m female $\Delta 9$ -THC exposed offspring. It is therefore likely that the glucometabolic effects of $\Delta 9$ -THC incurred earlier during prenatal development. For example in a model of maternal malnutrition (total caloric intake reduced to 50 % of controls), fetal growth restricted offspring exhibited reduced β -cell mass at birth (PND1) and PND21, attributable to reduced islet density and β -cell size, with no alternations present in β -cell replication at PND1 or PND21 [46,47]. $\Delta 9$ -THC may be particularly deleterious to β -cell development during prenatal life, first due to the effects of $\Delta 9$ -THC on placental insufficiency leading to less maternal transfer of oxygen and nutrients for fetal pancreatic development [6,7]. This is consistent with the thrifty phenotype hypothesis, in which adaptations suited to malnutrition *in utero* contribute to metabolic disease in later life [48]. Noteworthy for the present study are adaptations concerning the endocrine pancreas, namely reduced β -cell mass and reduced β -cell number at 90 % gestation, as observed in growth-restricted fetal sheep [49]. Similar adaptations may contribute to the altered endocrine pancreatic development as seen in our $\Delta 9$ -THC-exposed female offspring, warranting further investigation related to perinatal windows of $\Delta 9$ -THC exposure. Secondly, $\Delta 9$ -THC *in utero*




Fig. 4. Gestational exposure to $\Delta 9$ -THC reduces small islet and total islet density at PND21 and 5 m, and reduces β -cell mass at 5 m exclusively in female offspring. Density of small ($500\text{--}5000\ \mu\text{m}^2$), medium ($5000\text{--}10000\ \mu\text{m}^2$), large ($>10000\ \mu\text{m}^2$), and total islets for female (A, G) and male (D, J) offspring at PND21 and 5 m. β -cell mass (B, E, H, K) and α -cell mass (C, F, I, L) in female and male offspring at PND21 and 5 m. *, $p < 0.05$; ***, $p < 0.001$ for vehicle vs $\Delta 9$ -THC assessed by repeated measures two-way ANOVA and Holm-Sidak-corrected post-hoc tests. Values are mean \pm SEM. $n = 4\text{--}7$ per treatment per sex per age (2–3 technical replicates per animal), where each n value represents one offspring per dam.

Table 2

Endocrine pancreas morphometry in female and male offspring at PND21 and 5 m. S, $p < 0.05$ for effect of sex by two-way ANOVA; ns, not significant. Values are mean \pm SEM. $n = 4\text{--}7$ per treatment per sex (2–3 slides per animal).

		Female		Male		Two-way ANOVA
		$\Delta 9$ -THC	Vehicle	$\Delta 9$ -THC	Vehicle	
Mean Islet Size (μm^2)	PND21	3627 \pm 272.5	3592 \pm 349.5	4537 \pm 471.3	3995 \pm 147.4	ns
	5 m	6322 \pm 533.2	7341 \pm 514.2	8394 \pm 861	7228 \pm 236.4	ns
Mean β -cell Area per Islet (μm^2)	PND21	2226 \pm 117.8	2750 \pm 381.6	3326 \pm 243.9	2812 \pm 163.5	S
	5 m	4999 \pm 488.1	5781 \pm 453.1	6691 \pm 538.8	6000 \pm 34.89	ns
Clusters per mm ² Pancreas	PND21	4.092 \pm 0.5584	3.256 \pm 0.6557	2.603 \pm 0.1859	2.852 \pm 0.1772	S
	5 m	0.795 \pm 0.1543	0.802 \pm 0.0814	0.644 \pm 0.1359	0.689 \pm 0.134	ns




Fig. 5. Gestational exposure to Δ9-THC does not alter replication or apoptosis in β- or α-cells in females at PND21. Representative immunohistochemistry of an islet immunostained for Ki67 (A) and TUNEL (B). Percentage of β-cells positive for Ki67 (C); percentage of α-cells positive for Ki67 (D); percentage of β-cells positive for TUNEL (E). Values are mean ± SEM. $n = 4-5$ per treatment (2-3 technical replicates per animal), where each n value represents one offspring per dam.

may have direct adverse glucometabolic effects on the developing fetal pancreas via its interaction with CB1R [11,12]. In particular, *in vitro* studies have shown that CB1R activation suppresses mitogenic signaling in β-cells via decreased expression of cyclin D2 and impaired downstream insulin signaling, and promotes apoptosis through decreased expression of Bcl-2 [11,12]. As such, there is potential for a direct impact of Δ9-THC on development of the fetal endocrine pancreas, but further work is needed to investigate if such alterations are apparent in the fetal pancreas after Δ9-THC exposure *in vivo*.

5. Conclusion

In summary, though to date no associations have yet been found between maternal *cannabis* use and offspring development of type 2 diabetes in humans, future prospective clinical studies are warranted given that the major increases in Δ9-THC content in *cannabis* have only occurred over the past two decades. This is the first study to demonstrate a link between maternal Δ9-THC exposure and predisposition to develop glucose intolerance in female offspring. This raises concern about the future metabolic health of offspring exposed to Δ9-THC *in utero*, particularly as this study examined the effects of Δ9-THC at levels which are conservative estimates of what is available in recreational *cannabis* today.

Funding

This work was supported by an operating grant from the Women's Developmental Council (LHSC) to DBH and the Canadian Institutes of Health Research (CRU-163023 to DBH, PJT-155981 to ACH).

Author contribution

R.G. and S.V. were involved in design, execution, and interpretation of experiments. K.L. was involved with implementing the animal model. D.B.H. and E.A. were involved in the design of this project and interpretation of results. S.R.L. contributed to the design of the animal experiments and in the preparation and dosing of vehicle and Δ9-THC *in vivo*. A.C.H. played a role in the design of the animal experiments and in the analysis of metabolic outcomes. R.G. wrote the first draft of the manuscript; all authors edited drafts of the manuscript.

Data availability

The datasets generated and analyzed in the current study are available from the corresponding author upon reasonable request.

Declaration of interest

There is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Acknowledgements

We thank Linda Jackson-Boeters for preparing tissue sections, Shannon Seney for assisting in the measurement of sex steroid hormones, and the Women's Development Council (LHSC) for funding this project.

References

- [1] K C Young-Wolff, L Y Tucker, S Alexeeff, M A Armstrong, A Conway, C Weisner, N Goler, Trends in self-reported and biochemically tested Marijuana use among pregnant females in California from 2009–2016, *J. Am. Med. Assoc.* 318 (24) (2017) 2490–2491.
- [2] R E Westfall, P A Janssen, P Lucas, R Capler, Survey of medicinal cannabis use among childbearing women: patterns of its use in pregnancy and retrospective self-assessment of its efficacy against 'morning sickness', *Complement. Ther. Clin. Pract.* 12 (1) (2006) 27–33.
- [3] C Saint Louis, A Balm When You're Expecting: Sometimes Pot Does the Trick, *The New York Times*, 2017.
- [4] D S Hasin, US epidemiology of Cannabis use and associated problems, *Neuropsychopharmacology* 43 (1) (2018) 195–212.
- [5] H Bayrampour, M Zahradnik, S Lisonkova, P Janssen, Women's perspectives about cannabis use during pregnancy and the postpartum period: an integrative review, *Prev. Med. (Baltim.)* 119 (2019) 17–23.
- [6] X Chang, Y Bian, Q He, J Yao, J Zhu, J Wu, K Wang, T Duan, Suppression of STAT3 signaling by Δ9-tetrahydrocannabinol (THC) induces trophoblast dysfunction, *Cell. Physiol. Biochem.* 42 (2) (2017) 537–550.
- [7] B V V Natale, K N Gustin, K Lee, A C Holloway, S R Laviolette, D R C Natale, D B Hardy, Δ9-tetrahydrocannabinol exposure during rat pregnancy leads to symmetrical fetal growth restriction and labyrinth-specific vascular defects in the placenta, *Eur. J. Pharmacol.* 10 (2010) 544.
- [8] M A Elsohly, Z Mehmedic, S Foster, C Gon, S Chandra, J C Church, Archival report changes in Cannabis potency over the last 2 decades (1995–2014): analysis of current data in the United States, *Biol. Psychiatry* 79 (2016) 613–619.
- [9] C N Hales, D J P Barker, Type 2 (non-insulin-dependent) diabetes mellitus: the thrifty phenotype hypothesis, *Diabetologia* 35 (1992) 595–601.
- [10] K Malenczyk, E Keimpema, F Piscitelli, D Calvigioni, P Björklund, K Mackie, V Di Marzo, T G M Hökfelt, A Dobryzn, T Harkany, Fetal endocannabinoids orchestrate

- the organization of pancreatic islet microarchitecture, *Proc. Natl. Acad. Sci.* 112 (45) (2015) E6185–E6194.
- [11] W Kim, M E Doyle, Z Liu, Q Lao, Y-K-K Shin, O D Carlson, H S Kim, S Thomas, J K Napora, E K Lee, R Moaddel, Y Wang, S Maudsley, B Martin, R N Kulkarni, J M Egan, Cannabinoids inhibit insulin receptor signaling in pancreatic β -cells, *Diabetes* 60 (2011) 1198–1209.
 - [12] J Kim, K J Lee, J S Kim, J G Rho, J J Shin, W K Song, E K Lee, J M Egan, W Kim, Cannabinoids regulate Bcl-2 and cyclin D2 expression in pancreatic β cells, *PLoS One* 11 (3) (2016) e0150981.
 - [13] F J Bermúdez-Silva, J Suárez, E Baixeras, N Cobo, D Bautista, A L Cuesta-Muñoz, E Fuentes, P Juan-Pico, M J Castro, G Milman, R Mechoulam, A Nadal, F Rodríguez De Fonseca, Presence of functional cannabinoid receptors in human endocrine pancreas, *Diabetologia* 51 (3) (2008) 476–487.
 - [14] B Brocato, A A Zoerner, Z Janjetovic, C Skobowiat, S Gupta, B M Moore, A Slominski, J Zhang, M Schenone, R Phinehas, R J Ferry, E Dick, G B Hubbard, G Mari, N Schlambitz-Loutsevitch, Endocannabinoid crosstalk between placenta and maternal fat in a baboon model (*Papio* spp.) of obesity, *Placenta* 34 (11) (2013) 983–989.
 - [15] F Berrendero, N Sepe, Ja Ramos, V Di Marzo, J J Fernández-Ruiz, Analysis of cannabinoid receptor binding and mRNA expression and endogenous cannabinoid contents in the developing rat brain during late gestation and early postnatal period, *Synapse* 33 (3) (1999) 181–191.
 - [16] D E Hutchings, B R Martin, Z Gamagaris, N Miller, T Fico, Plasma concentrations of delta-9-tetrahydrocannabinol in dams and fetuses following acute or multiple prenatal dosing in rats, *Life Sci.* 44 (11) (1989) 697–701.
 - [17] J Bailey, Fetal disposition of Δ -9-tetrahydrocannabinol (THC) during late pregnancy in the rhesus monkey, *Toxicol. Appl. Pharmacol.* 90 (2) (1987) 315–321.
 - [18] K S Grant, R Petroff, N Isoherranen, N Stella, T M Burbacher, Cannabis use during pregnancy: pharmacokinetics and effects on child development, *Pharmacol. Ther.* J. (182) (2018) 133–151.
 - [19] D M Schwope, E L Karschner, D A Gorelick, M A Huestis, Identification of recent cannabis use: whole-blood and plasma free and glucuronidated cannabinoid pharmacokinetics following controlled smoked cannabis administration, *Clin. Chem.* 57 (10) (2011) 1406–1414.
 - [20] M Falcon, S Pichini, J Joya, M Pujadas, A Sanchez, O Vall, O García Algar, A Luna, R De La Torre, M C Rotolo, M C Pellegrini, Maternal hair testing for the assessment of fetal exposure to drug of abuse during early pregnancy: comparison with testing in placental and fetal remains §, *Forensic Sci. Int.* 218 (2011) 92–96.
 - [21] C Klein, E Karanges, A Spiro, A Wong, J Spencer, T Huynh, N Gunasekaran, T Karl, L E Long, X F Huang, K Liu, J C Arnold, I S McGregor, Cannabidiol potentiates Δ 9-tetrahydrocannabinol (THC) behavioural effects and alters THC pharmacokinetics during acute and chronic treatment in adolescent rats, *Psychopharmacology (Berl.)* 218 (2) (2011) 443–457.
 - [22] R D Harbison, B Mantilla-Plata, Prenatal toxicity, maternal distribution and placental transfer of tetrahydrocannabinol, *J. Pharmacol. Exp. Ther.* 180 (2) (1972) 446–453.
 - [23] A Chamson-Reig, S M Thyssen, D J Hill, E Arany, Exposure of the pregnant rat to low protein diet causes impaired glucose homeostasis in the young adult offspring by different mechanisms in males and females, *Exp. Biol. Med.* 234 (12) (2009) 1425–1436.
 - [24] T M Wallace, J C Levy, D R Matthews, Use and abuse of HOMA modeling, *Diabetes Care* 27 (6) (2004) 1487–1495.
 - [25] R W Mackenzie, B T Elliott, Akt/PKB activation and insulin signaling: a novel insulin signaling pathway in the treatment of type 2 diabetes, *Diabetes Metab. Syndr. Obes. Targets Ther.* 7 (2014) 55.
 - [26] T Mezza, G Muscogiuri, G P Sorice, G Clemente, J Hu, A Pontecorvi, J J Holst, A Giaccari, R N Kulkarni, Insulin resistance alters islet morphology in nondiabetic humans, *Diabetes* 63 (3) (2014) 994–1007.
 - [27] C A Beamish, L Zhang, S K Szlapinski, B J Strutt, D J Hill, An increase in immature β -cells lacking Glut2 precedes the expansion of β -cell mass in the pregnant mouse, 2017.
 - [28] N Ben-Othman, A Vieira, M Courtney, et al., Long-term GABA administration induces alpha cell-mediated beta-like cell neogenesis, *Cell* 168 (1–2) (2017) 73–85.e11.
 - [29] C J Bailey, H Ahmed-Sorour, Role of ovarian hormones in the long-term control of glucose homeostasis effects on insulin secretion, *Diabetologia* 19 (1980) 475–481.
 - [30] C Le May, K Chu, M Hu, C S Ortega, E R Simpson, K S Korach, M-J-J Tsai, F Mauvais-Jarvis, Estrogens protect pancreatic-cells from apoptosis and prevent insulin-deficient diabetes mellitus in mice, *PNAS* 103 (24) (2006) 9232–9237.
 - [31] K Yabiku, K Nakamoto, A Tokushige, Reintroducing testosterone in the db/db mouse partially restores normal glucose metabolism and insulin resistance in a leptin-independent manner, *BMC Endocr. Disord.* 18 (38) (2018).
 - [32] S Bonner-Weir, C Aguayo-Mazzucato, G C Weir, Dynamic development of the pancreas from birth to adulthood, *Ups J. Med. Sci.* 121 (2) (2016) 155–158.
 - [33] J Petrik, E Arany, T J McDonald, D J Hill, Apoptosis in the pancreatic islet cells of the neonatal rat is associated with a reduced expression of insulin-like growth factor II that may act as a survival factor, *Endocrinology* 139 (6) (1998) 2994–3004.
 - [34] G Tortoriello, C V V Morris, A Alpar, J Fuzik, S L Shirran, D Calvigioni, E Keimpema, C H Botting, K Reinecke, T Herdegen, M Courtney, Y L Hurd, T Harkany, Miswiring the brain: Δ 9-tetrahydrocannabinol disrupts cortical development by inducing an SCG10/stathmin-2 degradation pathway, *EMBO J.* 33 (2014) 668–685.
 - [35] S Mato, V Chevalere, D Robbe, A Pazos, P E Castillo, O J Manzoni, A single in-vivo exposure to Δ 9THC blocks endocannabinoid-mediated synaptic plasticity, *Nat. Neurosci.* 7 (6) (2004) 585–586.
 - [36] T U C Järbe, R S Gifford, A Makriyannis, Antagonism of Δ^9 -THC induced behavioral effects by rimonabant: time course studies in rats, *Eur. J. Pharmacol.* 648 (1–3) (2010) 133–138.
 - [37] R M Abrams, C E Cook, K H Davis, K Niederreither, M J Jaeger, H H Szeto, Plasma delta-9-tetrahydrocannabinol in pregnant sheep and fetus after inhalation of smoke from a marijuana cigarette, *Alcohol Drug Res.* 6 (5) (1985) 361–369.
 - [38] B R Martin, W L Dewey, L S Harris, J S Beckner, 3H-delta9-tetrahydrocannabinol distribution in pregnant dogs and their fetuses, *Res. Commun. Chem. Pathol. Pharmacol.* 17 (3) (1977) 457–470.
 - [39] F Grotenhermen, Clinical pharmacokinetics of cannabinoids, *J. Cannabis Ther.* 3 (1) (2003) 3–51.
 - [40] J J Meier, R C Bonadonna, Role of reduced β -cell mass versus impaired β -cell function in the pathogenesis of type 2 diabetes, *Diabetes Pathophysiol.* 36 (S2) (2013) 113–119.
 - [41] C Chen, C M Cohrs, J Stertmann, R Bozsak, S Speier, Human beta cell mass and function in diabetes: recent advances in knowledge and technologies to understand disease pathogenesis, *Mol. Metab.* 6 (9) (2017) 943–957.
 - [42] A E Butler, J Janson, S Bonner-Weir, R Ritzel, R A Rizza, P C Butler, Beta-cell deficit and increased beta-cell apoptosis in humans with type 2 diabetes, *Diabetes* 52 (1) (2003) 102–110.
 - [43] Z Wang, N W York, C G Nichols, M S Remedi, Pancreatic β cell dedifferentiation in diabetes and redifferentiation following insulin therapy, *Cell Metab.* 19 (5) (2014) 872–882.
 - [44] G Vela, S Martín, L García-Gil, J Antonio Crespo, M Ruiz-Gayo, J Javier Fernandez-Ruiz, C García-Lecumberri, D Pelaprat, J Angel Fuentes, J Antonio Ramos, E Ambrosio, Maternal exposure to Δ 9-tetrahydrocannabinol facilitates morphine self-administration behavior and changes regional binding to central m opioid receptors in adult offspring female rats, *Brain Res.* 807 (1998) 101–109.
 - [45] J E Bruin, L D Kellenberger, H C Gerstein, K M Morrison, A C Holloway, Fetal and neonatal nicotine exposure and postnatal glucose homeostasis: identifying critical windows of exposure, *J. Endocrinol.* 194 (1) (2007) 171–178.
 - [46] T Inoue, Y Kido, S Asahara, T Matsuda, Y Shibutani, M Koyanagi, M Kasuga, Effect of intrauterine undernutrition during late gestation on pancreatic beta cell mass, *Biomed. Res.* 30 (6) (2009) 325–330.
 - [47] A Garofano, P Czernichow, B Bréant, In utero undernutrition impairs rat beta-cell development, *Diabetologia* 40 (10) (1997) 1231–1234.
 - [48] C N Hales, D J P Barker, The thrifty phenotype hypothesis, *Br. Med. Bull.* 60 (1) (2001) 5–20.
 - [49] S W Limesand, J Jensen, J C Hutton, W W Hay, Diminished-cell replication contributes to reduced-cell mass in fetal sheep with intrauterine growth restriction, *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 288 (2005) 1297–1305.