

10-1-2014

Factors affecting transfusion requirement after hip fracture: Can we reduce the need for blood?

Sagar J. Desai
Western University

Kristi S. Wood
Western University

Jackie Marsh
Western University

Dianne Bryant
Western University

Hussein Abdo
Western University

See next page for additional authors

Follow this and additional works at: <https://ir.lib.uwo.ca/boneandjointpub>



Part of the [Medicine and Health Sciences Commons](#)

Citation of this paper:

Desai, Sagar J.; Wood, Kristi S.; Marsh, Jackie; Bryant, Dianne; Abdo, Hussein; Lawendy, Abdel Rahman; and Sanders, David W., "Factors affecting transfusion requirement after hip fracture: Can we reduce the need for blood?" (2014). *Bone and Joint Institute*. 175.

<https://ir.lib.uwo.ca/boneandjointpub/175>

Authors

Sagar J. Desai, Kristi S. Wood, Jackie Marsh, Dianne Bryant, Hussein Abdo, Abdel Rahman Lawendy, and David W. Sanders

Factors affecting transfusion requirement after hip fracture: Can we reduce the need for blood?

Sagar J. Desai, MD
 Kristi S. Wood, MSc, MD
 Jackie Marsh, MSc, PhD
 Dianne Bryant, MSc, PhD
 Hussein Abdo, BSc
 Abdel-Rahman Lawendy, MSc,
 MD
 David W. Sanders, MD, MSc

From Western University, London, Ont.

Accepted for publication
 Apr. 7, 2014

Correspondence to:

S.J. Desai
 Chief Resident, Orthopaedic Surgery
 St. Joseph's Health Centre
 268 Grosvenor St., Rm D0-213
 London ON N6A 4L6
 sjdesai@uwo.ca

DOI: 10.1503/cjs.030413

Background: Hip fractures are common injuries that result in blood loss and frequently require the transfusion of blood products. We sought to identify risk factors leading to increased blood transfusion in patients presenting with hip fractures, especially those factors that are modifiable.

Methods: We retrospectively reviewed the cases of all patients who had fixation of their hip fractures between October 2005 and February 2010. The need for transfusion was correlated with potential risk factors, including age, sex, preoperative hemoglobin, fracture type, fixation method and more.

Results: A total of 835 patients had fixation of their hip fractures during the study period; 631 met the inclusion criteria and 249 of them (39.5%) were transfused. We found an association between need for blood transfusion and female sex ($p = 0.018$), lower preoperative hemoglobin ($p < 0.001$), fracture type ($p < 0.001$) and fixation method ($p < 0.001$). Compared with femoral neck fractures, there was a 2.37 times greater risk of blood transfusion in patients with intertrochanteric fractures ($p < 0.001$) and a 4.03 times greater risk in those with subtrochanteric fractures ($p < 0.001$). Dynamic hip screw (DHS) fixation decreased the risk of transfusion by about half compared with intramedullary nail or hemiarthroplasty. We found no association with age, delay to operation ($p = 0.17$) or duration of surgery ($p = 0.30$).

Conclusion: The only modifiable risk factor identified was fixation method. When considering blood transfusion requirements in isolation, we suggest a potential benefit in using a DHS for intertrochanteric and femoral neck fractures amenable to DHS fixation.

Contexte : La fracture de la hanche est un traumatisme fréquent, qui cause une perte sanguine et nécessite souvent la transfusion de produits sanguins. Nous avons tenté d'identifier les facteurs de risque associés à une hausse du nombre des transfusions sanguines chez des patients ayant subi une fracture de la hanche, en particulier les facteurs modifiables.

Méthodes : Au cours d'une étude rétrospective, on a revu les cas de tous les patients chez qui on avait pratiqué une ostéosynthèse pour une fracture de la hanche survenue entre octobre 2005 et février 2010. La nécessité d'une transfusion sanguine a été associée à d'éventuels facteurs de risque, dont l'âge, le sexe, le taux d'hémoglobine préopératoire, le type de fracture, la technique d'ostéosynthèse, et d'autres facteurs encore.

Résultats : Au total, 835 patients avaient subi une ostéosynthèse pour fracture de la hanche au cours de la période à l'étude; 631 satisfaisaient les critères d'inclusion à l'étude et parmi eux, 249 (39,5 %) ont reçu une transfusion sanguine. On a observé l'existence d'un lien entre la nécessité d'une transfusion sanguine et le sexe féminin ($p = 0,018$), une plus faible concentration d'hémoglobine préopératoire ($p < 0,001$), le type de fracture ($p < 0,001$) et la technique d'ostéosynthèse ($p < 0,001$). Par rapport aux fractures du col fémoral, le risque de transfusion sanguine était 2,37 fois plus élevé chez les patients présentant une fracture intertrochantérienne ($p < 0,001$) et 4,03 fois plus élevé chez ceux présentant une fracture sous-trochantérienne ($p < 0,001$). En utilisant une vis dynamique de hanche, le risque de transfusion sanguine a diminué d'environ 50 % par rapport à l'enclouage centromédullaire ou à l'hémiarthroplastie. Aucun lien n'a été observé avec l'âge, le délai de l'intervention chirurgicale ($p = 0,17$), ni avec sa durée ($p = 0,30$).

Conclusion : La technique d'ostéosynthèse est l'unique facteur de risque modifiable ayant été identifié. Mais lorsqu'on évalue la nécessité d'une transfusion sanguine sans tenir compte des facteurs de risque, nos résultats semblent indiquer qu'on aurait avantage à utiliser une vis dynamique de hanche pour consolider les fractures intertrochantériennes et les fractures du col fémoral.

Hip fractures are among the most common fractures treated by orthopedic surgeons. With increasing life expectancy, it is estimated that the number of people aged 65 and older will increase from the recent estimate of 323 million to 1555 million by the year 2050. As a result, it is estimated that the number of hip fractures occurring worldwide will increase from 1.66 million in 1990 to 6.26 million in 2050.¹

Hip fractures result in blood loss and frequently require the transfusion of blood products. While blood transfusions are potentially life-saving interventions, they can also cause patient morbidity. Blood transfusions are correlated with an increased risk of bacterial infections²⁻⁷ and possibly increased mortality.^{8,9} There are also substantial costs involved in the collection, preparation, transport and administration of blood.

In the United States, more than 15 million units of blood are transfused annually.¹⁰ Many of these transfusions are given to surgical patients, including elderly patients with hip fractures. We sought to determine whether any modifiable risk factors for transfusion exist. The purpose of this study, therefore, was to assess risk factors for blood transfusion requirements in patients presenting with hip fractures. We performed a retrospective study at a single level I trauma centre from 2005 to 2010. Blood transfusion requirements were correlated with patient variables, such as age, sex, delay to surgery, duration of surgery, preoperative hemoglobin, fracture type and fixation method.

METHODS

All patients undergoing surgical fixation of their hip fractures in a single academic trauma centre between October 2005 and February 2010 were included in this retrospective study. The fracture patterns included were femoral neck, intertrochanteric and subtrochanteric fractures. Fixation methods used included hemiarthroplasty, dynamic hip screw (DHS), cannulated screws and cephalomedullary nails. The study was approved by our institution's review board. All patient information, including laboratory values and operative notes, were collected from our institution's electronic patient database, Powerchart (Cerner Corporation). Blood transfusion information was collected from our institution's blood transfusion laboratory.

The type of hip fracture was documented based on review of patients' preoperative and postoperative radiographs. These included femoral neck fractures (AO-OTA 31-B¹⁻³), intertrochanteric fractures (AO-OTA 31-A¹⁻³) and subtrochanteric fractures (AO-OTA 32A,¹⁻³ 32-B[1-3] and 32-C[1-3]). Surgical fixation was documented based on review of patients' preoperative and postoperative radiographs. The DHS used was from Synthes. All hemiarthroplasties were performed with the Conquest system (Smith & Nephew) through a standard Hardinge approach. The intramedullary nails were either Trigen or Intertan (both from Smith & Nephew). The cannulated screws were 7.3-

mm screws (Synthes Inc). The following measures were recorded from the patients' electronic charts: sex; age; American Society of Anesthesiologist (ASA) score; duration of surgery; and need for preoperative, intraoperative and postoperative blood transfusion. Time from admission to operation was also documented and was defined as the time from admission to our institution's emergency department to the start of the operation. Criteria for administration of blood transfusion were a hemoglobin value less than 70 g/L or less than 80 g/L with signs/symptoms of anemia.

Patients were excluded from this study if they were younger than 60 years; had a known cancer in the region of the fracture; had a hemorrhagic complication in another anatomic site, such as a gastrointestinal bleed, pre- or postoperatively; were admitted to hospital for other clinically important comorbidities, such as sepsis or polytrauma; were undergoing revision surgery; had an intraoperative complication, such as a trochanteric fracture; were therapeutically anticoagulated or had documented hematologic disease before surgery; or had a delay in diagnosis of their fracture longer than 1 week.

Notably, all patients received thromboembolic prophylaxis with 5000 units of low molecular-weight heparin (dalteparin sodium, Pfizer Canada). This therapy was started on admission, withheld on the day of surgery and restarted on the first postoperative day.

Statistical analysis

We performed our statistical analyses using SPSS software. We conducted a univariate analysis of all independent variables (i.e., age, sex, duration of surgery, preoperative hemoglobin level, fracture type, fixation method) to establish an association with blood transfusion requirement. Variables that were found to have a significant association were then included in a multivariate analysis. We considered results to be significant at $p < 0.05$.

RESULTS

A total of 835 patients had fixation of their hip fractures during the study period; 631 patients met the inclusion criteria (Fig. 1). Forty-one patients were excluded because of multiple injuries, 73 patients were anticoagulated, 32 had pathologic or pending pathologic fractures, 23 had a significant hemorrhagic complication unrelated to their hip fracture, 16 underwent revision surgery, 9 experienced an intraoperative complication, 4 were excluded because of poor documentation and 6 were delayed to surgery because the fracture diagnosis was delayed longer than 1 week (Fig. 1).

The mean age was 81.6 (range 60–100) years. The sample comprised 455 women (72.1%, mean age 82.4 yr) and 176 men (27.9%, mean age 79.3 yr; Table 1). The mean delay from admission to surgery was 48.9 hours. Of the 631 patients in the study, 249 patients were transfused (39.5%). This group included 26% of patients with femoral




Fig. 1. Patients meeting inclusion criteria. Cann screws = cannulated screws; DHS = dynamic hip screw; hemi = hemiarthroplasty; intertroch = intertrochanteric fracture; nail = intramedullary nail; subtroch = subtrochanteric fracture.

Table 1. Characteristics of study participants

Characteristic	No. (%),* n = 618
Age, mean ± SD yr	81.6 ± 8.7
Sex, female	444 (71.8)
Delay to OR in h	
mean ± SD	48.9 ± 33.4
median (IQR)	43.3 (26.1–63.2)
Duration of surgery, mean ± SD min	63.3 ± 23.8
Preoperative hemoglobin, mean ± SD	118.8 ± 16.0
Fracture type	
Femoral neck	318 (51.5)
Intertrochanteric	254 (41.1)
Subtrochanteric	44 (7.1)
Fixation type	
Hemiarthroplasty	262 (42.4)
DHS	246 (39.8)
IM nail	108 (17.5)
Fracture by fixation	
Femoral neck — hemiarthroplasty	262 (42.5)
Femoral neck — DHS	56 (9.1)
Intertrochanteric — DHS	190 (30.8)
Intertrochanteric — IM nail	64 (10.4)
Subtrochanteric — IM nail	44 (7.1)

DHS = dynamic hip screw; IM = intramedullary; IQR = interquartile range; OR = operating room; SD = standard deviation.
*Unless otherwise indicated.

neck fractures, 52% with intertrochanteric fractures and 71% with subtrochanteric fractures. The majority of blood transfusions were administered on postoperative day 1, 2 or 3 (Fig. 2).

The results of the univariate analysis are presented in Table 2. The 13 patients with femoral neck fractures treated with cannulated screws were removed from the analysis because of the small number of patients. The univariate analysis demonstrated an association between need for blood transfusion and increased age ($p = 0.004$), female sex ($p =$

0.018), lower preoperative hemoglobin level ($p < 0.001$), fracture type ($p < 0.001$) and fixation method ($p < 0.001$). Patients requiring a blood transfusion had an average ASA score of 3.4, whereas patients not receiving a transfusion had an average score of 3.3 ($p < 0.001$). We found no association with delay to operation ($p = 0.17$) or duration of surgery ($p = 0.30$).

Multivariate analysis demonstrated an association between blood transfusion requirement and 3 variables (Table 3). Men were at half the risk of women (1.54 odds ratio [OR], 95% confidence interval [CI] 1.002–2.36,




Fig. 2. Number of patients receiving blood transfusions preoperatively, intraoperatively and postoperatively. Most transfusions were performed on postoperative day 1, 2 or 3. OR = intraoperative transfusion; preop 0 = preoperative transfusion day of surgery; postop 0 = postoperative transfusion day of surgery.

Table 2. Univariate logistic regression analyses to predict transfusion, $n = 618$

Potential predictor	OR (95% CI)	p value
Age, 5-yr increase	1.16 (1.05–1.28)	0.004
Sex, women v. men	1.61 (1.08–2.38)	0.018
Delay to operating room, 8-h increase	0.97 (0.93–1.01)	0.17
Duration of surgery, 5-min increase	0.98 (0.95–1.02)	0.30
Preoperative hemoglobin, 10-point increase	0.65 (0.58–0.73)	< 0.001
Fracture type		< 0.001
Intertrochanteric v. femoral neck	2.37 (1.65–3.39)	
Subtrochanteric v. femoral neck	4.03 (2.11–7.70)	
Fracture by fixation		< 0.001
Femoral neck, DHS v. hemiarthroplasty	0.51 (0.23–1.12)	
Intertrochanteric, DHS v. intramedullary nail	0.57 (0.32–1.02)	

CI = confidence interval; DHS = dynamic hip screw; OR = odds ratio.

Table 3. Multivariable analyses to predict transfusion. Logistic regression with only those variables that were significant at $p = 0.05$ in univariate analyses, $n = 618$

Potential predictor	Adjusted OR (95% CI)	p value
Age, 5-yr increase	1.06 (0.95–1.19)	0.27
Sex, women v. men	1.54 (1.002–2.36)	0.049
Preoperative hemoglobin, 10-point increase	0.69 (0.61–0.78)	< 0.001
Fracture, fixation		< 0.001
Femoral neck, DHS v. hemiarthroplasty	0.49 (0.21–1.12)	
Intertrochanteric, DHS v. intramedullary nail	0.52 (0.29–0.95)	

CI = confidence interval; DHS = dynamic hip screw; OR = odds ratio.

$p = 0.049$). For every 10 g/L increase in preoperative hemoglobin, the risk of transfusion decreased by about 30% (OR 0.69, 95% CI 0.61–0.78, $p < 0.001$). Femoral neck fractures treated with DHS reduced the risk of transfusion by half compared with hemiarthroplasty (OR 0.49, 95% CI 0.61–0.78), and intertrochanteric fractures treated with DHS reduced the risk of transfusion by half compared with intramedullary nail (OR 0.52, 95% CI 0.29–0.95). This indicates that DHS fixation in either femoral neck fractures or intertrochanteric fractures decreases the risk of transfusion by about half compared with treatment with an intramedullary nail or hemiarthroplasty.

DISCUSSION

We found a correlation between blood transfusion requirement and sex, preoperative hemoglobin, ASA score, fracture type and fixation method. No correlation was found with age, duration of surgery or delay from admission to operation.

Low preoperative hemoglobin is an independent risk factor for the need for blood transfusion. In the present study, preoperative hemoglobin was 112.0 ± 14.3 g/L in transfused patients and 122.1 ± 15.8 g/L in those not transfused ($p < 0.001$). Multivariate analysis showed that for every 10 g/L decrease in hemoglobin, patients had a 30% increased risk of blood transfusion ($p < 0.001$). This is consistent with a study by Adunsky and colleagues,¹¹ who studied blood transfusion patterns in 302 patients with hip fractures. They found that a hemoglobin value less than 12 g/dL (120 g/L) increased transfusion risk 5-fold. They suggested that patients with a hemoglobin greater than 12 g/dL (120 g/L) did not require crossmatching preoperatively. Other studies have also identified low preoperative hemoglobin as a risk factor for transfusion.^{12–15}

Our univariate analysis showed that older patients were more likely to receive a blood transfusion, but age was not important in our multivariate analysis, which suggests that other factors explain the same variability in need for transfusion as age but were even more predictive. The average age of patients in our study who received a blood transfusion was 83.1 ± 7.9 years compared with 80.9 ± 9.0 years in patients who did not ($p < 0.003$). Swain and colleagues¹⁶ studied the transfusion requirements for 249 patients with femoral neck fractures and found that patients aged 80 years and older were transfused significantly more than those aged younger than 80 years. Similarly, Dillon and colleagues¹² found that patients older than 75 years were at higher risk of receiving blood transfusion.

Patients receiving a blood transfusion had an average ASA score of 3.4 compared with 3.3 in patients who did not receive a transfusion ($p < 0.001$). This result was significant; however, it is difficult to determine the clinical significance with such a minor difference between scores (0.1). Previous studies examining ASA scores and transfusion requirements did not find any association.^{11,12}

Delay from admission to operation in elderly patients with hip fractures carries significant morbidity and mortality. Delay from admission to operation has been shown to increase mortality,^{17–19} postoperative infections^{19,20} and length of stay in hospital.^{20–22} However, the present study did not find a correlation with delay to operation and transfusion requirements. One would assume that while a patient is awaiting surgery, continued blood loss would be present at the fracture site, leading to greater blood loss and greater risk of transfusion. The equivocal risk of transfusion found in our study may be related to the formation of hematoma around the fracture site. The delay to surgery may have allowed the fracture hematoma to fully stabilize, minimizing the active bleeding before surgery. Intraoperatively, this may have resulted in less surgical blood loss and a lower rate of blood transfusion.

When comparing fracture types, patients with intertrochanteric or subtrochanteric fractures were at an increased risk of requiring a blood transfusion. When compared with femoral neck fractures, there was a 2.37 times greater risk of blood transfusion in patients with an intertrochanteric fracture ($p < 0.001$) and a 4.03 times greater risk in patients with a subtrochanteric fracture ($p < 0.001$). These results are consistent with those of Adunsky and colleagues,¹¹ who found that patients with pertrochanteric fractures were transfused significantly more than patients with subcapital fractures. Swain and colleagues¹⁶ also found an increased transfusion requirement in patients with intertrochanteric fractures compared with those with intracapsular fractures, and Dillon and colleagues¹² reported an increased risk for blood transfusion in patients with pertrochanteric fractures.

When comparing fixation methods for the different fracture types, we found a significant difference in blood transfusion requirements. In patients with femoral neck fractures, 24.8% treated with hemiarthroplasty were transfused compared with 14.3% of patients treated with DHS ($p < 0.001$). This may be partially related to severity of injury, as the more displaced fractures were likely treated with hemiarthroplasty. This finding is consistent with those of a meta-analysis performed by Wang and colleagues,²³ who compared the outcomes of patients who underwent arthroplasty with those of patients who underwent internal fixation for displaced femoral neck fractures. They found greater operative blood loss and increased transfusion requirement in the hemiarthroplasty group. Similarly, Parker and colleagues²⁴ found lower operative blood loss and lower transfusion requirements with internal fixation (using cannulated screws) than with hemiarthroplasty in patients with displaced femoral neck fractures. Similar to femoral neck fractures, intertrochanteric fractures treated with a DHS had a lower risk of transfusion than those treated with an intramedullary device. In our study, 37.9% of intertrochanteric fractures treated with a DHS were transfused compared with 51.6%

treated with an intramedullary device ($p < 0.001$). The literature comparing blood loss in intertrochanteric hip fractures treated with intramedullary nail or DHS is inconclusive. Authors have reported reduced blood loss,²⁵⁻²⁷ increased blood loss²⁸ and no difference^{29,30} in patients with intertrochanteric fractures treated with an intramedullary nail. The variability among studies may be related to differences in fracture severity, as patients with more displaced fractures are generally more likely to be treated with an intramedullary device.

Limitations

There are limitations to this study that should be considered. First, there may be inherent differences in medical comorbidities between the transfused and nontransfused patient samples that were not discerned by this retrospective, observational study. As mentioned, we attempted to stratify by ASA score, but it is difficult to determine if such a small difference in ASA score between groups has any clinical importance. Second, patients were generally given a transfusion based on the criteria of hemoglobin less than 70 g/L or less than 80 g/L with signs/symptoms of anemia. This threshold is consistent with the restrictive strategy group in a study by Carson and colleagues,³¹ who compared restricted and liberal transfusion thresholds. Similar to Carson and colleagues,³¹ we attempted to make the decision to transfuse as uniform as possible; however, there are difficulties with this. Determining signs/symptoms of anemia is subjective, and a lack of uniformity inherently exists among treating physicians. This may have resulted in some inconsistencies in decision to transfuse. Finally, we did not separately analyze patients with stable and unstable intertrochanteric fractures. One would assume that 3- and 4-part intertrochanteric fractures are most likely to cause greater blood loss and more likely to be treated with an intramedullary device. This would potentially affect the results of transfusion requirements between the intramedullary nail and DHS groups. However, previously published literature has failed to find convincing evidence of greater blood loss in patients with unstable fracture patterns.^{32,33}

CONCLUSION

We found a correlation with blood transfusion requirement and sex, preoperative hemoglobin, fracture type and fixation method. The only modifiable risk factor found in the present study was fixation method. Further studies would be beneficial to determine other potential modifiable risk factors. Based on our findings, when considering blood transfusion requirements in isolation, we suggest a potential benefit in using a DHS for intertrochanteric and femoral neck fractures amenable to DHS fixation.

Competing interests: None declared.

Contributors: S. Desai, D. Bryant, A. Lawendy and D.W. Sanders designed the study. S. Desai, K.S. Wood and H. Abdo acquired the data, which S. Desai, J. Marsh, D. Bryant and A. Lawendy analyzed. S. Desai wrote the article, which all authors reviewed and approved for publication.

References

- Dennison E, Mohamed MA, Cooper C. Epidemiology of osteoporosis. *Rheum Dis Clin North Am* 2006;32:617-29.
- Carson JL, Altman DG, Duff A, et al. Risk of bacterial infection associated with allogeneic blood transfusion among patients undergoing hip fracture repair. *Transfusion* 1999;39:694-700.
- Agarwal N, Murphy JG, Cayten CG, et al. Blood transfusion increases the risk of infection after trauma. *Arch Surg* 1993;128:171-6, discussion 176-7.
- Edna TH, Bjerkeset T. Association between blood transfusion and infection in injured patients. *J Trauma* 1992;33:659-61.
- Edna TH, Bjerkeset T. Association between transfusion of stored blood and infective bacterial complications after resection for colorectal cancer. *Eur J Surg* 1998;164:449-56.
- Hill GE, Frawley WH, Griffith KE, et al. Allogeneic blood transfusion increases the risk of postoperative bacterial infection: a meta-analysis. *J Trauma* 2003;54:908-14.
- Koval KJ, Rosenberg AD, Zuckerman JD, et al. Does blood transfusion increase the risk of infection after hip fracture? *J Orthop Trauma* 1997;11:260-5, discussion 265-6.
- Engoren M, Mitchell E, Perring P, et al. The effect of erythrocyte blood transfusion on survival after surgery for hip fracture. *J Trauma* 2008;65:1411-5.
- Vincent JL, Baron JF, Reinhart K, et al. Anemia and blood transfusion in critically ill patients. *JAMA* 2002;288:1499-507.
- Report from Department of Health and Human Services. *The 2009 National Blood Collection And Utilization Survey Report*. Washington (DC): Department of Health and Human Services, Office of the Assistant Secretary for Health; 2011.
- Adunsky A, Lichtenstein A, Mizrahi E, et al. Blood transfusion requirements in elderly hip fracture patients. *Arch Gerontol Geriatr* 2003;36:75-81.
- Dillon MF, Collins MB, Rice J, et al. Preoperative characteristics identifying patients with hip fractures at risk of transfusion. *Clin Orthop Relat Res* 2005; (439):201-6.
- Kurdy NM, Hokan R. A cross-matching policy for fractures of the proximal third of the femur. *Injury* 1993;24:521-4.
- Levi N. Blood transfusion requirements in intracapsular femoral neck fractures. *Injury* 1997;27:709-11.
- Robbins J, Steingold RF. Blood use in urgent operation for patients with fractures of the femoral neck. *Injury* 1986;17:265-6.
- Swain DG, Nightingale PG, Patel JV. Blood transfusion requirements in femoral neck fracture. *Injury* 2000;31:7-10.
- Zuckerman JD, Skovron ML, Koval KJ, et al. Postoperative complications and mortality associated with operative delay in older patients who have a fracture of the hip. *J Bone Joint Surg Am* 1995;77:1551-6.
- Moran CG, Wenn RT, Sikand M, et al. Early mortality after hip fracture: Is delay before surgery important? *J Bone Joint Surg Am* 2005;87:483-9.
- Simunovic N, Devereaux PJ, Sprague S, et al. Effect of early surgery after hip fracture on mortality and complications: systematic review and meta-analysis. *CMAJ* 2010;182:1609-16.
- Verbeek DO, Ponsen KJ, Goslings JC, et al. Effect of surgical delay on outcome in hip fracture patients: a retrospective multivariate analysis of 192 patients. *Int Orthop* 2008;32:13-8.
- Rogers FB, Shackford SR, Keller MS. Early fixation reduces morbidity and mortality in elderly patients with hip fractures from

- low-impact falls. *J Trauma* 1995;39:261-5.
22. Orosz GM, Magaziner J, Hannan EL, et al. Association of timing of surgery for hip fracture and patient outcomes. *JAMA* 2004; 291:1738-43.
 23. Wang J, Jiang B, Marshall RJ, et al. Arthroplasty of internal fixation for displaced femoral neck fractures: Which is the optimal alternative for elderly patients? A meta-analysis. *Int Orthop* 2009;33:1179-87.
 24. Parker MJ, Khan JK, Crawford J, et al. Hemiarthroplasty versus internal fixation for displaced intracapsular hip fractures in the elderly. *J Bone Joint Surg Br* 2002;84:1150-5.
 25. Giancola R, Antonini G, Rose GD, et al. Percutaneous compression plating versus gamma nail for the treatment of pertrochanteric hip fractures. *Strategies Trauma Limb Reconstr* 2008;3:9-14.
 26. Dujardin FH, Benewz C, Polle G, et al. Prospective randomized comparison between a dynamic hip screw and a mini-invasive static nail in fractures of the trochanteric area: preliminary results. *J Orthop Trauma* 2001;15:401-6.
 27. Little NJ, Verma V, Fernando C, et al. A prospective trial comparing the Holland nail with the dynamic hip screw in the treatment of intertrochanteric fractures of the hip. *J Bone Joint Surg Br* 2008; 90:1073-8.
 28. Foulongne E, Gilleron M, Roussignol X, et al. Mini-invasive nail versus DHS to fix pertrochanteric fractures: a case-control study. *Orthop Traumatol Surg Res.* 2009;95:592-8.
 29. Varela-Egocheaga JR, Iglesias-Colao R, Suarez-Suarez MA, et al. Minimally invasive osteosynthesis in stable trochanteric fractures: a comparative study between Gotfried percutaneous compression plate and gamma 3 intramedullary nail. *Arch Orthop Trauma Surg* 2009;129:1401-7.
 30. Harrington P, Nihal A, Singhanian AK, et al. Intramedullary hip screw versus sliding hip screw for unstable intertrochanteric femoral fractures in the elderly. *Injury* 2002;33:23-8.
 31. Carson JL, Terrin ML, Noveck H, et al. Liberal or restrictive transfusion in high-risk patients after hip surgery. *N Engl J Med* 2011;365:2453-62.
 32. Lavini F, Renzi-Brivio L, Aulisa R, et al. The treatment of stable and unstable proximal femoral fractures with a new trochanteric nail: results of a multicentre study with the Veronail. *Strategies Trauma Limb Reconstr* 2008;3:15-22.
 33. Leung KS, So WS, Shen WY, et al. Gamma nails and dynamic hip screws for pertrochanteric fractures. A randomised prospective study in elderly patients. *J Bone Joint Surg Br* 1992;74:345-51.

Essential reading

from the Canadian Medical Association

- *CMAJ*
- *Canadian Journal of Surgery*
- *Journal of Psychiatry & Neuroscience*

For information contact

CMA Member Service Centre
888 855-2555
cmamsc@cma.ca

