

4-1-2008

## **Meta-analysis of alexithymia in posttraumatic stress disorder.**

Paul A Frewen

David J A Dozois

Richard W J Neufeld

Ruth A Lanius

Follow this and additional works at: <https://ir.lib.uwo.ca/psychologypub>



Part of the [Psychology Commons](#)

---

### **Citation of this paper:**

Frewen, Paul A; Dozois, David J A; Neufeld, Richard W J; and Lanius, Ruth A, "Meta-analysis of alexithymia in posttraumatic stress disorder." (2008). *Psychology Publications*. 223.

<https://ir.lib.uwo.ca/psychologypub/223>

Running Head: "ALEXITHYMIA IN PTSD"

Word Count:

Text, Footnotes, Table 1 Note: 1640

References: 790

Tables (1): 2.0 pages x 200 words = 400 words

Total Word Count = 2830 words

**Meta-analysis of Alexithymia in Posttraumatic Stress Disorder**

Paul A. Frewen<sup>1</sup>, David J. A. Dozois<sup>1,2</sup>, Richard W. J. Neufeld<sup>1,2,3</sup>, & Ruth A. Lanius<sup>2,3</sup>

Departments of Psychology<sup>1</sup>, Psychiatry<sup>2</sup>, and Neuroscience<sup>3</sup>

The University of Western Ontario

Acknowledgment: The authors would like to acknowledge *The Canadian Institutes of Health Research* (Grant #MOP49543 to R.A.L.), the *Ontario Mental Health Foundation* (Grant #M931D5 to R.A.L.; Grant #X0180A27 to D.J.A.D.), the *Canadian Psychiatric Research Foundation* (Grant #MA94A7 to R.A.L.), and the *Social Sciences & Humanities Research Council of Canada* (Grant # R2180A08 to R.W.J.N., and in the form of a Canada Graduate Scholarship to P.A.F.), for their support.

Corresponding Author: David J. A. Dozois, PhD, University of Western Ontario, London, Ontario, Canada, N6A 5C2, Email: [dzois@uwo.ca](mailto:dzois@uwo.ca), Tel: (519) 661-2111 x84678.

### **Abstract**

The present meta-analytic review investigated the prevalence of alexithymia in individuals with posttraumatic stress disorder (PTSD). A large effect size was found associating PTSD with alexithymia. Effect sizes were higher in studies of male combat PTSD samples in comparison with studies of other PTSD samples. Clinical and research directions are discussed.

**Key words:** Alexithymia, Emotional Awareness, Posttraumatic Stress Disorder, Meta-Analysis.

## Meta-analysis of Alexithymia in Posttraumatic Stress Disorder

Disturbances in emotional experience and expression are often present in individuals with posttraumatic stress disorder (PTSD), and several theories have been put forward to account for these disturbances (reviews by Brewin & Holmes, 2003; Dalgleish, 2004). One form of emotional-experiential and -expressive disturbance is alexithymia, a psychological construct defined by difficulties in identifying and labeling emotional feelings and a tendency toward externally-oriented thinking (Sifneos, 1973; Taylor, Bagby, & Parker, 1997).

Since the seminal clinical observations of Krystal (1968, Krystal & Krystal, 1988), empirical studies have confirmed an increased prevalence of alexithymia in PTSD samples. The present article meta-analyzes these studies to evaluate the size of these effects, as well as to test whether they vary by trauma type and gender.

### Method

#### *Literature Review*

The *PsychINFO* and *PubMed* databases were searched using the keywords '*alexithymia*', '*posttraumatic stress disorder/PTSD*' and '*trauma*'. Articles referenced by studies identified by these searches were also reviewed. To the authors' knowledge, the present review encompasses all English-written published studies that have reported *alexithymia scores* in a PTSD sample up to July 2007 other than that of Cloitre et al. (1997), which could not be included due to their use of non-parametric statistics. Study authors were contacted to provide information that was unavailable in the published manuscripts. Studies of PTSD symptoms in non-diagnosed samples (e.g., McCaslin et al., 2006) were not included to insure a minimum level of PTSD severity. '*Alexithymia scores*' in this context refer to scores on psychometric scales of alexithymia that are widely recognized within the literature as such, most notably, versions of the *Toronto Alexithymia Scale* (see Taylor et al., 1997 for review). This measure has

demonstrated strong psychometric properties, particularly the 20-item version (Bagby, Parker, Taylor, 1994; Parker, Taylor, & Bagby, 2003).

#### *Meta-analytic calculations*

Weighted-mean effect sizes ( $d$ ) were calculated following Hedges and Olkin (1985) with diagnosed PTSD samples compared with study sample control groups (referred to hereafter as  $d_{SAMP}$ ) as well as population norms for the respective psychometric alexithymia scale used (referred to hereafter as  $d_{POP}$ ). The population norms used for the  $d_{POP}$  calculations were the following: *TAS-20* (from: Parker, Taylor, & Bagby, 2003, Table 3):  $\mu_{M(N=868)} = 47.30$ ,  $\sigma = 11.32$ ,  $\mu_{F(N=1065)} = 44.15$ ,  $\sigma = 11.19$ ,  $\mu_{M\&F(N=1933)} = 45.57$ ,  $\sigma = 11.35$ ; *TAS-23* (from: Taylor, Bagby, Parker, 1992, Table 2):  $\mu_{M(N=55)} = 59.76$ ,  $\sigma = 11.00$ ,  $\mu_{F(N=84)} = 59.11$ ,  $\sigma = 10.44$ ,  $\mu_{M\&F(N=139)} = 59.40$ ,  $\sigma = 10.70$ ; *TAS-26* (from: Taylor, Ryan, & Bagby, 1985, Table 3):  $\mu_{M(N=126)} = 63.32$ ,  $\sigma = 10.90$ ,  $\mu_{F(N=370)} = 61.11$ ,  $\sigma = 11.33$ ,  $\mu_{M\&F(N=542)} = 61.80$ ,  $\sigma = 11.27$ .

$d$  values were classified following Cohen (1988) as follows: small ( $\geq .20$ ), medium ( $\geq .50$ ), and large ( $\geq .80$ ). Significant heterogeneity evident in the combined effect-sizes was parsed a priori by examining trauma-type and gender as moderators. For  $d_{SAMP}$ , Rosenthal's (1979) corresponding 'Fail Safe N' (FSN) was calculated, which is interpreted as the number of studies with null effects ( $d = 0$ ) required to render  $d_{SAMP}$  zero.

## **Results**

### *Study Descriptive Information*

Twelve studies were located in the literature searches that met study inclusion criteria totaling a PTSD sample of 1095. Study sample control groups in these studies varied as follows: Hyer et al. (1990) compared PTSD veterans to general psychiatric outpatient veterans; Phan et al. (2006) compared veterans with versus without PTSD; Zeitlin et al. (1989) compared combat veterans with PTSD to healthy non-combat controls; Söngergaard & Theorell compared refugees with versus without PTSD; Yehuda et al. (1997) compared holocaust survivors with versus without PTSD; Zeitlin et al. (1993) compared rape victims

with versus without PTSD; and Zlotnick et al. (2001) compared individuals with PTSD to other general psychiatric outpatient samples.

### *Effect Sizes*

Table 1 reports descriptive information and effect sizes for the 12 studies. The mean effect size when PTSD samples were compared with study samples was  $d_{SAMP} = 0.80$  ( $SE = 0.10$ , 95% C.I. =  $.60 \leq d_{SAMP} \leq 1.00$ ), a large effect. The number of studies with null effects that would be required to render this effect size to zero (FSN) = 370. The mean effect size when PTSD samples were compared with normative values for non-psychiatric populations was  $d_{POP} = 1.20$  ( $SE = 0.04$  95% C.I. =  $1.10 \leq d_{POP} \leq 1.30$ ), also a large effect.

### *Moderators*

When calculated relative to non-psychiatric population norms, combat veteran PTSD samples displayed a larger mean effect size than non-combat veteran PTSD samples:  $d_{POP} = 1.71$  ( $SD = 0.55$ ) vs.  $d_{POP} = 1.02$  ( $SD = 0.41$ ), respectively,  $t(9) = 2.37$ ,  $p = .04$  (2-tailed). Similarly, the ratio of males to females within each study PTSD sample significantly correlated with the  $d_{POP}$  value,  $r(9) = .70$ ,  $p = .03$  (2-tailed).

## **Discussion**

This meta-analysis supports the hypothesis that individuals with PTSD experience symptoms of alexithymia. Alexithymic symptomatology was particularly characteristic of males with combat-related PTSD although individuals with non-combat related PTSD also displayed clinically-elevated alexithymia. Future research will need to take gender differences in alexithymia within the normal population into account in tests of the effects of alexithymia in PTSD in relation to gender and trauma-type, as in the present review these factors were confounded (combat samples were 100% male, non-combat samples were 20% male).

These findings may have significant clinical implications. For example, it has been proposed that certain individuals with PTSD may require modularized interventions not only for reprocessing traumatic memories but also for ameliorating more general deficits in emotional awareness and regulation such as

alexithymia symptoms (e.g., Cloitre, Cohen, & Koenen, 2006). However, it remains to be shown whether individual differences in alexithymia represent negative prognostic indicators in PTSD.

Future studies should assess the impact alexithymia may have on emotional experience and expression in response to emotion-activating stimuli in PTSD (Litz, Orsillo, & Kaloupek, 2000; Miller & Litz, 2004; Orsillo, Batten, Plumb, Luterek, & Roessner, 2004; Wagner, Roemer, Orsillo, & Litz, 2003). It may be helpful for future studies of this type to employ self-referential paradigms, detailed assessments of participants' emotional experience, and non-self-report measurements of emotional response (e.g., psychophysiology, neuroimaging).

The present review has limitations. For example, the population norms used for calculation of the  $d_{POP}$  effect sizes may have been biased by demographic and/or other non-specific differences between the PTSD sample and normative group that were not consistently reported by study authors. The  $d_{SAMP}$  and  $d_{POP}$  values found for certain studies also varied suggesting that the magnitude of the effects may depend on the control group studied, particularly if the control group is another psychiatric group rather than a healthy sample (although Hyer et al., 1990 and Zlotnick et al., 2001 showed PTSD samples scored higher than other psychiatric outpatients). The fact that participants in the studies comprising this meta-analysis were not randomly sampled from the PTSD population also means that the results may not generalize beyond the specific samples studied. Additionally, studies rarely uniquely compared the distinctive sub-dimensions of alexithymia, making it unclear whether individuals with PTSD are more likely to display all characteristics of alexithymia or only specific sub-dimensions<sup>2</sup>. Finally, the degree of trauma exposure has typically not been evaluated as a moderating variable, and in turn the timing of trauma exposure within emotional and cognitive development<sup>3</sup>. However, it is conceivable that symptoms of alexithymia relate to developmental trauma exposure independent of PTSD symptoms. For example, developmental models such as that of Lane and Schwartz (1987; Lane et al., 1997) posit that cognitive abilities with respect to emotional awareness and expression undergo developmental progression within the context of attachment

relationships, and traumatic experiences and neglect occurring during childhood may disturb this natural development, culminating in severe cases in alexithymia in adulthood. Consistent with this proposal Frewen et al. (2007) and Zlotnick et al. (2001) found that alexithymia symptoms correlated with retrospectively-reported experiences of childhood emotional neglect.

In conclusion, the present meta-analytic results provide support for an association between PTSD and alexithymia. It is nevertheless important not to misconstrue the present results as suggesting that *all* PTSD subjects are alexithymic. Instead, it is hoped that this review will spark continuing clinical and research interest into the complex disturbances in emotional experience and expression, such as of alexithymia, that may be present in certain individuals with PTSD and complex trauma histories.



## References

- Badura, A. S. (2003). Theoretical and empirical exploration of the similarities between emotional numbing in posttraumatic stress disorder and alexithymia. *Anxiety Disorders, 17*, 349-360.
- Bagby, R. M., Taylor, G. J., & Parker, J. D. A. (1994a). The twenty-item Toronto Alexithymia Scale-1: Item selection and cross-validation of the factor structure. *Journal of Psychosomatic Research, 38*, 23-32.
- Brewin, C. R., & Holmes, E. A. (2003). Psychological theories of posttraumatic stress disorder. *Clinical Psychology Review, 23*, 1041-1065.
- Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences, 2<sup>nd</sup> Ed.* Hillsdale, NJ: Lawrence Erlbaum).
- Cloitre, M., Cohen, L. R., & Koenen, K. C. (2006). *Treating Survivors of Childhood Abuse: Psychotherapy for the Interrupted Life.* New York, NY: Guilford Press.
- Cloitre, M., Scarvalone, P., & Difede, J. (1997). Posttraumatic stress disorder, self- and interpersonal dysfunction among sexually retraumatized women. *Journal of Traumatic Stress, 10*, 437-452.
- Dalgleish, T. (2004). Cognitive approaches to posttraumatic stress disorder: The evolution of multi-representational theorizing. *Psychological Bulletin, 130*, 228-260.
- Frewen, P. A., Lanius, R. A., Dozois, D. J. A., Neufeld, R. W. J., Pain, C., Hopper, J. W., Densmore, M., & Stevens, T. K. (in press). Clinical and neural correlates of alexithymia in PTSD. *Journal of Abnormal Psychology.*
- Hedges, I. V., & Olkin, I. (1985). *Statistical Methods for Meta-Analysis.* New York, NY: Academic Press.
- Hyer, L, Woods, G., Summers, M. N., Boudewyns, P., & Harrison, W. R. (1990). Alexithymia among Vietnam veterans with posttraumatic stress disorder. *Journal of Clinical Psychiatry, 51*, 243-247.
- Kleiger, J. H., & Kinsman, R. A. (1980). The development of an MMPI alexithymia scale. *Psychotherapy & Psychosomatics, 34*, 17-24.
- Krystal, H. (1968). *Massive Psychic Trauma.* New York, NY: International University Press.

- Krystal, H., & Krystal, J. H. (1988). *Integration and self-healing: Affect, trauma, alexithymia*. Hillsdale, NJ, England: Analytic Press, Inc.
- Litz, B. T. & Gray, M. J. (2002). Emotional numbing in posttraumatic stress disorder: Current and future directions. *Australian and New Zealand Journal of Psychiatry, 36*, 198-204.
- McCaslin, S.E., Metzler, T.J., Best, S.R., Liberman, A., Weiss, D.S., Fagan, J., & Marmar, C.R. (2006). Alexithymia and PTSD symptoms in urban police officers: Cross-sectional and prospective findings. *Journal of Traumatic Stress, 19*, 361-373.
- Monson, C. M., Price, J. L., Rodriguez, B. F., Ripley, M. P., & Warner, R. A. (2004). Emotional deficits in military-related PTSD: An investigation of content and process disturbances. *Journal of Traumatic Stress, 17*, 275-279.
- Orsillo, S. M., Batten, S. V., Plumb, J. C., Luterek, J. A., & Roessner, B. M. (2004). An experimental study of emotional responding in women with posttraumatic stress disorder related to interpersonal violence. *Journal of Traumatic Stress, 17*, 241-248.
- Parker, J. D. A., Taylor, G. J., & Bagby, R. M. (2003). The 20-item Toronto Alexithymia Scale III: Reliability and factorial validity in a community population. *Journal of Psychosomatic Research, 55*, 269-275.
- Phan, K. L., Britton, J. C., Taylor, S. F., Fig, L. M., & Liberzon, I. (2006). Corticolimbic blood flow during nontraumatic emotional processing in posttraumatic stress disorder. *Archives of General Psychiatry, 63*, 184-192.
- Ramirez, S. M., Glover, H., Ohlde, C., Mercer, R., Goodnick, P., Hamlin, C., & Perez-Rivera, M. (2001). Relationship of Numbing to Alexithymia, Apathy, and Depression. *Psychological Reports, 88*, 189-200.
- Sifneos, P. E. (1973). The prevalence of 'alexithymic' characteristics in psychosomatic patients. *Psychotherapy & Psychosomatics, 22*, 255-262.
- Söndergaard, H. P., & Theorell, T. (2004). Alexithymia, emotions and PTSD: Findings from a longitudinal study of refugees. *Nordic Journal of Psychiatry, 58*, 185-191.

- Taylor, G. J., Bagby, R. M., & Parker, J. D. A. (1992). The Revised Toronto Alexithymia Scale: Some reliability, validity, and normative data. *Psychotherapy & Psychosomatics*, *57*, 34-41.
- Taylor, G. J., Bagby, R. M., & Parker, J. D. A. (1997). *Disorders of Affect Regulation: Alexithymia in Medical and Psychiatric Illness*. Cambridge, UK: Cambridge University Press.
- Taylor, G. J., Ryan, D., & Bagby, R. M. (1985). Toward the development of a new self-report alexithymia scale. *Psychotherapy & Psychosomatics*, *44*, 191-199.
- Wagner, A. W., Roemer, L., Orsillo, S. M., & Litz, B. T. (2003). Emotional experiencing in women with posttraumatic stress disorder: Congruence between facial expressivity and self-report. *Journal of Traumatic Stress*, *16*, 67-75.
- Yehuda, R., Steiner, A., Kahana, B., Binder-Brynes, K., Southwick, S. M., Zelman, S., & Giller, E. L. (1997). Alexithymia in holocaust survivors with and without PTSD. *Journal of Traumatic Stress*, *10*, 93-100.
- Zeitlin, S. B., Lane, R. D., O'Leary, D. S., & Schrift, M. J. (1989). Interhemispheric transfer deficit and alexithymia. *American Journal of Psychiatry*, *146*, 1434-1439.
- Zeitlin, S. B., McNally, R. J., & Cassiday, K. L. (1993). Alexithymia in victims of sexual assault: An effect of repeated traumatization? *American Journal of Psychiatry*, *150*, 661-663.
- Zlotnick, C., Mattia, J. L., & Zimmerman, M. (2001). The relationship between posttraumatic stress disorder, childhood trauma and alexithymia in an outpatient sample. *Journal of Traumatic Stress*, *14*, 177-188.

## Footnotes

<sup>1</sup> The corresponding value for  $d_{POP}$  is not presented due to it being artificially elevated due to the large control group  $n$ -size.

<sup>2</sup> Sondergaard & Theorell found refugees with versus without PTSD to differ only on the *Difficulty Identifying Feelings* factor of the TAS-20, whereas Monson et al. (2004) found that *Externally-Oriented Thinking* scores were the only significant individual predictor of CAPS total scores when each of the three TAS-20 subscale scores were entered individually in a multiple regression analysis.

<sup>3</sup> Zietlin et al. (1993) found that, whereas rape victims with versus without PTSD did not differ in alexithymia, rape victims were more alexithymic than were individuals who had not experienced rape. Moreover, rape victims who had experienced more than one assault were more alexithymic than were victims of only a single assault. In contrast, neither Yeduda et al. (1997) nor Badura (2003) found that trauma exposure correlated with degree of alexithymia.

## Table 1 Note

Note: M = Male, F = Female, SE = Standard Error of the Effect Size, TAS = *Toronto Alexithymia Scale*, MMPI = *Minnesota Multiphasic Personality Inventory*, <sup>1</sup> see Kleiger & Kinsman (1980), low scores reflect greater alexithymia therefore sign of  $d$ -values was reversed; n-av = data not available due to lack of a control group.

**Table 1****Study descriptive statistics and effect sizes for self-report studies of alexithymia**

#	Reference	Trauma Type	Control Group	Scale	$n_{PTSD}$ (M:F)	$M_{PTSD}$ (SD)	$n_{CONT}$ (M:F)	$M_{CONT}$ (SD)	$d_{SAMP}$ (SE)	$d_{POP}$ (SE)
1	Badura (2003)	Combat	None	TAS-23	274 (274:0)	75.14 (11.67)	n-av	n-av	n-av	1.40 (0.2)
2	Monson et al. (2004)	Combat	None	TAS-20	85 (85:0)	74.35 (9.55)	n-av	n-av	n-av	2.40 (0.1)
3	Ramirez et al. (2001)	Combat	None	TAS-20	353 (353:0)	72.12 (12.04)	n-av	n-av	n-av	2.20 (0.1)
4	Zeitlin et al. (1989)	Combat	Non- psychiatric	TAS-26	25 (25:0)	78.20 (16.60)	10 (10:0)	54.60 (6.60)	3.27 (0.9)	1.37 (0.2)
5	Phan et al. (2006)	Combat	Non- psychiatric	TAS-20	16 (16:0)	61.30 (19.90)	15 (15:0)	52.50 (7.90)	1.59 (0.5)	1.20 (0.1)
6	Hyer et al. (1990)	Combat	Psychiatric Alexithymia <sup>1</sup>	MMPI	76 (76:0)	8.90 (2.50)	75 (75:0)	10.40 (3.20)	0.47 (0.2)	n-av

Alexithymia in PTSD 13

7	Cloitre et al. (2002)	Childhood Abuse	None	TAS-20	58 (0:58)	55.43 (11.77)	n-av	n-av	n-av	1.00 (0.1)
8	Söndergaard & Theorell (2004)	Refugees	Non- psychiatric	TAS-20	32 (23:9)	61.41 (12.57)	54 (31:23)	55.44 (12.17)	0.48 (0.2)	1.40 (0.1)
9	Yehuda et al. (1997)	Holocaust	Non- psychiatric	TAS-20	30 (9:21)	56.93 (14.37)	26 (13:13)	47.15 (12.69)	0.93 (0.3)	1.00 (0.1)
10	Zeitlin et al. (1993)	Rape	Non- psychiatric	TAS-26	12 (1:11)	64.67 (11.62)	12 (0:12)	65.50 (14.66)	-0.09 (.4)	0.25 (0.3)
11	Frewen et al. (2007)	Mixed	Non- psychiatric	TAS-20	105 (12:93)	59.38 (13.67)	45 (16:29)	35.39 (8.86)	2.71 (0.3)	1.20 (0.1)
12	Zlotnick et al. (2001)	Mixed	Psychiatric	TAS-26	29 (n-av)	76.69 (14.43)	223 (n-av)	70.20 (12.45)	0.52 (0.2)	1.32 (0.2)