

7-16-2020

Teaching NeuroImages: Reversible neuroimaging findings during treatment of infantile spasms with vigabatrin

David Dongkyung Kim

Amit Kumar Sharma

Manas Sharma

Andrea Andrade
aandrad6@uwo.ca

Follow this and additional works at: <https://ir.lib.uwo.ca/paedpub>



Part of the [Pediatrics Commons](#)

Citation of this paper:

Kim, David Dongkyung; Sharma, Amit Kumar; Sharma, Manas; and Andrade, Andrea, "Teaching NeuroImages: Reversible neuroimaging findings during treatment of infantile spasms with vigabatrin" (2020). *Paediatrics Publications*. 527.
<https://ir.lib.uwo.ca/paedpub/527>

Teaching NeuroImages: Reversible neuroimaging findings during treatment of infantile spasms with vigabatrin


David Dongkyung Kim, MD, Amit Kumar Sharma, MD, Manas Sharma, MD, and Andrea Andrade, MD

Neurology® 2020;95:e2314-e2315. doi:10.1212/WNL.0000000000010370

Correspondence

Dr. Kim
david.kim@medportal.ca

Figure Reversible abnormal MRI findings during treatment of infantile spasms with vigabatrin



Diffusion-weighted imaging showed symmetrical signal change in bilateral thalami (A, B), globus pallidi (B, C), cerebral peduncles (D), central tegmental tracts (D-F), and dentate nuclei (F). Apparent diffusion coefficient maps showed corresponding changes (not pictured). Follow-up imaging 4 months later showed resolution of the changes seen earlier (not pictured).

A 10-month-old boy with global developmental delay presented to clinic with a few months of infantile spasms occurring multiple times a day. His seizures continued despite vigabatrin (dosed at 133 mg/kg/d), levetiracetam, and steroid therapy. On vigabatrin, routine follow-up MRI showed abnormal signal change (figure), which may occur in 30.9% of patients.¹ Risk is associated with a high peak dose but not cumulative.² These findings are largely asymptomatic although rarely patients can present with hyperkinetic disorders.² The imaging findings resolved on 4-month follow-up after tapering vigabatrin. At 18 months of age, the patient continues to have 1 seizure every 2 weeks.

MORE ONLINE

→ **Teaching slides**
links.lww.com/WNL/B176

From the Department of Clinical Neurological Sciences (D.D.K., A.K.S.), Division of Neuroradiology, Department of Radiology (M.S.), and Division of Pediatric Neurology, Department of Pediatrics (A.A.), Western University, London, Canada.

Go to Neurology.org/N for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

Study funding

No targeted funding reported.

Disclosure

The authors report no disclosures relevant to the manuscript. Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures.

Appendix Authors

Name	Location	Contribution
David Dongkyung Kim, MD	Western University, London, Canada	Involved in the concept and writing of the manuscript
Amit Kumar Sharma, MD	Western University, London, Canada	Involved in the concept and writing of the manuscript

Appendix *(continued)*

Name	Location	Contribution
Manas Sharma, MD	Western University, London, Canada	Involved in critical revision and study supervision
Andrea Andrade, MD	Western University, London, Canada	Involved in critical revision and study supervision

References

1. Dracopoulos A, Widjaja E, Raybaud C, Westall CA, Snead OC III. Vigabatrin-associated reversible MRI signal changes in patients with infantile spasms. *Epilepsia* 2010;51:1297–1304.
2. Hussain SA, Tsao J, Li M, et al. Risk of vigabatrin-associated brain abnormalities on MRI in the treatment of infantile spasms is dose-dependent. *Epilepsia* 2017;58:674–682.

Neurology[®]

Teaching NeuroImages: Reversible neuroimaging findings during treatment of infantile spasms with vigabatrin

David Dongkyung Kim, Amit Kumar Sharma, Manas Sharma, et al.
Neurology 2020;95:e2314-e2315 Published Online before print July 16, 2020
DOI 10.1212/WNL.0000000000010370

This information is current as of July 16, 2020

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/95/16/e2314.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/95/16/e2314.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Epilepsy/Seizures http://n.neurology.org/cgi/collection/all_epilepsy_seizures All Pediatric http://n.neurology.org/cgi/collection/all_pediatric All Toxicology http://n.neurology.org/cgi/collection/all_toxicology Infantile spasms http://n.neurology.org/cgi/collection/infantile_spasms MRI http://n.neurology.org/cgi/collection/mri
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2020 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

