1988

Ultrastructure And Atpase Nature Of Polar Membrane In Campylobacter Jejuni

Frances Marguerite Brock

Follow this and additional works at: <https://ir.lib.uwo.ca/digitizedtheses>

Recommended Citation

Brock, Frances Marguerite, "Ultrastructure And Atpase Nature Of Polar Membrane In Campylobacter Jejuni" (1988). *Digitized Theses*. 1691.
<https://ir.lib.uwo.ca/digitizedtheses/1691>

This Dissertation is brought to you for free and open access by the Digitized Special Collections at Scholarship@Western. It has been accepted for inclusion in Digitized Theses by an authorized administrator of Scholarship@Western. For more information, please contact tadam@uwo.ca, wlsadmin@uwo.ca.

National Library
of Canada

Bibliothèque nationale
du Canada

Canadian Theses Service

Service des thèses canadiennes

Ottawa, Canada
K1A 0N4

NOTICE

The quality of this microform is heavily dependent upon the quality of the original thesis submitted for microfilming. Every effort has been made to ensure the highest quality of reproduction possible.

If pages are missing, contact the university which granted the degree.

Some pages may have indistinct print especially if the original pages were typed with a poor typewriter ribbon or if the university sent us an inferior photocopy.

Previously copyrighted materials (journal articles, published tests, etc.) are not filmed.

Reproduction in full or in part of this microform is governed by the Canadian Copyright Act, R.S.C. 1970, c. C-30.

AVIS

La qualité de cette microforme dépend grandement de la qualité de la thèse soumise au microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

S'il manque des pages, veuillez communiquer avec l'université qui a conféré le grade.

La qualité d'impression de certaines pages peut laisser à désirer, surtout si les pages originales ont été dactylographiées à l'aide d'un ruban usé ou si l'université nous a fait parvenir une photocopie de qualité inférieure.

Les documents qui font déjà l'objet d'un droit d'auteur (articles de revue, tests publiés, etc.) ne sont pas microfilmés.

La reproduction, même partielle, de cette microforme est soumise à la Loi canadienne sur le droit d'auteur, S.R.C. 1970, c. C-30.

ULTRASTRUCTURE AND ATPASE NATURE OF POLAR MEMBRANE IN
CAMPYLOBACTER JEJUNI

by

Frances Marguerite Brock

Department of Microbiology and Immunology

Submitted in partial fulfillment
of the requirements of the degree of
Doctor of Philosophy

Faculty of Graduate Studies
University of Western Ontario

London, Ontario

August 1987

© Frances Marguerite Brock 1988

Permission has been granted to the National Library of Canada to microfilm this thesis and to lend or sell copies of the film.

The author (copyright owner) has reserved other publication rights, and neither the thesis nor extensive extracts from it may be printed or otherwise reproduced without his/her written permission.

L'autorisation a été accordée à la Bibliothèque nationale du Canada de microfilmer cette thèse et de prêter ou de vendre des exemplaires du film.

L'auteur (titulaire du droit d'auteur) se réserve les autres droits de publication; ni la thèse ni de longs extraits de celle-ci ne doivent être imprimés ou autrement reproduits sans son autorisation écrite.

ISBN 0-315-40759-X

This thesis is dedicated to
Dr. C.F. Robinow whose teaching
and enthusiasm for microbiology
led me down this road.

Abstract

The ultrastructure of polar membrane in Campylobacter jejuni was examined and was shown to line the cytoplasmic membrane at the poles of the cell like an internal cylindrical cap. This was not found to be the case in Aquaspirillum serpens MW5, where the polar membrane was found in patches lining the cytoplasmic membrane near the flagellar tufts. Polar membrane in C. jejuni was found at the site of septum formation, although this was not a consistent result.

The component parts of polar membrane (stalks and knobs) were found to be consistent with previous reports of this structure, although their dimensions (stalks, 2-3 nm in diameter, and knobs, 5-6 nm in diameter) were smaller than those described for Ectothiorhodospira mobilis (stalks 3-4 nm in diameter, and knobs, 8-9 nm in diameter).

The association of polar membrane with the flagellar basal complex was examined and it was found that this structure encroaches on the lateral margins of the basal complex to varying degrees, but was never found to cover the area where the M ring was located.

An area containing fewer ribosomes than the rest of

cytoplasm was found in the location of the polar membrane in *C. jejuni* and *A. serpens* MW5. *C. jejuni* was found to have a densely staining spherical inclusion body, probably consisting of phosphate moieties, in the cytoplasm beneath the polar membrane structure.

Polar membrane in *C. jejuni* has been visualized on membrane vesicles. It is composed of dough-nut shaped particles 5-6 nm in diameter, arranged in an hexagonal array. This structure was stabilized on the membranes by a high ionic strength buffer in the presence of 2-mercaptoethanol.

C. jejuni sheds its flagella and varying proportions of the poles of the cell late in the growth cycle, resulting in the production of very small flagellated structures 0.1 to 0.3 um in diameter. Electron microscopy revealed that these structures were minicells possessing outer membrane, cytoplasmic membrane, flagellar basal complex, and polar membrane; nucleoplasms were not seen. The initial event in the formation of these minicells involved a constriction of the cytoplasmic membrane, segregating the polar regions of the cell. The peptidoglycan layer of the cell wall was not visible, but was presumed to lyse at the separation site of minicell formation, and to reform or remain intact along the

main length of the cell because the rods did not spheroplast. Finally, rupture and resealing of the outer membrane component of the wall resulted in the release of fully enclosed minicells and nonflagellated rods.

Histochemical staining of whole cells of C. jejuni indicated localized Mg^{2+} -ATPase activity at the poles of the cells. Ca^{2+} -ATPase activity was randomly distributed throughout the cell. Histochemical staining for Mg^{2+} -ATPase and Ca^{2+} -ATPase activity was performed in Escherichia coli K12, Bacillus cereus, Micrococcus luteus, Deinococcus radiodurans, and A. serpens MW5 for comparison with that found in C. jejuni. A. serpens MW5, which also possessed polar membrane, showed localized Ca^{2+} -ATPase activity at the poles of the cells and Mg^{2+} -ATPase activity randomly distributed throughout the cell.

An ATPase with distinctive properties was isolated and purified from C. jejuni giving a specific activity of approximately 0.3 units/mg of protein. Electron microscopy showed dough-nut shaped particles 5-6 nm in diameter. Non-dissociating and sodium dodecyl sulfate polyacrylamide gel electrophoresis of the purified enzyme revealed, respectively, a single band with ATPase activity and a molecular weight of ca. 75,000 daltons. The enzyme was cold labile and activity was abolished by trypsin.

Dicyclohexylcarbodiimide inhibited the membrane-bound form of the enzyme, but did not inhibit the soluble form. Oligomycin had no inhibitory activity on either form of the enzyme. The enzyme specifically hydrolysed ATP, other nucleotide substrates were not degraded. The enzyme was activated by Mg^{2+} and inhibited by Ca^{2+} , other ions had no effect on activity.

Antibodies prepared to this enzyme bound to the polar regions of whole cells of C. jejuni as shown by protein A-colloidal gold immunoelectron microscopy. The antibodies to this ATPase cross reacted (shown by Western blotting) with four proteins from a whole cell extract of this organism, two proteins in A. serpens MW5, and three proteins for E. coli K12. They did not cross react with any proteins from Spirillum volutans, Methanococcus voltae, Vibrio cholerae, or rat liver mitochondria. Antibodies raised against the F₁-ATPase of E. coli K12 cross reacted with six proteins in a whole cell extract of this organism, and one protein species in each of the whole cell extracts of V. cholerae, A. serpens MW5, S. volutans, and rat liver mitochondria. These antibodies did not recognize any whole cell proteins from either C. jejuni, or M. voltae.

These results along with the ATPase activity localized by histochemical staining indicate that polar membrane functions as an ATPase.

Acknowledgements

The author wishes to thank Dr. R.G.E. Murray for his scientific input, support, patience, and long-suffering throughout the course of this work.

The technical assistance of Mickey Hall and Dianne Moyles was also greatly appreciated.

As well, the author is indebted to Zobeeda (Zo) Hosein, and Marion Kist for their scientific and social support during the years at Western.

Finally, and perhaps most important, the author wishes to thank her Mother and Father for their undying support (both moral and financial) and for their prayers.

The author was financially supported by Ontario Graduate Scholarships and the Medical Research Council of Canada.

This work was financed by the Medical Research Council of Canada.

This thesis is dedicated to
Dr. C.F. Robinow whose teaching
and enthusiasm for microbiology
led me down this road.

"For now we see in a mirror dimly, but then face to face,
now I know in part, but then I shall know fully just as I
also have been fully known." I Corinthians 13:12.

TABLE OF CONTENTS

	Page
CERTIFICATE OF EXAMINATION.....	ii
ABSTRACT.....	iii
ACKNOWLEDGEMENTS.....	vii
TABLE OF CONTENTS.....	x
LIST OF TABLES.....	xiv
LIST OF FIGURES.....	xv
CHAPTER 1 - INTRODUCTION.....	1
1.1. <u>Campylobacter jejuni</u> : Description of the organism.....	3
1.2. Ultrastructure of the bacterial cell envelope.....	4
1.3. Components of the bacterial cell envelope..	6
.1 Outer membrane.....	6
.2 Peptidoglycan.....	9
.3 Periplasmic space.....	11
.4 Cytoplasmic membrane.....	11
.5 Flagella.....	15
1.4. Polar membrane.....	17
1.5. Eucaryotic and procaryotic adenosine triphosphatases (ATPases).....	19
.1 Ion-transporting ATPases.....	20
.1 Monovalent and divalent cation- transporting ATPases.....	21
.2 Anion-transporting ATPases.....	24
.2 ATPases associated with contractile systems.....	24
.3 Proton-Translocating ATPases.....	24
.4 Bacterial H ⁺ -ATPases.....	28
.5 Solubilization and purification of the F ₁ portion.....	29
.6 Solubilization and purification of the FoF ₁ complex.....	31
.7 Molecular Weights of F ₁ -ATPases.....	32
.8 Electron Microscopic Studies on F ₁ - ATPases.....	32
.9 Subunit composition and arrangement of F ₁ -ATPases.....	33

1.5.10	Activation of the F1-ATPase.....	35
.11	Substrates of F1-ATPases.....	36
.12	Inhibition of F1-ATPases.....	36
1.6	Goals.....	38
CHAPTER 2 - MATERIALS AND METHODS.....		39
2.1	Organisms.....	39
.1	Cultivation and harvesting.....	40
.2	Preparation of cell envelopes.....	43
2.2	Electron microscopy.....	43
.1	Negative staining.....	43
.2	Fixation, embedding and sectioning.....	44
2.3	Visualization of polar membrane on membrane vesicles by negative staining....	45
.1	Stabilization of polar membrane on membrane vesicles.....	45
.2	Preparation of lipid-depleted membrane sheets.....	46
.3	Application of a modified cell monolayer technique of Nermut (1983) for visualiz- ing the inner surface of the plasma membrane.....	46
2.4	Separation of minicells from whole cells by sucrose density centrifugation...	47
2.5	Extraction and purification of ATPase from <u>C. jejuni</u>	48
.1	Shock-wash procedure.....	48
.2	Extraction with n-butanol.....	51
2.6	Determination of protein.....	53
2.7	Electrophoresis and molecular weight determinations.....	53
2.8	Measurement of ATPase activity.....	55
2.9	Substrate specificity.....	56
2.10	Effect of ions on enzyme activation.....	56
2.11	Cold lability.....	57
2.12	Effect of inhibitors on soluble and membrane bound ATPase activity.....	57
2.13	Effect of trypsin on the soluble ATPase activity.....	58
2.14	Reconstitution of ATPase activity in ATPase depleted membranes.....	58
2.15	Histochemical staining for ATPase activity.....	59
2.16	Antigen localization of ATPase in thin section.....	60
.1	Preparation of ANTisera and isolation of gamma globulin (IgG) fractions.....	60
.2	Antigen localization in thin sections of Spurr resin-embedded cells.....	61

2.16.3	Low-temperature embedding for antigen localization.....	63
.4	Antigenic relatedness of ATPase from <u>C. jejuni</u> to ATPases of other bacterial species and rat liver mitochondria.....	65
.5	Immunoblotting.....	65
CHAPTER 3	- RESULTS.....	67
3.1	Growth characteristics of <u>C. jejuni</u>	67
3.2	Ultrastructure of polar membrane.....	67
.1	Location in whole cells.....	67
.2	Association of polar membrane with the flagellar basal complex.....	69
.3	Fine structure and dimensions of polar membrane.....	70
.1	Observations on thin sections of <u>C. jejuni</u>	70
.2	Observations by negative stain on membrane preparations of <u>C. jejuni</u>	71
3.3	Minicell formation by <u>C. jejuni</u>	73
3.4	Localization of ATPase activity in whole cells by histochemical staining.....	75
3.5	Isolation and characterization of an F1-like ATPase from <u>C. jejuni</u>	77
.1	Analysis of ATP hydrolysing proteins in shock-wash membrane supernatants.....	77
.2	Purification of the F1-like ATPase.....	78
.3	Electron microscopy of ATPase.....	80
.4	Reconstitution of ATPase activity on depleted membranes.....	80
.5	Subunit composition.....	80
.6	Inhibition of membrane-bound and soluble F1-ATPase activity.....	82
.7	Enzyme activation and specificity.....	82
3.6	Immunocytochemical localization of ATPase from <u>C. jejuni</u>	85
3.7	Antigenic relatedness of the ATPase from <u>C. jejuni</u> to ATPases of other bacterial species and rat liver mitochondria.....	87
CHAPTER 4	- DISCUSSION.....	90
4.1	Ultrastructure of polar membrane.....	91
4.2	Minicell formation by <u>C. jejuni</u>	95
4.3	Localization of ATPase activity in whole cells by histochemical staining.....	98
4.4	Isolation and characterization of an F1-like ATPase from <u>C. jejuni</u>	101
4.5	Immunocytochemical localization of ATPase in <u>C. jejuni</u>	109

4.6	Antigenic relatedness of the ATPase from <u>C. jejuni</u> to ATPases of other bacterial species and rat liver mitochondria.....	111
4.7	Conclusions.....	112
FIGURES.....		116
REFERENCES.....		210
VITA.....		262

LIST OF TABLES

<u>Table</u>	<u>Description</u>	<u>Page</u>
1	Purification of soluble ATPase	79
2	Reconstitution of ATPase activity in depleted membranes	81
3	Effects of inhibitors on the membrane bound and soluble purified ATPase from <u>C. jejuni</u>	83
4	Effect of trypsin on soluble ATPase activity	84
5	Effect of freezing (-20°C) on soluble ATPase activity	96

LIST OF FIGURES

<u>Figure</u>	<u>Description</u>	<u>Page</u>
1	Diagram of the polar membrane in <u>Ectothiorhodospira mobilis</u> .	117
2	Conceptual model of FOF1 of <u>Escherichia coli</u> .	119
3	Growth Curve for <u>C. jejuni</u> .	121
4	Thin section of <u>C. jejuni</u> fixed with 5% acrolein and 0.25% glutaraldehyde.	123
5	Thin section of <u>A. serpens</u> MW5 fixed with 5% acrolein and 0.25% glutaraldehyde.	125
6	Cross section of <u>C. jejuni</u> .	127
7	Cross section of <u>A. serpens</u> MW5	129
8	Thin section of <u>C. jejuni</u> showing septum and polar membrane.	131
9	Thin section of <u>C. jejuni</u> showing septum with no associated polar membrane.	133
10	Thin section of polar region of <u>C. jejuni</u> showing association of polar membrane with the flagellar basal complex.	135
11	Thin section of the polar region of <u>C. jejuni</u> showing association of polar membrane and the flagellar basal complex.	137
12	Thin section of the polar region of <u>C. jejuni</u> showing association of polar membrane to the flagellar basal complex.	139
13	Thin section of the polar region of <u>C. jejuni</u> showing the association of polar membrane to the flagellar basal complex.	141

14	Thin section of the polar region of <u>C. jejuni</u> showing the association of polar membrane to the flagellar basal complex.	143
15	Thin section of the polar region of <u>C. jejuni</u> showing the association of polar membrane to the flagellar basal complex.	145
16	Longitudinal and cross section of <u>C. jejuni</u> showing polar membrane and rigid cytoplasmic membrane.	147
17	French pressure cell disrupted membrane preparations of <u>C. jejuni</u> negatively stained with 1% phosphotungstic acid.	149
18	French pressure cell disrupted membrane preparation of <u>C. jejuni</u> negatively stained with 1% phosphotungstic acid.	151
19	Negatively stained membrane preparations of <u>C. jejuni</u> .	153
20	Thin section of membrane preparations from <u>C. jejuni</u> pre-fixed with 1% tannic acid.	155
21	Negatively stained minicells from <u>C. jejuni</u> bearing flagella.	157
22	Thin sections of minicells from <u>C. jejuni</u> .	159
23	Thin sections of <u>C. jejuni</u> showing formation of minicells.	161
24	Negatively stained whole cell of <u>C. jejuni</u> showing minicell forming in the polar region.	163
25	Thin section of minicell preparation showing contamination with whole cells, membrane vesicles and flagellar debris.	165
26	Negative stain of a minicell showing hexagonal pattern in the interior.	167
27	Thin sections of <u>C. jejuni</u> histochemically stained for the presence of Mg ²⁺ -ATPase activity.	169

28	Thin sections of spheroplasts of <u>C. jejuni</u> histochemically stained for Mg ²⁺ -ATPase activity.	171
29	Thin sections of <u>C. jejuni</u> histochemically stained for the presence of Ca ²⁺ -ATPase activity.	173
30	Thin sections of <u>E. coli</u> K12 histochemically stained for the presence of Ca ²⁺ - and Mg ²⁺ -ATPase activity.	175
31	Thin section of <u>B. cereus</u> histochemically stained for the presence of Mg ²⁺ -ATPase activity.	177
32	Thin sections of <u>M. luteus</u> histochemically stained for the presence of Mg ²⁺ - and Ca ²⁺ -ATPase activity.	179
33	Thin sections of <u>D. radiodurans</u> histochemically stained for the presence of Mg ²⁺ - and Ca ²⁺ -ATPase activity.	181
34	Thin sections of <u>A. serpens</u> MW5 histochemically stained for the presence of Mg ²⁺ - and Ca ²⁺ -ATPase activity.	183
35	Non-dissociating gel analysis of protein profiles and ATPase activity in polyethylene glycol precipitated material from the initial shock-wash supernatant from membranes of <u>C. jejuni</u> .	185
36	Non-dissociating-PAGE of purified ATPase.	187
37	Negatively stained solubilized ATPase from <u>C. jejuni</u> .	189
38	SDS-PAGE of RNA polymerase and <u>C. jejuni</u> ATPase. Urea-PAGE of <u>C. jejuni</u> ATPase.	191
39	Non-dissociating-PAGE of the effect of cations on <u>C. jejuni</u> ATPase activity.	193
40	Non-dissociating-PAGE of the substrate specificity of the ATPase of <u>C. jejuni</u> .	195
41	Immunocytochemical localization of <u>C. jejuni</u> ATPase on thin sections of Spurr embedded whole cells.	197

42	Immunocytochemical localization of <u>C. jejuni</u> ATPase on thin sections of Spurr embedded whole cells.	199
43	Immunocytochemical localization of <u>C. jejuni</u> ATPase on hydrogen peroxide etched sections of Spurr embedded whole cells.	201
44	Immunocytochemical localization of <u>C. jejuni</u> ATPase on thin sections of Lowicryl KM4 embedded whole cells.	203
45	Immunocytochemical localization of <u>C. jejuni</u> ATPase on thin sections of Lowicryl KM4 embedded whole cells.	205
46	Western blots of whole cell proteins reacted with antibody prepared against the ATPase isolated from <u>C. jejuni</u> .	207
47	Western blots of whole cell proteins reacted with antibody prepared against the F1-ATPase of <u>E. coli</u> K12.	209

The author of this thesis has granted The University of Western Ontario a non-exclusive license to reproduce and distribute copies of this thesis to users of Western Libraries. Copyright remains with the author.

Electronic theses and dissertations available in The University of Western Ontario's institutional repository (Scholarship@Western) are solely for the purpose of private study and research. They may not be copied or reproduced, except as permitted by copyright laws, without written authority of the copyright owner. Any commercial use or publication is strictly prohibited.

The original copyright license attesting to these terms and signed by the author of this thesis may be found in the original print version of the thesis, held by Western Libraries.

The thesis approval page signed by the examining committee may also be found in the original print version of the thesis held in Western Libraries.

Please contact Western Libraries for further information:

E-mail: libadmin@uwo.ca

Telephone: (519) 661-2111 Ext. 84796

Web site: <http://www.lib.uwo.ca/>

Chapter 1

Introduction

In 1963, Murray and Birch-Andersen observed, in thin sections of Aquaspirillum serpens, "an extremely thin membranous component 200 Å underneath the plasma membrane". This structure appeared to be connected to the overlying plasma membrane by short bar-like links of regular length, and was found in association with the poles of the cell. The membranous appearance and location of this structure led the authors to name this specialized region "polar membrane". In 1966, Ritchie, et. al., described a similar structure in Vibrio fetus (now, Campylobacter fetus) and in 1968, Remsen, et. al., further defined the structure in Ectothiorhodospira mobilis as being composed of "a series of projections" arising out of the plasma membrane "which had stalks 50 Å long, which terminated with a knob-like structure approximately 75-80 Å in diameter". In the twenty-three years which have passed since the initial reports of polar membrane the structure has been observed in a variety of polarly flagellated Gram-negative bacterial genera, including Rhodospirillum, Selenomonas, Chromatium,

Vibrio and Methanococcus (Hickman and Frenkel, 1965; Chalcroft, et. al., 1973; Smibert, 1978; Ferris and Beveridge, 1985; Koval and Jarrel, 1987). However, to date, the biochemical composition and function of polar membrane is unknown. Its location at the poles of the cell surrounding the flagellar insertion point has led to the hypothesis that polar membrane functions in energy transductions (Murray and Birch-Andersen, 1963). As well, the component units are structurally similar to the ATPase of the inner mitochondrial membrane and the cytoplasmic membrane of various bacterial species (Racker 1967; Harold, 1972) because they are all composed of stalks and knobs. A study of polar membrane in Campylobacter jejuni has been pursued to try and resolve both its structure and function.

The purpose of this introduction is to acquaint the reader with 1) the organism under investigation, 2) the structure and function of bacterial cell walls and envelopes, with emphasis on the cell envelopes of Gram-negative bacteria and flagella basal complex, which the polar membrane is most closely associated with, 3) the currently known facts of polar membrane, including further elaboration on its structure, and possible function(s), 4) eucaryotic and procaryotic adenosine triphosphatases and finally 5) an outline of the goals of this study.

1.1 Campylobacter jejuni : Description of the organism.

C. jejuni is a Gram-negative; nonsporeforming, microaerophilic, helically curved, slender bacillus (Smibert, 1978 and 1984). Various morphological forms have been observed in cultures of this organism, including s-shaped, gull-winged, comma, dimpled, and coccoid (Karmali and Skirrow, 1984; Merrell, et al., 1981; Smibert, 1978 and 1984; Ng, et al., 1985). This organism readily produces coccoid forms or bodies late in its growth cycle or upon exposure to atmospheric levels of oxygen (Ogg, 1962; Smibert, 1965 and 1984). This phenomenon has led to questioning of the purity of cultures being examined.

C. jejuni is motile by means of a single, unsheathed, polar flagellum at one or both ends of the cell. The flagellum may be two to three times the length of the cell, and propels the organism in a characteristic corkscrew-like motion (Smibert, 1984).

C. jejuni causes abortion in sheep and enteritis with fever in man. It has been found, over the last few years, to cause as much enteritic disease in man as Salmonella and Shigella and is therefore considered an important human intestinal pathogen (Smibert, 1984). This organism can be found in the intestinal tract of poultry, dogs, cats, sheep, rabbits, monkeys, sea gulls, cattle, and pigs (Smibert,

1965, 1969, 1978, and 1984).

1.2. Ultrastructure of the Bacterial Cell Envelope

The taxonomic classification of bacteria primarily begins with the use of the Gram reaction (Gram, 1884). This staining procedure has resulted in the division of bacteria into two major groups, Gram-positive and Gram-negative. This staining procedure showed initial structural and molecular differences between the outer layers of bacteria that retain the purple iodine-gentian violet complex after extraction with polar solvents such as alcohol or acetone (Gram-positive), as compared to those that do not do so (Gram-negative). The lack of retention of this dye complex in Gram-negative bacteria is now thought to be a result of the dissolution of the outer membrane by the polar solvent allowing the cell to become leaky (Beveridge and Davies, 1983). A few anomalies to this system of classification do exist. The archaebacteria possess cell walls of a simple organization. The cell envelopes of the archaebacteria Sulfolobus acidocaldarius and Halobacterium cutirubrum consist of a cytoplasmic membrane and external to this a regularly-structured array of glycoprotein subunits (Kandler, 1982; Stoeckenius and Rowen, 1967; Weiss, 1974). Butyrivibrio fibrisolvens possesses a Gram-positive wall

type but stains Gram-negative (Cheng and Costerton, 1977) and Deinococcus radiodurans has a Gram-negative wall type but stains Gram-positive (Thompson and Murray, 1982)

Gram-positive eubacteria possess a relatively simple cell wall profile. Initial studies on the cell wall of Bacillus cereus revealed the presence of a thick, densely staining, mucopeptide layer. At first, a distinct plasma membrane was not seen (Chapman and Hillier, 1953), but the development of better fixation and embedding techniques for the electron microscope resulted in the visualization of a 7.5-8.0 nm double-track cytoplasmic membrane in a variety of Gram-positive eubacteria (Murray, 1957; Kellenberger and Ryter, 1958; Van Iterson, 1961)

The cell walls of Gram-negative bacteria are more complex than those of Gram-positive organisms. In the early studies of the surfaces of Gram-negative bacteria there were difficulties in observing more than a single dense surface layer. The first clear demonstration of the plasma membrane and the cell wall as two separate structures was accomplished by Kellenberger and Ryter (1958). By the use of special fixation and embedding techniques they showed that, in Escherichia coli, the plasma membrane was a single dense layer, 6.0-8.0 nm thick, and the cell wall was a triple-layered structure, consisting of two outer dense layers separated by a less dense layer, with a total diameter of 6.0-9.0 nm. One of the components of this

triple-layered wall was a second double-track membrane which was termed the outer membrane (Birch-Andersen *et al.*, 1953). As techniques improved further it was found that an additional layer was present. A dense layer was seen between the plasma membrane and the cell wall, first in Vitreoscilla (Costerton, *et al.*, 1961), Spirillum serpens (Murray, 1963), and Moraxella (Ryter and Piechaud, 1963), and later in a wide range of Gram-negative bacteria (Glauert and Thornley, 1969). This was a 2.0-3.0 nm thick layer of murein, or peptidoglycan, aligned between the two membranes (De Petris, 1965, 1967; Murray *et al.*, 1965).

Today it is known that many bacteria possess cell envelopes of even greater complexity. Additional layers have been found external to the peptidoglycan layer of Gram-positive and the outer membrane of Gram-negative eubacteria. These layers include capsules, slime layers, and regularly-structured arrays of protein (Beveridge, 1981).

1.3. Components of the Bacterial Cell Envelope

1.3.1. Outer Membrane

The outer membrane of Gram-negative bacteria appears in thin section as a wavy, double tracked membrane varying from 6 to 10 nm in diameter according to the species.

Projections and blebs from this membrane are frequently seen (De Petris, 1967). The outer membrane can be separated from the cytoplasmic membrane because of its greater density due to the presence of lipopolysaccharide (LPS) (Osborn *et al.*, 1972). The isolated outer membrane has been found to be composed of approximately 20-25% phospholipid, 30% lipopolysaccharide, and 45-50% protein (Di Rienzo *et al.*, 1978).

The phospholipid is found mainly in the inner leaflet of the membrane bilayer, and the lipopolysaccharide is associated with the outer leaflet (Funahara and Nikaido, 1980). The phospholipids of the outer membrane are quantitatively similar to, but qualitatively different from those of the cytoplasmic membrane in *E. coli* (White, *et al.*, 1972; Osborn, *et al.*, 1972). Outer membrane is enriched in phosphatidylethanolamine as compared to the cytoplasmic membrane (Rottem *et al.*, 1975).

Lipopolysaccharides are common components of the cell envelopes of all Gram-negative eubacteria and are apparently unique to these organisms (Osborn, 1969). LPS is an amphipathic molecule consisting of complex phosphorylated heteropolysaccharides covalently linked to a specific glucosamine-containing lipid, Lipid A (Luderitz *et al.*, 1974; Osborn, 1969). Lipid A is embedded in the outer leaflet of the outer membrane and the heteropolysaccharides

extend outwards from the cell surface (Beveridge, 1981; Costerton et al., 1974).

The protein composition of the outer membrane differs from that of the cytoplasmic membrane in that it contains a small variety of proteins in large quantities (Inouye, 1979). It has, therefore, been relatively easy to purify and characterize these major proteins. The outer membrane of wild-type *E. coli* K-12 contains at least three classes of major proteins: (i) matrix proteins (Rosenbusch, 1974; Di Rienzo et al., 1978; Osborn and Wu, 1980) which when viewed by negative stain on the electron microscope appear as a tightly packed lattice with a 7.7 nm repeat (Rosenbusch, 1974; Steven et al., 1977). These proteins are thought to form pores allowing passive diffusion (Nakae, 1976a, 1976b); (ii) The omp A protein (also known as the tolG protein) is found in large quantities in the surface of the outer leaflet of the outer membrane (Inouye and Yee, 1972). Omp A is thought to function in F pilus-mediated conjugation (Manning et al., 1976; Manning and Reeves, 1976), and as bacteriophage receptors (Datta et al., 1977). (iii) lipoprotein (also known as the Braun protein) which may function in maintaining the integrity of the outer membrane structure (Braun and Rehn, 1969; Braun and Sieglin, 1970; Braun and Wolff, 1970; Di Rienzo et al., 1978). Ten to

9

twenty minor protein species are also present in the outer membranes of Gram-negative eubacteria (Inouye, 1979). These proteins function in the transport of nutrients, vitamins and metals, and may also act as bacteriophage and bacteriocin receptors (Di Rienzo *et al.*, 1978).

The outer membrane is remarkably devoid of enzymatic activity. A protease and a phospholipase have been isolated from the outer membrane of *E. coli* (MacGregor *et al.*, 1979; Bell *et al.*, 1971).

1.3 2 Peptidoglycan

Peptidoglycan is a fundamental polymer that is a common component of the cell walls of Gram-positive and Gram-negative eubacteria, Rickettsiae and blue-green bacteria. This structure is composed of N-substituted glucosamine and muramic acid (3-O lactylglucosamine) linked together by 1-4 beta bonds (Rogers, 1983). It is the peptidoglycan that is responsible for the rigidity and integrity of the shape of the cell. When peptidoglycan is degraded, for example by lysozyme, the bacterium tends to lose its characteristic shape and to form a spherical body known as a protoplast in Gram-positive organisms and a spheroplast in Gram-negatives (Rogers *et al.*, 1980). The

chemical composition of peptidoglycan has been worked out over a period of thirty-five years following the classical studies of Weidel (1950) and of Salton (1953).

The polysaccharide or glycan chains of peptidoglycan are crosslinked by short peptides, in any one peptidoglycan, of a limited number of amino acids

Four basic types of peptidoglycan have so far been determined (Ghuysen, 1968) and are differentiated by the way in which the peptides are linked together. However, the range of amino acids found in peptidoglycans from different species of bacteria is large. A complicated classification of the different types of peptidoglycan taking into account this wide variation has been devised by Schleifer and Kandler (1972).

The peptidoglycan of Campylobacter species has been found to contain alanine, lysine, glycine, glutamic acid, aspartic acid, and either cysteine or diaminopimelic acid (Keeler et al., 1966). The peptidoglycan of intestinal and venereal strains of C. fetus is composed of muramic acid, glucosamine, alanine, glutamic acid, and diaminopimelic acid (1:1:2:1:1) and the repeating unit in the polymer was presented as Glc NAc-MurNAc-L-Ala-D-Glu-meso-Dpm-D-Ala (Smibert, 1978).

1.3.3 Periplasmic Space

The periplasmic space is the region between the plasma and outer membranes of Gram-negative eubacteria. This space constitutes some 20-40 per cent of the total cell volume and is distinguished by a distinct ionic composition and the presence of a unique series of proteins (Beveridge, 1981). This space may be compartmentalized by the presence of adhesion zones or sites connecting the plasma and outer membrane (Bayer, 1968). The proteins found within this area perform a variety of functions. Some have a catalytic function, converting various substrates into forms for translocation by specific carriers present in the cytoplasmic membrane (Kellermann and Szmecman, 1980; Lo, 1979); others can be classed as hydrolytic enzymes, some of which are responsible for the destruction of antibacterial agents penetrating the outer membrane (Wetzel et al., 1910; Curtis et al., 1972).

1.3.4. Cytoplasmic Membrane

The cytoplasmic membrane appears, on thin section, as a 7.0-8.0 nm thick double track layer surrounding the cytoplasm (De Petris, 1965, 1967). There is little difference between the plasma membranes of Gram-positive

and Gram-negative bacteria except for the presence of lipoteichoic acid in the membranes of the Gram-positive organisms (Ward, 1981). Bacterial cytoplasmic membranes are composed of approximately 50-70% protein and 20-30% lipid (as reviewed by Rogers *et al.*, 1980).

Most of the lipid portion of the membrane consists of phospholipids such as phosphatidylglycerol, diphosphatidylglycerol (cardiolipin) and phosphatidylethanolamine (Lechevalier, 1977). The phospholipid composition of the cytoplasmic membrane of Campylobacter species includes, in addition to those mentioned above, phosphatidyl inositol, phosphatidyl serine and the glycolipid digalactosyldiglyceride (Smibert, 1978). The fatty acids of bacterial lipids are generally of the saturated, monounsaturated, or cyclopropane variety (Kaneda, 1977). The major fatty acids found in the plasma membrane of C. jejuni were C-10:, C-12:0, C-14:0 present in small amounts and in larger amounts C-16:0, C-16:1, and C-18:1 (Smibert, 1978).

When cytoplasmic membrane is disaggregated by heating at 100 °C for one minute in 1% SDS and electrophoresis is applied in a polyacrylamide gel, approximately 30-40 protein bands can be separated (as reviewed by Rogers *et al.*, 1980). Rottem and Razin (1967) have suggested that gel patterns from unidirectional electrophoresis of membrane proteins are

sufficiently characteristic to be used as "finger prints" in the classification of Mycoplasmas. Using the O'Farrell (1975) two dimensional electrophoresis method, a very much larger number of spots can be seen in plasma membrane preparations. Approximately 140 spots were visualized in membrane preparations from Acholeplasma laidlawii (Archer et al., 1978). The number and variety of proteins present in this membrane suggest it to be multifunctional in nature (Salton, 1974). To date, plasma membrane has been shown to possess the enzymes responsible for oxidative phosphorylation, electron transport, wall biosynthesis and active transport of elements, nutrients and metabolites (Salton, 1971, 1974). Most of these enzymes have been identified by using crossed immunoelectrophoresis and include dehydrogenases utilizing succinate, 1,6-phosphogluconate, dihydro-orotate, two utilizing NADH, malate, glycerol-3-phosphate, lactate and glutamate as substrates; an adenosine triphosphatase and a protease (Johansson and Hjerten, 1974; Johansson and Wroblewski, 1978; Owen and Salton, 1975, 1977; Owen and Smyth, 1977; Salton, 1978; Salton and Owen, 1976; Smyth et al., 1976, 1978).

Abram (1965) showed that, when negatively stained with phosphotungstate (pH 7.0-7.2), the plasma membranes of Bacillus stearothermophilus were studded with particles

approximately 6.5-8.5 nm in diameter that were deduced to be on the inner surface of the membrane. These particles were attached to the membrane by fine stalks of the order of 5 nm long. Similar particles, found on the plasma membranes of Streptococcus faecalis and Micrococcus lysodeikticus, have been stripped from the membranes (by washing in a low ionic strength buffer plus EDTA, ethylenediamine tetraacetate), purified and shown to possess ATPase activity (Oppenheim and Salton, 1973; Schnebli et al., 1970; Munoz et al., 1968). The location of the ATPase activity on the cytoplasmic side of the plasma membrane has been demonstrated in M. lysodeikticus by two methods; (1) antiserum prepared against the purified ATPase was reacted with protoplasts and isolated membrane preparations inhibiting the ATPase activity in the membrane preparations but not in the intact protoplasts (Oppenheim and Salton, 1973; Salton, 1978), and (2) ferritin labelled antibodies to the purified ATPase were found to bind to membranes in a pattern similar to that of the particles seen by negative staining on unreacted membranes (Oppenheim and Salton, 1973).

The inner leaflet of the cytoplasmic membrane of Campylobacter species is thickened at the polar region (Ritchie et al., 1966). It possesses on its inner surface a structure termed polar membrane. This structure, as

mentioned earlier, is also found in a variety of polarally flagellated bacteria and will be discussed at length later in this review.

1.3.5. Flagella

C. jejuni is actively motile by means of a single polar flagellum at one or both ends of the cell (Smibert, 1978). The flagella are composed of three parts: (1) the basal body, (2) the hook, and (3) the filament.

The flagellar basal complex in Gram-negative bacteria consists of four rings (or discs) surrounding a central axial rod (Doetsch and Sjobald, 1980). These rings were originally designated by DePamphilis and Adler (1971) according to their position and association with various structures in the cell envelope of E. coli: the L ring for its association with the outer (lipopolysaccharide) membrane, the P ring with the peptidoglycan, the S ring (for supramembrane) just above the cytoplasmic membrane, and M ring associated with the plasma membrane. Some basal bodies are more complex. This structure in Caulobacter crescentus has five rings (Johnson et al., 1979) and in A. serpens, V. cholerae, and C. fetus a number of concentric rings (CMR's) are found surrounding the central axial rod and appear to be located between the L and P rings (Coulton and Murray, 1977,

1978; Ferris and Beveridge, 1984). In C. fetus the S and M rings were found to be associated with the plasma membrane (possibly with the outer and inner hydrophilic surfaces, respectively), the L ring was embedded in the outer membrane and the P ring was associated with the peptidoglycan (Ferris and Beveridge, 1984). In both A. serpens and C. fetus fibrous material was found associated with the proximal end of the basal complex emanating from the M ring (Coulton and Murray, 1977, 1978; Ferris and Beveridge, 1984). The basal body is thought to function in two ways: (1) as an anchor of the flagella to the cell, and (2) as a "motor", with the M ring generating torque for rotation by interacting with a "stator" (possibly the S ring or the outer surface of the axial rod). The L and P rings are thought to function as bushings allowing free rotation of the axial rod in the cell wall (Berg, 1974, 1975; Lauger, 1977; Macnab, 1979).

The hook is located at the base of the filament. It is approximately 900 nm long and consists of a single protein of 42,000 daltons in E. coli and 33,000 daltons in Bacillus subtilis (Kagawa et al., 1976; Silverman and Simon, 1972; Dimmitt and Simon, 1971). This structure is thought to act as a "universal joint" between the basal body and the filament (Silverman and Simon, 1977).

The filament is a tubule varying in size from 12 to 15 nm in diameter, composed of identical protein subunits (flagellin) in one of two configurations (Silvermon and Simon, 1977). The arrangement of the flagellin subunits in intact flagella gives rise to two different types of filament configuration: an A form, which is characterized by helically connected globules, and a thick-lined B form in which helical organization is not observed (Lowy and Hanson, 1965). The flagella of C. fetus are approximately 21 nm in diameter during the mid to late stationary phase of growth and about 18 nm in diameter after 24 hours of growth (McCoy et al., 1976).

1.4. Polar Membrane

A structure comprised of an ordered array of densely staining short bar-like particles arising out of the cytoplasmic membrane was first described in A. serpens by Murray and Birch-Andersen (1963). They termed this structure "polar membrane". Cohen-Bazire and Kunisawa (1963) observed a similar structure in Rhodospirillum rubrum and termed it a "polar cap". Ritchie et al. (1966) noted the presence of this structure in V. fetus (now C. fetus) and chose to describe it as a complex cytoplasmic membrane. To date the term "polar membrane" seems to have been adopted, although

it is slightly misleading in that the structure is not always found at the poles of the cell and when well resolved in thin sections it does not appear to have the structure of membrane. This structure has also been found in a variety of other bacterial species including Ectothiorhodospira mobilis (Remsen et al., 1968), Selenomonas ruminantium (Chalcroft et al., 1973), Vibro cholera (Ferris and Beveridge, 1984), and Methanococcus voltae (Koval and Jarrel, 1987).

The best description to date of polar membrane comes from the work of Remsen et al. (1968) on Ectothiorhodospira mobilis. This structure was about 30 nm thick and consisted of plasma membrane (7.5 nm), a dense area about 10 nm thick, and series of projections which terminated with knob-like structures approximately 7.5 to 8.0 nm in diameter. Figure 1 illustrates the pertinent morphology of polar membrane and proximity to the flagellar basal complex (Figure 4). In C. fetus the polar membrane is usually found at the flagellated pole, however, its presence has also been observed near the point of septum formation and presumably the site of insertion of a future flagellum (Ritchie et al., 1966). Polar membrane covers the inner surface of the cytoplasmic membrane and is near to the flagellar basal complex, but has never been seen to cover the membrane at the base of this structure or to join up with any component of it (Murray and

Birch-Andersen, 1963; Ritchie et al., 1966; Coulton and Murray 1978) It has also been observed that the cytoplasm in the polar regions associated with polar membrane is remarkably devoid of ribosomes (Murray and Birch-Andersen, 1963; Ritchie et al., 1966).

The location of this structure has led to the hypothesis that polar membrane may be involved in energy transformations, such as the liberation of energy for flagellar rotation or the synthesis of new cell wall (Murray and Birch-Andersen, 1963). Ritchie et al (1966) proposed that the complex cytoplasmic membrane (polar membrane) may contribute a torsional force at some stage of development and influence the conformation of the organism, as at the time this structure had been identified in only spirally curved bacterial species. This hypothesis was deemed unlikely with the discovery of polar membrane in the straight rod Chromatium (Smibert, 1978).

1.5. Eucaryotic and Procaryotic Adenosine Triphosphatases (ATPases)

All forms of living cells conserve energy in chemical compounds, especially in ATP. Enzymes are required for the synthesis of and hydrolysis of these energy rich compounds.

All ATP-hydrolysing enzymes are not usefully regarded as ATPases. There is a large group of non-specific phosphatases for which ATP is only one of a wide range of possible substrates (e.g. Alkaline phosphatases (Leonard and Provenza, 1972)). Therefore for the purposes of this review, an enzyme will only be regarded as an ATPase if its activity is largely confined to nucleoside triphosphates and ATP is one of its optimal substrates.

To date three broad groups of ATPases have been distinguished: (1) ion-transporting ATPases, (2) ATPases associated with contractile systems, and (3) proton-translocating ATPases.

1.5.1. Ion-transporting ATPases

A large group of enzymes of this class have been identified in the plasma membranes and other membrane sites of eucaryotic cells. These enzymes normally are involved in ATP hydrolysis, using the energy so derived for active ion transport. However, when ion gradients become steep enough, these enzymes may function in a synthetic mode (e.g. the Ca^{2+} -transporting ATPase of the sarcoplasmic reticulum (Hasselbach and Suko, 1974)).

This class of enzymes includes (i) monovalent and divalent cation transporting ATPases, and (ii) anion transporting ATPases.

1.5.1.1. Monovalent and divalent cation-transporting ATPases

This group of enzymes includes the Na^+, K^+ -dependent ATPase of mammalian plasma membranes, the K^+, H^+ -ATPase of fungal, plant and gastric parietal cell plasma membranes, Ca^{2+} -ATPase of the sarcoplasmic reticulum, vacuolar proton ATPases and a K^+ -ATPase found in *E. coli*.

The existence of a Na^+, K^+ -dependent ATPase in the plasma membranes of mammalian cells was first recognized when it was found that the sodium pump of nerve axons was coupled to ATP hydrolysis (Hodgkin and Keynes, 1955). A similar enzyme system in erythrocyte plasma membranes was found to be dependent on Mg^{2+} , Na^+ , and K^+ (Glynn, 1957). These enzymes function in the maintenance of Na^+ and K^+ gradients across plasma membranes, bulk transport of monovalent cations (Epstein, 1969), and less directly, in the transport of other substances coupled to electrochemical gradients set up by the Na^+, K^+ -ATPases (Bronk and Lesse, 1974).

These Na^+, K^+ -dependent ATPases, when purified, are found to be composed of a 106,000 dalton alpha-subunit (Hastings and Reynolds, 1979; Esmann *et al.*, 1980) and a 37,000 to 40,000 dalton beta subunit (Hastings and Reynolds, 1979; Esmann *et al.*, 1980; Sabatine *et al.*, 1981).

As well, a 10,000 to 12,000 dalton proteolipid has been identified in purified preparations of these enzymes and has been termed the gamma subunit (Hokin *et al.*, 1973; Forbush *et al.*, 1978).

Na^+/K^+ -dependent ATPases have been crystallized in the membrane, forming two dimensional crystals, by using vanadate (Skriver *et al.*, 1981; Maunsbach *et al.*, 1983) and inorganic phosphate (P_i) (Herbert *et al.*, 1982) as stabilizing ligands. Negative staining of these membrane crystals showed distinct surface particles of 30 to 50 nm in diameter (Deguchi *et al.*, 1977; Skriver *et al.*, 1981).

Na^+/K^+ -dependent ATPases are inhibited by ouabain, vanadate and diethylstilbesterol (De Meis and Vianna, 1979; Sarkadi, 1980).

The plasma membranes of fungi, plants, and gastric parietal cells possess K^+/H^+ -ATPases. These enzymes are composed of a major subunit of approximately 100,000 daltons (Malpartida and Serrano, 1980, 1981; Dufour and Goffeau, 1978; Addison and Scarborough, 1981; Bowman *et al.*, 1981; Bara and Serrano, 1982; Uchida *et al.*, 1985), and are inhibited specifically by vanadate, diethylstilbesterol, and dicyclohexylcarbodiimide (DCCD), but not by ouabain, azide, oligomycin or molybdate (Goffeau and Slagman, 1981; Delhez *et al.*, 1977; Serrano, 1978; Gallagher and Leonard, 1982; Hodges, 1976; Bowman *et al.*, 1978; Bowman, 1983).

The Ca^{2+} -transporting ATPase of sarcoplasmic reticulum is responsible for the cessation of muscle contraction cycles by removing Ca^{2+} from the sarcoplasm into the sarcoplasmic reticulum cavities (as reviewed by Moller *et al.*, 1982). These enzymes range in molecular weight from 90,000 to 150,000 daltons and appear to be composed of a single subunit type (MacLennan *et al.*, 1971; MacFarland and Inesi, 1971; Meissner *et al.*, 1973). They are located on the outer surface of the sarcoplasmic reticulum and have been crystallized in the membrane, forming two dimensional crystals, by using vanadate as a ligand (Dux and Martonosi, 1983). When negatively stained, these enzymes appear as particles of approximately 4.0 nm in diameter connected to the membrane via a stalk 2.0 nm long (Jilka *et al.*, 1975; Dux and Martonosi, 1983). These Ca^{2+} -ATPases have the same inhibition pattern as the mammalian Na^+, K^+ -ATPases (Josephson and Cantley, 1977; O'Neal *et al.*, 1979).

The vacuolar proton ATPases are found in eucaryotic membranes such as endoplasmic reticulum, the Golgi complex and secretory vacoules. These enzymes are distinguished from the aforementioned ones by virtue of their inhibitor specificities. They are not inhibited by oligomycin, efrapeptin, azide, aurovertin (mitochondrial F_1 -ATPase inhibitors), vanadate or ouabain (Na^+, K^+ -ATPase

inhibitors), but are specifically inhibited by alkylating reagents such as N-ethylmaleimide (NEM) and 4-chloro-7-nitrobenzo-2-oxa-1,3-diazol (NBD-Cl) (Stone *et al.*, 1983; Forgae *et al.*, 1983; Galloway *et al.*, 1983; Glickman *et al.*, 1983; Ohkuma *et al.*, 1982; Bowman, 1983).

A K^+ transporting ATPase has also been isolated from the plasma membranes of the bacterium *E. coli* and found to be associated with the Kdp transport system (Epstein *et al.*, 1978). This enzyme is stimulated by K^+ and is not inhibited on the membrane by DCCD which inhibits the Ca^{2+} , Mg^{2+} -ATPase (F_1F_0 -ATPase) of this organism (Epstein *et al.*, 1978).

1.5.1.2. Anion-Transporting ATPases

These enzymes are stimulated by HCO_3^- and differ from Na^+ , K^+ -dependent ATPases in that they are not inhibited by ouabain (Spennay *et al.*, 1973; Kinne-Saffran and Kine, 1974; Simon *et al.*, 1971).

1.5.2. ATPases associated with contractile systems

This group of ATPase activities are associated with filamentous and other non-membranous structures in the cytoplasmic matrix of eucaryotic cells. The activity of these enzymes is assumed to be related to motility and

contraction events, such as the myosin ATPase of muscle cells (Tonomura and Inoue, 1974). These enzymes are not inhibited by ouabain or oligomycin (Tonomura and Inoue, 1974).

1.5.3. Proton-Translocating ATPases

Enzymes of this group form part of the inner mitochondrial membrane, the grana membranes of chloroplasts and the plasma membranes of bacteria. In mitochondria and chloroplasts of eucaryotes the ATPase is required for the formation of ATP from ADP. In procaryotes the ATPase is located in the plasma membranes of both aerobic and anaerobic species (Downie *et al.*, 1979).

These proton-translocating ATPases are a part of a complex of two enzyme systems found in energy-transducing membranes: they consist of the electron transport proteins and the proton ATPase complex (H^+ -ATPase) (Amzel and Pedersen, 1983), and they appear to be coupled by an electrochemical gradient of protons generated by the redox components (Mitchell, 1966, 1981). The H^+ -ATPases function by (1) allowing a flow of protons across the membrane (i.e. the translocation step), (2) catalysing the formation of ATP from ADP and inorganic phosphate (P_i), and (3) directing protons in such a way as to use the energy stored in the

electrochemical gradient to produce high ATP concentrations (i.e. the coupling step) (Amzel and Pedersen, 1983).

Oxidative phosphorylation is, as with many enzymatic reactions, reversible and ATP hydrolysis can be used to induce the formation of an electrochemical gradient of protons (Amzel and Pedersen, 1983). In the anaerobic bacterium Strep. faecalis the in vivo direction of the H^+ -ATPase is toward hydrolysis of ATP (Abrams, et al. 1976). In the mitochondria of animal cells and in most aerobic bacteria the physiological direction of these enzymes is primarily toward ATP synthesis (Amzel and Pedersen, 1983). For this reason, these enzymes have been termed ATP synthase (or synthetase) and those that function primarily in ATP hydrolysis have been called Mg^{2+} , Ca^{2+} -ATPases (Futai and Kanazawa, 1983). This latter term is somewhat confusing, because it tends to suggest the transport of divalent cations rather than the translocation of protons. These proton-translocating ATPases are commonly referred to as H^+ -ATPases (Futai and Kanazawa, 1983).

The H^+ -ATPase complex consists of two main portions, the F_1 or peripheral membrane portion and the F_0 or integral membrane portion (Figure 2). F_1 , an extrinsic membrane protein, is the catalytic portion and consists of typically five subunits, alpha, beta, gamma, delta, and epsilon (Futai

and Kanazawa, 1983). The F_o portion is a transmembrane complex, which mediates proton translocation between the external and internal compartments of the organelle (or organism) (Futai and Kanazawa, 1983). The F_o portion has not been as extensively studied as the F_1 portion, but the subunits (a, b, and c) from *E. coli* have been identified in biochemical and genetic studies (Foster and Fillingame, 1979; Gay and Walker, 1981; Hansen *et al.*, 1981; Kanazawa *et al.*, 1981). A third portion, called I' , has been identified and functions as a regulator protein preventing the hydrolysis of newly synthesized ATP (Pederson *et al.*, 1981; Ernster *et al.*, 1979).

Studies of the H^+ -ATPase complex began in organelles such as mitochondria and chloroplasts and followed in bacteria. The results from these studies suggested a universal nature for these enzyme complexes, with their structure and basic functions having remained virtually unchanged throughout evolution (Abrams, 1965; Schnebli and Abrams, 1970; Harold *et al.*, 1969; Steck and Fox, 1972; Kagawa and Racker, 1966; Catterall and Pedersen, 1971). The remainder of this review will concentrate on the H^+ -ATPases of bacteria with any differences to the eucaryotic enzymes being mentioned throughout.

1.5.4. Bacterial H⁺-ATPases

The H⁺-ATPases isolated from procaryotic microorganisms are generally very similar, although there are specific differences. Therefore, in order to examine the development of the current ideas about the structure and function of bacterial ATPases, this section will generally provide an overall picture rather than separate reports on individual bacteria.

Pinchot and Racker (1951) provided the first tentative report of oxidative phosphorylation in a fractionated bacterial system using cell free extracts of E. coli. It was later shown, in Alcaligenes, that this activity could be fractionated into membranes and a soluble fraction (Pinchot, 1953) which could be reconstituted to form a coupled phosphorylating system (Pinchot, 1957).

The presence of an H⁺-ATPase localized in the plasma membranes of bacteria was demonstrated in a study of Strep. faecalis (Abrams et al., 1960). Voelz (1964) demonstrated the presence of ATPase activity on membranes in thin sections of E. coli and B. cereus by using a cytochemical stain specific for ATPase activity. Negative staining of membrane preparations of several bacteria has shown the presence of "knobs" protruding from the membrane (Oppenheim and Salton, 1973; Abram, 1965; Asano et al., 1973; Jones and

Redfearn, 1967). These knobs were found to be ATPase, confirming the observation previously made with mitochondria, and they were localized on the inner face of the plasma membrane (Oppenheim and Salton, 1973).

1.5.5. Solubilization and Purification of the F₁ portion

It was first shown, in 1962, that the membrane-bound ATPase activity of M. lysodeikticus could be released by a shock wash using cold water and that this solubilized fraction could be reconstituted in the shock washed membranes producing oxidative phosphorylation (Ishikawa and Lehninger, 1962). Since then, similar procedures using low ionic strength buffers in the presence of EDTA have been employed to solubilize the membrane-bound ATPases of many bacterial species including Strep. faecalis (Abrams, 1965), B. megaterium (Mirsky and Barlow 1971; Ishida and Mizushima, 1969), B. stearothermophilus (Hachimori et al., 1970), Azotobacter vinelandii (Pandit-Hovenkamp 1967; Ackrell and Jones, 1971), M. lysodeikticus (Munoz and Salton, 1969), E. coli (Kobayashi and Anraku, 1972; Davies and Bragg, 1972; Vogel and Steinhart, 1976), Rhodopseudomonas sphaeroides (Melandri et al., 1972), Alcaligenes faecalis (Adolfson and Moudrianakis, 1971), Lactobacillus casei (Biketov et al., 1982), Clostridium thermoaceticum (Mack Ivey and Ljungdahl,

1986), Mycobacterium phlei (Kalra, et al., 1975). In some microorganisms, such Rhodospirillum rubrum (Bengis-Garber and Gromet-Elhanan, 1979) and Mycoplasma landlawii (Cox et al., 1978), it appears that, as with chloroplasts or mitochondria, the ATPase cannot be solubilized by shock wash procedures.

Alternative methods for solubilizing the F₁ ATPase from membranes include (i) extracting the membranes with N-butanol or chloroform (Salton and Schor, 1972, 1974; Ishikawa, 1970), (ii) treatment of the membranes with Triton X-100 (Hanson and Kennedy, 1973; Reed and Raveed, 1972), (iii) passing washed membranes through a French pressure cell (Lucke and Klemme, 1976), and (iv) sonication (Baccarini-Melandri et al., 1970).

Purification methods include precipitation with ammonium sulfate (Futai et al., 1974; Nelson et al., 1974) or polyethylene glycol (Vogel and Steinhart, 1976; Vogel et al., 1978), followed by ion exchange chromatography on DEAE cellulose (Bragg and Hou, 1972, 1975; Futai et al., 1974; Senior et al., 1979) or DEAE Sephadex (Hanson and Kennedy, 1973), and gel filtration using Sepharose 6B (Bragg and Hou, 1972; Vogel and Steinhart, 1976; Vogel et al., 1978) or Biogel A (Futai et al., 1974; Nelson et al., 1974). Most

of these methods yield F_1 preparations that can bind to F_1 -depleted membranes and reconstitute energy-linked reactions and it is this result which determines the feasibility of a purification method. Presumably, a purified enzyme that cannot bind to depleted membranes or reconstitute ATPase activity on the membrane has been altered in some form during preparation.

The F_1 from M. phlei was purified in a single step by chromatography on a Sepharose-ADP affinity column (Kalra et al., 1975).

1.5.6. Solubilization and Purification of the F_1F_0 complex

The F_1F_0 -ATPase complex has been solubilized from E. coli membranes using cholate or deoxycholate (Niewenhuis et al., 1974) and from the membranes of C. pasteurianum and M. phlei using Triton X-100 (Clarke and Morris, 1976; Lee et al., 1976). This complex has also been isolated from the thermophilic bacterium PS3 (Sone et al., 1975).

Purification of this complex has been accomplished by sucrose density-gradient centrifugation (Hare, 1975), DEAE-cellulose chromatography (Foster and Fillingame, 1978), Sepharose-4B column chromatography (Clarke and Morris, 1976) and ADP-Sepharose affinity chromatography (Lee et al., 1976).

The F_1F_0 complex, unlike the solubilized F_1 portion, is sensitive to inhibition by DCCD (N,N'-dicyclohexylcarbodiimide). This chemical is, therefore, used to check for the integrity of the ATPase complex during the purification procedure (Downie *et al.*, 1979). It is often necessary to add phospholipids to the ATPase complex to obtain maximal ATPase activity and DCCD sensitivity (Hare, 1975).

1.5.7. Molecular Weights of F_1 -ATPases

The earliest measurement of the molecular weight of an F_1 -ATPase was made in *Strep. faecalis*. Using rate-zonal sedimentation of a semipurified enzyme preparation in a density gradient, Abrams (1965) found the weight of this enzyme to be 350,000 daltons. A more precise value of the molecular weight was determined by Schnebli and Abrams (1969) and found to be $385,000 \pm 9,000$. To date the reported values for the molecular weights of the purified F_1 from bacterial membranes ranges from 250,000 to 400,000 daltons (Downie *et al.*, 1979).

1.5.8. Electron Microscopic Studies on F_1 -ATPases

Negatively stained electron micrographs of the membranes from mitochondria (Kagawa and Racker, 1966; Hinkle

and McCarty, 1978), chloroplasts (Lien and Racker, 1971; Garber and Steponkus, 1974), *E. coli* (Hinkle and McCarty, 1978), *M. lysodeikticus* (Munoz, *et al.*, 1968), and the thermophilic bacterium PS3 (Kagawa *et al.*, 1976) all show spheres or doughnut-like structures of 8 to 10 nm in diameter attached to the membrane through a portion called the stalk. The F_1 from the thermophilic bacterium PS3 has been crystallized and exhibits a periodic array of the molecules when viewed with the electron microscope (Kagawa *et al.*, 1976; Wakabayashi *et al.*, 1977). Image analysis of this F_1 portion of the enzyme, by light diffraction, showed considerable six-fold and three-fold symmetry and a hexagonal structure with a low-density region, or hole, near the center of the structure (Wakabayashi *et al.*, 1977).

1.5.9. Subunit Composition and Arrangement of F_1 -ATPases

Early analyses of F_1 preparations suggested the presence of only two subunits (Downie *et al.*, 1979). However, it is now generally believed that most bacterial F_1 -ATPases have at least five subunits (alpha, beta, gamma, delta and epsilon in order of decreasing molecular weight) (Futai and Kanazawa, 1980). The molecular weights reported

for these subunits from E. coli fall within the following ranges: alpha, 54,000 to 60,000 daltons (Kobayashi and Anraku, 1972; Hanson and Kennedy, 1973); beta, 52,000 to 56,000 daltons (Vogel and Steinhart, 1976; Bragg and Hou, 1975; Hanson and Kennedy, 1973); gamma, 32,000 to 35,000 daltons (Vogel and Steinhart, 1976; Bragg and Hou, 1975; Hanson and Kennedy, 1973); delta, approximately 21,000 daltons (Vogel and Steinhart, 1976; Bragg and Hou, 1975); epsilon, 11,000 to 13,000 daltons (Hanson and Kennedy, 1973; Bragg and Hou, 1975). F₁ preparations with five subunits have been described for Strep. faecalis (Abrams et al., 1976), S. typhimurium (Bragg and Hou, 1975), R. rubrum (Johansson and Baltscheffsky, 1975; Lucke and Klemme, 1976), M. phlei (Kalra et al., 1975), A. faecalis (Adolfson et al., 1975; Adolfson and Moudrianakis, 1976), and M. lysodeikticus (Huberman and Salton, 1979). It has recently been shown that not all F₁-ATPases possess the typical five subunit profile. The F₁ purified from C. thermoaceticum has four subunits with molecular weights corresponding to 60,000, 55,000, 37,000, and 17,000 daltons in an apparent molar ratio of 3:3:1:1 (Mack Ivey and Ljungdahl, 1986). The F₁ of L. casei has a single subunit (Biketov et al., 1982; Tikhonova et al., 1983) and the enzyme of C. pasteurianum possesses possibly two subunit types (Clarke and Morris, 1976).

The stoichiometry of the different subunits in the five subunit type enzymes is still unclear, and may be of the following order, $\alpha_3\beta_3\gamma_1\delta_1\epsilon_1$ (Bragg and Hou, 1975) or $\alpha_2\beta_2\gamma_2\delta_1\epsilon_2$ and ϵ_2 (Vogel and Steinhart, 1976).

1.5.10. Activation of the F₁-ATPase

Bacterial F₁-ATPases are activated by cations. Mg^{2+} is usually a major activator of these enzymes, however, Ca^{2+} , Mn^{2+} , Zn^{2+} , Co^{2+} , and Cd^{2+} have also been found to activate these molecules in a minor way (Abrams and Smith, 1974). Munoz and Salton (1968) found that Ca^{2+} was the main cationic activator of the F₁ of *M. lysodeikticus* and this activity was inhibited when Ca^{2+} was added.

The solubilized F₁ portion of the ATPase from *M. phlei* is inactive or latent in the hydrolysis of ATP and can be activated by treatment with trypsin (Kalra *et al.*, 1975). Activation by proteases has resulted in the necessity of the inclusion of protease inhibitors during the extraction and purification steps of these enzymes (Abrams and Smith, 1974).

1.5.11. Substrates of F₁-ATPases

The F₁ ATPases of bacteria are generally able to hydrolyse nucleotide substrates other than ATP. They show activity with GTP and to a lesser extent with UTP and CTP. The F₁ of B. megaterium exhibits active hydrolysis of ITP (Mirsky and Barlow, 1971). The E. coli enzyme degrades ITP and ADP (Davies and Bragg, 1972; Hanson and Kennedy, 1973), and Strep. faecalis F₁ hydrolysed ITP and deoxyATP (Abrams and Smith, 1974).

1.5.12. Inhibition of F₁-ATPases

Unlike the F₁F₀ complex, the soluble F₁ enzyme is cold labile (i.e., activity is rapidly lost after storage at 0 to 4°C (Haddock and Jones, 1977)). This property is shared with mitochondrial F₁ (Penefsky and Warner, 1965), but not with the purified ATPases from PS3 and M. phlei (Hakimi et al., 1974; Haddock and Jones, 1977).

A number of chemical inhibitors affect the ATPases of procaryotic microorganisms. DCCD inhibits the ATPase activity of membrane-bound, but not of soluble F₁ (Clarke and Morris, 1976; Harold et al., 1969; Bragg and Hou, 1975). This inhibitor has been shown to react with the F₀ portion of the ATPase (Abrams and Baron, 1970; Abrams et al., 1972).

Bathophenanthroline inhibits the membrane-bound ATPase of E. coli but not the soluble F_1 and can be reversed by the CCCP (carbonyl cyanide *m*-chlorophenylhydrazine) which is an uncoupler (Sun et al., 1975). Uncoupler sensitive inhibition of soluble mitochondrial F_1 has also been observed (Phelps et al., 1975). Oligomycin inhibits the membrane bound F_1 from mitochondria, but does not affect the ATPase activities of most of the bacteria studied to date (Downie et al., 1979). The membrane-bound ATPases of R. rubrum and Rhodopseudomonas capsulata are inhibited by oligomycin (Downie et al., 1979).

Inhibitors that affect the solubilized ATPase (F_1) activity of bacteria are azide (Muñoz, et al., 1968; Evans, 1969), the antibiotic dio-9 (Harold et al., 1969; Bragg and Hou, 1975), quercetin (Futai et al., 1974), and the zinc chelator Zincon (Sun et al., 1975). Aurövertin also inhibits soluble ATPase activity in bacteria (Adolfsen and Moudrianakis, 1976), but unlike its binding to mitochondrial F_1 it does not show fluorescence (Van De Stadt et al., 1974).

1.6. Goals

The purpose of this study was an examination of the polar membrane of C. jejuni to: (i) further elaborate on its ultrastructure from that described by Ritchie et al. (1966), (ii) to analyse minicell formation with the hope of using this fraction to obtain cytoplasmic membrane enriched in the polar membrane structure, and (iii) to determine the function of this structure.

It was recognized that the the component units of polar membrane, the stalks and knobs, were structurally similar to the ATPases of the inner mitochondrial membrane and the cytoplasmic membrane of various bacterial species (Racker, 1967; Harold, 1972). Therefore, a study of ATPase activity in C. jejuni was undertaken to clarify the nature and suggest the function of this interesting structure.

Since other bacteria possess polar membrane a study of the relationship between these structures was also initiated, along with a comparison of the relatedness of the ATPase of C. jejuni to that of other organisms.

CHAPTER 2

Materials and Methods

2.1. Organisms

The polar membrane and F_1 -ATPase of Campylobacter jejuni a clinical isolate obtained from Victoria Hospital London, Ontario (University of Western Ontario Culture Collection no. 1210) was examined in this study. Aquaspirillum serpens MW5 (U.W.O. no. 377) and A. serpens MW11 (U.W.O. no. 374) were used for structural comparisons of the polar membrane with C. jejuni. The following organisms were used for comparison of ATPase activity by histochemical staining i) Escherichia coli K12 (U.W.O. no. 1025), ii) A. serpens MW5, iii) Micrococcus luteus (U.W.O. no. 286, a "lysodeikticus" strain), iv) Deinococcus radiodurans (U.W.O. no. 1071) and v) Bacillus cereus (U.W.O. no. 317). The following organisms were used for comparisons of F_1 -ATPases by immunoblotting i) E. coli K12, ii) A. serpens MW5, iii) Spirillum volutans (American Type Culture Collection no. 19554), iv) Vibrio cholerae CA401 (biotype El Tor, serotype INABA) (provided by T.J. Beveridge, Dept. Microbiology, Univ. Guelph, Guelph, Ontario), v) Methanococcus voltae strain PS (obtained from K. Jarrell, Queens Univ., Kingston, Ontario), and vi) rat liver mitochondria (provided by G. McDougall, Dept.

Biochem., Univ. Western Ontario, London, Ontario)

2.1.1. Cultivation and Harvesting

C. jejuni was cultivated in 500 ml volumes of Brucella Broth (Difco Laboratories, Detroit, Mich.), without shaking, at 40 °C. A growth curve for this organism was prepared by removing 1 ml samples at various time intervals post-inoculation and measuring absorbance at 675 nm. Alternately, for the production of minicells, this organism was grown on Brain Heart Infusion Agar (Difco Laboratories, Detroit, Mich.) at 40°C in an anaerobic jar from which the atmosphere had been removed by a vacuum.

A. serpens MW5 and MW11 were grown on peptone-succinate salts slants (PSS broth containing 1.5% (wt/vol) agar) at 30°C (Hylemon, et al., 1973). The PSS broth, pH 7.2, contained, as a final concentration (wt/vol): 1.0% peptone (Difco Laboratories, Detroit, Mich.), 0.1% succinate, 0.1% magnesium sulphate, 0.1% ammonium sulphate, 0.0002% ferric chloride, and 0.0002% manganese chloride. The succinate and salts were all obtained from Fisher Scientific Co., Fair Lawn, New Jersey. After autoclaving, sterile calcium chloride was added to a final concentration of 0.025% (wt/vol).

E. coli K12 was grown in nutrient broth (Difco Laboratories, Detroit, Mich.) at 37°C with shaking (150

rpm; Psychrotherm incubator-shaker; New Brunswick Scientific Co., New Brunswick, N.J.).

M. luteus and D. radiodurans were grown at 30°C on tryptone-yeast extract-glucose agar slants, pH 7.0, with the final concentrations (wt/vol) of the constituents; 0.5% tryptone, 0.3% yeast extract, 0.1% glucose, and 1.5% agar. The tryptone, yeast extract and agar were obtained from Difco Laboratories, Detroit, Mich., and the glucose from the McArthur Chemical Co., Montreal, Can..

S. volutans was grown at 30°C in MPSS broth, which contained 1.0 g/l $(\text{NH}_4)_2\text{SO}_4$ (McArthur Chem. Co., Montreal, Can.), 1.0 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (Fisher Sci. Co., Fair Lawn, N.J.), 0.002 g/l $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (J.T. Baker Chemical Co., Phillipsburg, N.J.), 5 g/l peptone (Difco Laboratories, Detroit, Mich.), 1.0 g/l succinic acid (Fisher Sci. Co., Fairlawn, N.J.), 0.0002 g/l $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ (Fisher Sci. Co., Fair Lawn, N.J.) and adjusted to pH 6.8 with 2N KOH (Fisher Sci. Co., Fair Lawn, N.J.).

V. cholerae CA401 was grown at 30°C on Brain Heart Infusion Agar slants (Difco Laboratories, Detroit, Mich.).

B. cereus was grown on trypticase soy agar slants (Difco Laboratories, Detroit, Mich.) at 37°C.

Methanococcus voltae was obtained in 10 ml volumes of Balch medium III (Balch and Wolfe, 1979) in sealed

serum bottles with a gas atmosphere of CO₂/H₂ (1:4, vol/vol). The cells were harvested by centrifugation at 3,500 x g for 10 min. at 4°C, and frozen as packed cells at -70°C until used for immunoblotting.

Cells were routinely harvested in the mid-logarithmic phase of growth by centrifugation at 3,500 x g for 15 min. at 4°C and washed in 50 mM HEPES (N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid) buffer at pH 7.2 containing 10 mM MgCl₂ (HEPES-Mg²⁺-buffer). Organisms grown on agar slants or plates were washed off using the aforementioned buffer and centrifuged as above.

Examination of *C. jejuni* cultures for minicell formation was performed as follows; samples (5ml) were collected at 27, 30, 33, 37, and 48 h after inoculation, mixed with an equal volume of glutaraldehyde-acrolein fixative (Burdett and Murray, 1974) for 3 h at room temperature, harvested by centrifugation at 5,000 x g for 5 min. at 4°C, washed 3 times in 5ml volumes of 0.05 M sodium cacodylate buffer, pH 7.5, and enrobed in 2% (wt/vol) Noble agar (Difco Laboratories, Detroit, Mich.) for embedding and sectioning.

2.1.2. Preparation of Cell Envelopes

Envelopes were prepared as follows. Washed cells

were homogenized with a Potter-Elvehjem tissue homogenizer using a Teflon pestle in order to remove their flagella. The homogenized cells were then centrifuged at $3.5000 \times g$ for 10 min at 4°C and the supernatant fraction was discarded. The cell pellets were resuspended in ice cold HEPES- Mg^{2+} buffer containing 10 $\mu\text{g}/\text{ml}$ DNase and 20 $\mu\text{g}/\text{ml}$ RNase. Both enzymes were obtained from the Sigma Chemical Co. (St. Louis, Mo.). The cells were then broken by passage through a French pressure cell (Aminco) at 16,000 lb/in^2 . Whole cells and debris were removed by centrifugation at $3,500 \times g$ for 10 min. at 4°C . The supernatant was then centrifuged at $120,000 \times g$ for 1 h at 4°C in a Ti-50 rotor on a Beckman L-8 ultra centrifuge. The cell envelopes were then washed twice in HEPES- Mg^{2+} buffer by homogenization with a tissue homogenizer and Teflon pestle, and centrifuged at $120,000 \times g$ for 1h each at 4°C .

2.2. Electron Microscopy

2.2.1. Negative Staining

Whole cells or minicells were negatively stained after adhering cells to formvar/carbon coated 200 mesh copper grids by floating the grids on droplets of the samples on a paraffin film. Excess fluid was removed by

touching the grids edgewise to a piece of filter paper. The adherent cells were then washed by floating the grids successively on 5 drops of glass distilled water, stained by floating the grids on a drop of 1% (wt/vol) aqueous uranyl acetate (Merck, Rahway, N.J.), and dried after removal of the excess stain with filter paper as above. Membrane preparation and purified ATPase were stained as above except no washing of the grids was performed and 1% (wt/vol) phosphotungstic acid pH6.8 was used as the stain.

2.2.2. Fixation, Embedding, and Sectioning

Whole cells and minicells were routinely fixed in 5.0% acrolein-0.25% glutaraldehyde (both from Polysciences Inc., Warrington, Pa.) in 0.05 M sodium cacodylate buffer at pH 7.2, at room temperature for 3 h (Burdett and Murray, 1974). The cells were post-fixed, with 1% osmium tetroxide in 0.05 M sodium cacodylate buffer for 1 h at room temperature, washed 3 times in the same buffer, enrobed in 2% Noble agar (Difco Laboratories, Detroit, Mich.), cut into 2 mm blocks and stained with 1% uranyl acetate (British Drug Houses) in distilled water for 2 h at room temperature. The samples were dehydrated in a 30 to 100% (vol/vol) ethanol series and embedded in Spurr embedding medium (Marivac Ltd., Halifax, N.S., Can.). Sections were cut using a Porter-Blum ultramicrotome (Sorval, Norwalk,

Conn.) with glass knives, picked up on 400 mesh formvar carbon coated copper grids and stained for 3 min with 1% (wt/vol) aqueous uranyl acetate and for 6 min. with lead citrate (Reynolds, 1963). Samples were examined in a Phillips EM-300 electron microscope operating at 60 KV. Micrographs were taken on Kodak fine-grain positive film.

Some membrane preparations for embedding were prefixed with 0.5% tannic acid in 0.05 M sodium cacodylate buffer pH 7.2 prior to fixation with 5% acrolein-0.25% glutaraldehyde.

2.3. Visualization of Polar Membrane on Membrane Vesicles by Negative Staining

2.3.1. Stabilization of Polar Membrane on Membrane Vesicles

Polar membrane was stabilized on the membranes by preparing the membranes as described in section 2.1.2, except all buffers used contained 10 mM $MgCl_2$ and 2.5 mM 2-mercaptoethanol. The cell membranes were washed only once prior to negative staining.

2.3.2. Preparation of Lipid-Depleted Membrane Sheets

Cell membranes were prepared as described above (section 2.1.2.) and the lipids were extracted from the

membranes with 1% (wt/vol) cholic acid (Sigma) according to the method of Salton et al. (1968). The extraction buffer was 20 mM Tris-HCl, pH 7.5, plus 1% cholic acid, 10 mM MgCl₂ and 2.5 mM 2-mercaptoethanol. The cell membranes were extracted a total of 6 times with the cholic acid, washed twice in the same buffer minus the cholic acid and negatively stained with 1% phosphotungstic acid, pH 6.8.

2.3.3. Application of a Modified Cell Monolayer Technique of Nermut (1983) for Visualizing the Inner Surface of the Plasma Membrane

Whole cells of *C. jejuni* were harvested, washed and resuspended to a paste like consistency. The cell paste was spread on a piece of freshly cleaved mica (1 x 2 cm). Six 200 mesh formvar carbon-coated copper grids were placed on top of the cell paste, sandwiched with a piece of filter paper (Whatman no. 1, 1.3 x 2.3 cm) and immersed in liquid nitrogen for 5 minutes. The mica and filter paper were then torn apart (under the liquid nitrogen) and the grids adhering to the filter paper were stained immediately with 1% phosphotungstic acid, pH 6.8, to visualize the inner surface of the plasma membrane.

2.4. Separation of Minicells from Whole cells by Sucrose Density Centrifugation

Cultures (24 h) of *C. jejuni* grown on Brain Heart Infusion agar plates, in an anaerobic jar under a vacuum, were harvested by washing the cells from the plates with cold glass distilled water (500ml total volume). The cell suspensions were washed twice in cold distilled water and then layered on 10 ml of a 1.44 M sucrose solution in 15 ml Corex centrifuge tubes and centrifuged at $40,000 \times g$ for 1 h at 4°C . Whole cells pelleted to the bottom of the tube and a diffuse band was seen at the water-sucrose interface. This band was removed, diluted 1:3 with distilled water and centrifuged at $40,000 \times g$ for 1 h. A small pellet was recovered and washed in distilled water. A sample of this pellet was negatively stained for electron microscopic examination and the remainder was fixed and embedded as described in section 2.2.2.

This procedure was repeated with the inclusion of 10 ug/ml DNase, 20 ug/ml RNase and 10 mM MgCl_2 in distilled water.

When this procedure was scaled up, using cells grown in 500 volumes of Brucella broth and washed in distilled water containing DNase and RNase, an 0.1 g (wet weight) sample enriched in minicells was obtained from 10 g (wet weight) of whole cells.

2.5. Extraction and Purification of ATPase from C. jejuni

2.5.1. Shock-Wash Procedure

Extraction of the ATPase from C. jejuni was done according to the method of Vogel and Steinhart (1976). Fifty grams of cells (wet weight) were used for the extraction procedure. Cell membranes were prepared as described previously. The membrane pellets were washed once by resuspending in 300 ml of a 1 mM TRIS-HCl, pH 8.0, buffer containing 2.5 mM 2-mercaptoethanol, 10% (vol/vol) glycerol and 0.5 mM EDTA. Extractions were also done using the above buffer without the EDTA and others included the protease inhibitor phenylmethylsulfonyl fluoride (PMSF) at a concentration of 1 mM. Membranes were collected by centrifugation at 120,000 x g for 90 min. at 4°C. The first wash supernatant fraction was examined for ATP hydrolysing proteins after precipitation with polyethylene glycol as described later in this section. The washed membranes were homogenized in 600 ml of the washing buffer, described above and incubated with stirring for 1 h at room temperature. This solution was then centrifuged at 120,000 x g for 90 minutes. The washed or ATPase depleted membrane pellets were resuspended in 20 ml of 50 mM HEPES-MgCl₂ buffer and stored at -70°C. The supernatant

fraction, containing the released ATPase, was mixed with methanol (4:1) and brought to 50 mM MgCl₂ by the addition of 1 M MgCl₂. The resulting precipitate was removed by centrifugation at 10,000 xg for 20 minutes. To each 100 ml of supernatant 25 ml of 50% (wt/wt), aqueous polyethylene glycol (PEG) (average molecular weight 8,000) (Fisher Sci. Co., Fair Lawn, N.J.) was added slowly beneath the surface while the mixture was stirred. This solution was stirred for a further 20 min. and centrifuged for 30 min. at 10,000 x g.

The supernatant fraction was discarded and the precipitate was resuspended in 50 ml of 50 mM HEPES, pH 7.0, containing 20 % (vol/vol) methanol, 2.5 mM 2-mercaptoethanol and for some experiments 1 mM PMSF. MgCl₂ was added to 50 mM and the resulting precipitate was removed by centrifugation at 40,000 x g for 10 minutes. A 2.5 ml volume of 50% (wt/wt) aqueous PEG (average molecular weight 1000 (British Drug Houses)) was added to each 10 ml of the supernatant fraction. The precipitated protein was collected by centrifugation at 40,000 xg for 15 min. and dissolved in 5 ml of 50 mM HEPES (pH 7.0) buffer containing 20% (vol/vol) methanol or 10% (vol/vol) glycerol and 2.5 mM 2-mercaptoethanol and centrifuged at 10,000 x g for 10 min. to remove any undissolved material. The resulting supernatant was a clear, slightly yellow

solution. This solution was applied to a column of Sepharose 6B (2.6 x 100 cm) previously equilibrated with the above buffer. The protein was eluted at a flow rate of approximately 15 ml/h and 5 ml fractions were collected. The eluted protein was detected by reading absorbance of the fractions at 280 nm. A single protein peak was observed after an elution volume of about 300ml. This peak was pooled and the protein was precipitated by addition of $MgCl_2$ to 50 mM and 2.5 ml of 50% (wt/wt) aqueous PEG (avg. Mr 8000) per 10 ml of the protein solution. The precipitated protein was collected by centrifugation at 40,000 x g for 15 min. and dissolved in 3 ml of 30 mM Tris-HCl (Tris(hydroxymethyl) aminomethane hydrochloride, Fisher Sci. Co., Fair Lawn, N.J.), pH 7.0, containing 20% methanol and 2.5 mM 2-mercaptoethanol. This extraction and purification procedure resulted in enzyme fractions containing 1 to 3 mg protein/ml as determined by the modified Lowry method of Hess *et al.* (1978). These protein fractions were analysed for ATPase activity, homogeneity and subunit composition as described later in this chapter.

This method was also used for the extraction of the ATPase from *E. coli* K12. However, instead of chromatography on Sepharose 6B, the enzyme was partially purified by affinity chromatography on ADP-Sepharose 4B according to the method of Higashi, *et al.* (1975).

2.5.2. Extraction with n-Butanol

The ATPase from *C. jejuni* was also extracted using n-butanol according to the method of Salton, and Schor (1974). Cell membranes were prepared as previously described. The membranes were washed and resuspended in 30 mM Tris-HCl buffer, pH 7.5, plus 5 mM EDTA. The membranes were then collected by centrifugation at 120,000 x g for 90 min., resuspended in the same buffer to give approximately 5 mg membrane protein per ml and extracted at 0°C by mixing with ice-cold n-butanol (Fisher Sci. Co., Fair Lawn, N.J.). Three volumes of n-butanol were added, with stirring, to 4 volumes of the membrane suspension. This solution was allowed to stand for 15 min. in an ice bath. The organic solvent phase was then separated by centrifugation at 27,000 xg for 15 min. at -4°C. The clear upper n-butanol layer was removed and discarded. This extraction process was repeated seven times with fresh ice-cold n-butanol. The final upper n-butanol phase was removed and the aqueous phase above the pellet was carefully withdrawn and placed in a dialysis bag and dialysed against 6 changes (500 ml each) of ice-cold 5 mM Tris-HCL buffer, pH 7.5, to remove any remaining butanol dissolved in the aqueous phase. Dialysis was continued at 4°C until the preparation was free of the odour of butanol. The dialysed aqueous phase was then centrifuged

at 27,000 x g for 30 min. at 0°C and the supernatant was concentrated in an Amicon ultrafiltration apparatus (Amicon Corp.) fitted with a PM10 membrane filter. The concentrated supernatant was then centrifuged at 120,000 x g for 90 min. at 4°C. The pellets were discarded and the supernatants were further concentrated by ultrafiltration. The final protein concentration of the supernatant was estimated to be 3 mg/ml by the modified Lowry method of Hess et al. (1978). This fraction was dialysed against 30 mM Tris-HCl buffer, pH 7.5, and 0.35 g/ml of solid ammonium sulphate was added with stirring at 0°C. The pH of the solution was adjusted to 7.2 with the addition of 1 M NaOH and the suspension was held at 0°C for 1 h. The precipitated protein was collected by centrifugation at 10,000 x g for 20 minutes. The pellet was resuspended in 2 ml of Tris-HCl buffer, pH 7.5, and centrifuged at 27,000 x g for 30 min. to remove any insoluble material. The supernatant was then dialysed against 30 mM Tris-HCl buffer, pH 7.5, and applied to a Sepharose 6B column (2.6 x 100 cm) previously equilibrated in the same buffer. A protein peak was observed after an elution volume of approximately 320ml. This peak was concentrated by ultrafiltration as previously described and had a protein concentration of approximately 1.5 mg/ml. This fraction was then analysed for ATPase activity, homogeneity and subunit composition as described later in this chapter.

2.6. Determination of Protein

Protein concentrations were determined by the modified Lowry method of Hess, *et al.* (1978). Bovine serum albumin (Sigma) was used as a standard

2.7. Electrophoresis and Molecular Weight

Determinations

Polyacrylamide gel electrophoresis under non-dissociating conditions was carried out on 5% polyacrylamide (30:0.8; acrylamide:bis-acrylamide) discontinuous slab gels, with a 3% stacking gel, using the method of Laemmli (1970) as described by Koval and Murray (1981) without sodium dodecyl sulphate. Protein was detected by staining with Coomassie brilliant blue R250 in 25% methanol and 7% acetic acid at room temperature overnight. Destaining was accomplished by diffusion in several changes of 25% methanol-10% acetic acid. Samples were made in the sample buffer described by Laemmli (1970) without sodium dodecyl sulphate, 2-mercaptoethanol or heating. All chemicals for electrophoresis were obtained from Bio-Rad Laboratories (Richmond, Calif.). ATPase activity, in the gels, was detected by the method of Abrams, *et al.* (1976). Lanes to be assayed for ATPase activity were cut from the slab gel

and incubated in 30 mM Tris-HCl buffer, pH 7.5, plus 5.0 mM ATP or alternate substrates and 10 mM MgCl₂ or alternate ions and incubated for 2 h at 37°C. The incubation mixture was then removed and the gels were rinsed twice with distilled water. The gels were then stained for the presence of inorganic phosphate (Pi) with a mixture of 9.4% (vol/vol) perchloric acid, 1% (wt/vol) ammonium molybdate and 0.25% (wt/vol) 1-amino-1 naphthol-4 sulfonic acid (Fisher Sci., Co., Fair Lawn, N.J.). The Pi containing bands were allowed to develop to the point where the gel background remained clear (approximately 20 min). The stain was then removed and the gels were washed twice with distilled water, allowed to air dry for 20 min. and photographed with Kodak Panatomic X film.

Gel analysis under dissociating conditions was carried out on 12% polyacrylamide slab gels containing 0.1% (wt/vol) sodium dodecyl sulphate (SDS-PAGE) or 8 M urea (Urea-PAGE) using the system of Laemmli (1970). Samples for SDS-PAGE were routinely heated in the sample buffer described by Laemmli (1970) at 100°C for 3 minutes. SDS-PAGE gels were stained for protein with Coomassie brilliant blue as described above or by silver staining (Bio-Rad Laboratories, Richmond, Calif.). In some cases, in order to resolve subunit composition, SDS-PAGE was carried out for twice the amount of time it took for

the dye front to reach the bottom of the gel. Samples for Urea-PAGE were prepared in 8 M urea plus 2.5 mM 2-mercaptoethanol.

The molecular weight of the denatured ATPase was determined by comparing its migration in SDS-polyacrylamide gels to that of RNA polymerase (165,000, 155,000, and 39,000 daltons). The molecular weights of cell proteins visualized by immunoblotting were determined by comparing their migration in SDS-polyacrylamide gels to those of the following calibration proteins: beta -galactosidase (116,000), phosphorylase B (97,400), bovine serum albumin (66,000), and carbonic anhydrase (29,000) (Sigma Chemical Co., St. Louis, Mo.)

2.8. Measurement of ATPase Activity

ATPase activity was measured at 37° C in a reaction mixture containing 5 mM ATP (or alternate substrate), 10 mM MgCl₂ (or alternate ion) in 30 mM Tris-HCl buffer (pH 7.0), and 50 to 100 ug of protein in a volume of 1 ml. The reaction was initiated by adding ATP and stopped at various time intervals (0, 30, 60, 105, 120, 150, and 180 min.) by the addition of 200 ul of 10% (wt/vol) trichloroacetic acid (TCA). The precipitated protein was removed by centrifugation in an Eppendorf centrifuge and

500 μ l of the supernatant was removed for detection of inorganic phosphate by the method of Abrams et al. (1976) KH_2PO_4 was used as a standard and P_i levels were determined by reading the absorbance at 650 nm.

One unit of activity was defined as the release of 1 μM of P_i per min., and specific activity was defined as the units of activity per milligram of protein. The effect of inhibitors on ATPase activity was evaluated by incubating the ATPase (soluble and membrane bound) in the assay mixture with the inhibitors at 37°C for 10 min. prior to starting the reaction by addition of the substrate.

2.9. Substrate Specificity

Analysis of substrate specificity of the ATPase from C. jejuni was done using the non-denaturing PAGE activity assay described in section 2.2.8. GTP, CTP, TTP, ITP, AMP, ADP (all from Boehringer Mannheim, Dorval, Quebec, Can.), and glycerol-6-phosphate (Sigma Chemical Co., St. Louis, Mo.) were tested at a level of 5 mM in 30 mM Tris-HCl buffer, pH 7.0, plus 10 mM MgCl_2 .

2.10. Effect of Ions on Enzyme Activation

Ionic activation of the ATPase was analysed by the non-denaturing PAGE method described earlier. All cations

were used at a level of 10 mM. Mg^{2+} , Ca^{2+} , Co^{2+} , Fe^{3+} , Zn^{2+} , Mn^{2+} , Na^{+} , and K^{+} were tested for enzyme activation. Mg^{2+} and Ca^{2+} were tested as a mixture at a level of 10 mM each.

2.11. Cold Lability

Aliquots (100 ug of 2.5 mg protein/ml) of purified ATPase from *C. jejuni* were stored at $-20^{\circ}C$ for 0 to 10 days. One sample was removed every 24 h and assayed for ATPase activity as described in section 2.2.9.

2.12. Effect of Inhibitors on Soluble and Membrane Bound ATPase Activity

N,N'-dicyclohexylcarbodiimide (DCCD) and oligomycin (both from Sigma Chemical Co., St. Louis, Mo.) were tested for their effect on both the soluble and membrane bound ATPase from *C. jejuni*. Inhibition was evaluated by incubating 100 ug protein /ml of the soluble ATPase, or 100 ug protein /ml of the washed membranes (prior to extraction) with the inhibitors (DCCD: 1.0 and 5.0 uM; oligomycin: 10 and 20 uM) for 10 min. prior to the addition of ATP. The enzymes were incubated in the presence of ATP for 30 min., the reaction was stopped by addition of TCA and the levels of P_i released were measured as described in section 2.2.9. Control samples

containing neither inhibitor were used to estimate normal ATPase activity. This activity was considered to be 100% and inhibition was determined by a decrease in activity from this value. DCCD was dissolved in methanol and the control sample for this assay contained an equal concentration of methanol.

2.13. Effect of Trypsin on the Soluble ATPase Activity

Trypsin (Sigma Chemical Co., St. Louis, Mo.), at levels of 0.0, 0.1, and 0.5 mg trypsin per mg of ATPase protein, was incubated in the ATPase assay mixture described in section 2.2.9 for 0, 5, 15, and 30 min. at 37°C. P_i released was then measured as described earlier.

2.14. Reconstitution of ATPase Activity in ATPase Depleted Membranes

Purified ATPase (200 ug protein/ml), prepared as described in section 2.5.1, was added to 400 ug protein/ml of shock-washed or ATPase depleted membranes, which had been resuspended in 30 mM Tris-HCL buffer, pH 7.0, plus 10 mM $MgCl_2$, in a total assay volume of 1.5 ml. This suspension was incubated for 30 min., with agitation, at room temperature. The membranes were then collected by centrifugation for 20 min. in an Eppendorf centrifuge and the supernatant was removed and assayed for protein

content and ATPase activity. The membrane pellet was washed 5 times in the above buffer and assayed for protein content and ATPase activity.

2.15. Histochemical Staining for ATPase Activity

Cells to be histochemically stained for ATPase localization, after the method of Voelz (1964), were harvested in phosphate buffered saline (PBS) pH 7.2 and prefixed for 10 min. at room temperature with 1% glutaraldehyde in PBS. Alternatively the cells were prefixed in 1% formaldehyde for 20 min. at room temperature. The fixed cells were washed 5 times in glass distilled water and resuspended in the ATP medium of Otero-Vilardibo et al. (1963) and incubated at room temperature for 20, 30, and 60 min. The ATP medium contained in final concentrations 8.3×10^{-4} M ATP-disodium (Sigma Chemical Co., St. Louis, Mo.), 0.08 M Tris Malate buffer pH 7.3, 2.5×10^{-3} M $\text{Pb}(\text{NO}_3)_2$, no cations (control), 0.05 M MgCl_2 , or 0.05 M CaCl_2 . After incubation the cells were washed 3 times in glass distilled water and resuspended in 1% osmium tetroxide for 1 h at room temperature, washed 3 times with glass distilled water and enrobed and embedded as described above. The sections were stained with 2% uranyl acetate in 50% (vol/vol) ethanol at 60°C for 10 min.

2.16. Antigen Localization of ATPase in Thin Section

2.16.1. Preparation of Antisera and Isolation of Gamma Globulin (IgG) Fractions

Antisera, specific for the purified ATPase isolated from C. jejuni, was raised in 2.5 kg. New Zealand white female rabbits. A preparative non-dissociating polyacrylamide gel was loaded with 13.3 mg of the purified protein (prepared as described in section 2.5.1) and electrophoresis conducted as usual. The gel was briefly fixed, stained, and destained and the ATPase band excised. The protein-containing gel slice was washed in saline and then completely homogenized by, first, grinding with a mortar and pestle and then by passage through a syringes fitted with successively higher guage needles (18 to 26 guage). The suspension volume was then brought to 5 ml with sterile saline and mixed 1:1 with Freund's complete adjuvant. One ml (approximately 1 mg ATPase) of this antigen preparation was then injected subcutaneously into the neck region of each rabbit. Two weeks later the rabbits were rechallenged with 1 ml of the antigen by intramuscular injection. This procedure was repeated one week later. Five weeks following the initial injection, the blood was obtained from the rabbits by slitting the ear vein and using a mild vacuum collection apparatus.

The blood was allowed to clot at 4°C, and the antisera was stored at -70°C. Normal (preimmune) serum was collected prior to injection with the antigen. Rabbit antiserum specific for the native F₁-ATPase of E. coli K12 was a gift from Dr. S. Dunn, Dept. of Biochemistry, University of Western Ontario, London, Canada.

The IgG fractions were isolated by passing the crude antisera through a Protein A-Sepharose column (Bio-Rad). The bound IgG was eluted in 0.5 ml fractions with 0.1 M glycine-HCl, pH 3.0, into tubes containing 0.5 ml of 0.1 M Tris-HCl, pH 8.0, and the absorbance of the eluted fraction was measured at 280 nm. The eluted IgG fractions were pooled and simultaneously concentrated and dialysed against 10 mM phosphate buffered saline (PBS), pH 7.4, by Amicon ultrafiltration, using a YM30 membrane filter. The IgG samples were stored in 200 ul aliquots at -70°C.

2.16.2. Antigen Localization in Thin section of Spurr Resin-Embedded Cells

Thin sections of whole cells fixed and embedded as described in section 2.2.2. were picked up on naked 400 mesh nickel grids and antigen localization was carried out according to the method of Roth (1982). The grids with the attached thin sections were placed face down on a droplet of 4% (wt/vol) ovalbumin (Calbiochem, San Deigo, Calif.) in PBS (pH 7.4) for 5 min. at room temperature. the grids were then transferred, without rinsing onto

drops of the antibody solutions for 2 h at room temperature or 18 h at 4°C. These procedures were carried out in a moist chamber. Various dilutions of the antibody solution (in PBS plus 1% ovalbumin) ranging from 1:100 to 1:100,000 were used. The sections were then "jet-washed" by a mild spray of filter sterilized PBS (i.e. after passage through a 0.2 μ m millipore filter) from a plastic spray bottle, placed in PBS for about 2 min., "jet-washed" a second time with PBS and blotted by holding the grids edgewise on a piece of filter paper. The grids were then placed on drops of a 20 to 25 fold dilution of a 5 or 10 nm protein A colloidal gold stock solution (Janssen Pharmaceutica Life Sciences Products, Beerse, Belgium) for 30 to 60 min. at room temperature in a moist chamber. The grids were then "jet-washed" with PBS for 1 min., immersed for 5 min. in PBS in a porcelain depression dish, "jet-washed" and immersed in PBS a second time and finally "jet-washed" with filter sterilized (0.2 μ m millipore filter) double distilled water. The grids were never allowed to dry during the whole procedure. Finally, the section were stained with 5% aqueous uranyl acetate and lead citrate (Reynolds, 1963) for 5 and 10 min. respectively.

Controls for this method were as follows; i) PBS plus 1% ovalbumin was used in place of the antibody solution, and ii) diluted normal pre-immune IgG was used

in place of the antibody solution.

For some localization studies thin section of whole cells which had been post-fixed with osmium tetroxide were etched for 2 h at room temperature by floating grids with the attached sections on droplets of 20% (vol/vol) hydrogen peroxide (Fisher Scientific Co., Fair Lawn, N.J.)

2.16.3 Low-Temperature Embedding for Antigen Localization

Lowicryl KM4 (Chemische werke Lowi, D8264 Waldkraiburg, F.R.G.), an embedding resin which polymerizes at very low temperatures, was used as an alternative to Spurr resin embedding to preserve antigen structure for antibody localisation in sections.

Cultures were fixed for low temperature embedding using 5% acrolein-0.25% glutaraldehyde in 0.05 M sodium cacodylate buffer as previously described. However, post-fixation with osmium tetroxide and block staining with 1% uranyl acetate was omitted. Dehydration with ethanol and embedding in Lowicryl KM4 was done according to the method of Roth, *et al.*, (1981). The fixed cells were enrobed in 2% Noble agar as previously described, cut into 2 mm blocks and dehydrated in 30% (vol/vol) ethanol at 0°C (on crushed ice) with agitation for 30 min. The tissue blocks in the 30% ethanol were then transferred to

-22°C (Crushed ice:NaCl, 3:1 (wt/wt)) and dehydration was continued with 50%, 70%, 95% and 2 times 100% ethanol solutions for 1 h each at -22°C. The tissue blocks were then transferred to a precooled 1:1 (vol/vol) Lowicryl KM4:100% ethanol for 1 h at -22°C. This was followed by transfer of the blocks to a 2:1 solution of resin in 100% ethanol for 1 h at the same temperature and finally by transfer to pure resin for 1 h at -22°C. The tissue blocks were then placed in fresh resin and held at -22°C overnight. The resin was composed of 2.5 g of crosslinker A, 17.3 g of monomer B and 0.1 g of initiator C. Fresh resin was prepared, precooled to -22°C and placed in capped BEEM capsules leaving no air space at the top. The tissue blocks were transferred to the prefilled BEEM capsules and allowed to equilibrate for 10 minutes. These steps were performed in a precooled styrofoam chest which was fitted with a long wave ultraviolet lamp in its lid. This UV chamber was kept in a 4°C walk in cold room. The inside of the UV chamber was lined with aluminum foil as a reflector. The BEEM capsules containing the tissue blocks were placed in the bottom of the UV chamber (24 cm from the UV source), the chamber was cooled to -22°C with a crushed ice:NaCl, 3:1 mixture and curing by UV irradiation was carried out for 24 h at this temperature and for 48 h at room temperature. The blocks were sectioned as.

previously described. The sections were picked up on naked nickel grids, incubated with antibody and protein A colloidal gold (5 nm) as previously described and finally stained with 5% aqueous uranyl acetate for 30 min. at room temperature.

2.16.4. Antigenic Relatedness of ATPase from *C. jejuni* to ATPases of other Bacterial Species and Rat Liver Mitochondria

2.16.5. Immunoblotting

Whole cell suspension of *C. jejuni*, *E. coli* k12, *V. cholera*, *A. serpens* MW5, *S. volutans*, *M. voltae*, and rat liver mitochondria were prepared in glass distilled water. The protein content of each suspension was measured by the modified Lowry method of Hess *et al.* (1978) for proteins insoluble in 1N NaOH. 150 ug protein of each cell type was electrophoresed on a 12% SDS-polyacrylamide gel and then electrophoretically transferred to nitrocellulose paper (40 V, 3 h) by the method of Towbin, *et al.* (1979). The protein transferred to the nitrocellulose paper was visualized by briefly staining (2 min.) with 1% Amido black in 25% methanol, 10% acetic acid and destaining in 10% methanol, 10% acetic acid. The individual lanes were cut out and washed, first with distilled water, then with

PBS (pH 7.4). The nitrocellulose strips were then washed in PBS overnight at 4°C. The PBS wash solutions were removed and any available non-specific antibody binding sites were blocked by incubation of the strips in 1% (wt/vol) BSA in PBS for 2 h with gentle agitation. The strips were then washed 6 times in PBS for a period of 40 min. and incubated for 2 h in PBS containing either IgG prepared against the ATPase from C. jejuni (10 ug protein/ml), IgG prepared against the F₁-ATPase of E. coli K12 (10 ug protein/ml), or preimmune IgG (10 ug protein/ml). The antibody solutions were then removed, the strips were washed 8 times in PBS over 1 h with gentle agitation, and incubated for 2 h with goat anti-rabbit IgG-horse radish peroxidase conjugate (Bio-Rad) (8 ug/lane). The strips were washed overnight with PBS and the bound IgG was visualized with 4-chloro-1-naphthol. The strips were rinsed with distilled water, allowed to air dry, and photographed using Kodak Panatomic X film.

CHAPTER 3

RESULTS

3.1. Growth Characteristics of *C. jejuni*

A growth curve for *C. jejuni* cultivated in 500 ml volumes of Brucella broth, without shaking, at 40°C, is shown in Figure 3. This organism showed very little lag in growth when a large inoculum (100 ml of a 24 hour culture) was used and the culture was not disturbed for the first five hours of growth. The exponential phase of growth lasted approximately 15 hours and the culture entered stationary phase at approximately 20 hours post inoculation. Growth under these circumstances provided approximately 2.0 g of cells (wet weight) per 500 ml culture.

3.2. Ultrastructure of Polar Membrane

3.2.1. Location in Whole Cells

Thin sections of whole cells of *C. jejuni* from a 24 hour culture showed the typical helical shape and structures associated with this organism (Figure 4). A polar flagellum and adjacent polar membrane was visible at

the poles of the cell underlying the plasma membrane as described by Ritchie et al. (1966). Spherical inclusion bodies, with densely stained perimeters and lighter interiors, were seen in the polar regions of the cell and appeared to be located centrally within the region where polar membrane was found (Figure 4). The polar regions also contained a lower concentration of ribosomes than the rest of the cytoplasm (Figure 4) similar to the observation of Murray and Birch-Andersen (1963) in A. serpens. Figure 5 allows a comparison of the location and ultrastructure of polar membrane in A. serpens MW11. This organism possesses a polar tuft of flagella (not visible in this micrograph), polar membrane underlying the plasma membrane, and a ribosome free area at the poles.

A cross section through the polar region of C. jejuni suggests that the polar membrane structure covers the entire inner surface of the plasma membrane (Figure 6), whereas, a cross section through the pole of A. serpens MW5 shows the polar membrane to be located in patches (Figure 7).

Polar membrane has also been noted at the site of septum formation (Figure 8), which is in agreement with the observations of Murray and Birch-Andersen (1963) and Ritchie et al. (1966). However, the location of polar

membrane at this site is not a consistent result (Figure 9), as it is not known when polar membrane is laid down at the poles.

3.2.2. Association of Polar Membrane with the Flagellar Basal Complex

The flagellar basal complex in *C. jejuni*, when dissociated from the membranes, shows the typical four ring structure associated with Gram negative basal complexes and also possesses concentric membrane rings (CMR's) between the L and the P rings (Ferris and Beveridge, 1984). Figures 10 through 15 are high magnification micrographs of longitudinal thin sections of the polar region of *C. jejuni*. When the section passes through the centre of the basal complex it is possible to distinguish the location of the CMR's, the P ring and the M ring (Figures 10, 11, 12, and 13), however, an off centre cut blurs the position of the rings within the cell envelope (Figures 14 and 15). The position of the rings observed in these micrographs and the presence of fibres extending from the M ring (Figures 10 and 11) are in agreement with the observations of Coulton and Murray (1977, 1978) and Ferris and Beyeridge (1984).

Figures 10 through 15 also illustrate the association of polar membrane with the basal complex.

Polar membrane encroaches on the lateral margin of the basal complex to varying degrees, as can be seen by comparing Figures 10 through 15. In Figure 10 the polar membrane structure lines the inner surface of the plasma membrane starting approximately 800 nm away from the basal complex. In Figure 11 the structure begins approximately 600 nm down the plasma membrane from the basal complex and in Figures 12 through 15 the structure can be seen next to the basal complex but not covering the area where the M ring is located.

A distinct ribosome free area can be seen in Figures 10, 11, 12, 14 and 15. As well, a densely stained inclusion body with an ill defined perimeter was observed in the polar region (Figure 15).

3.2.3. Fine Structure and Dimensions of Polar Membrane

3.2.3.1 Observations on Thin Sections of *C. jejuni*

Polar membrane appears, on thin section, as a densely staining line approximately 200 nm internal to the plasma membrane and appears to be connected to the plasma membrane by bands or striations (Figures 4 and 6).

Observations of this structure in *A. serpens* MW11 clearly show the presense of stalks arising out of the plasma membrane joined to globular structures at the apposing end

(Figure 5). The structure in *C. jejuni* has not been as clearly resolved on thin section, but figure 16 shows a high magnification micrograph of this structure. Globular units or heads can be seen in the densely staining line 200 nm from the plasma membrane and stalks joining these heads to the plasma membrane are partially resolved. The heads are approximately 5 to 6 nm in diameter and the stalks appear to be half this size although measurement of these structures is hampered by the poor resolution.

It should also be noted that the plasma membrane in the polar membrane region has a smoother almost rigid form when compared to the lateral plasma membrane (Figures 10 to 16).

3.2.3.2. Observations by Negative Stain on Membrane preparations of *C. jejuni*

Negatively stained preparations of unwashed *C. jejuni* cell envelopes obtained by French pressure cell disruption showed the presence of small doughnut-shaped structures on the membranes associated with the flagellar basal complex. (Figure 17a). These structures corresponded in size (5 to 6 nm in diameter) and location to the globular head units seen on thin section. These structures were also observed in an intact polar vesicle (Figure 17b) further associating them with the polar membrane structure seen on thin section.

These structures were not frequently seen in washed membrane preparations. Occasionally they were observed in enclosed vesicles of membrane from the polar region, recognized because of the presence of CMR's (Figure 18).

The inclusion of 2.5 mM 2-mercaptoethanol and 10 mM $MgCl_2$ in the 50 mM HEPES wash buffer increased the frequency of observation of these structures on membrane preparations. Figure 19a shows a membrane vesicle partially covered with small dough-nut shaped structures (5 to 6 nm in diameter) which appear to be arranged in an hexagonal pattern.

These structures were also observed when whole cells were cleaved by a modification of the monolayer technique of Nermut (1983) and the resulting membrane fragments were negatively stained (Figure 19b). This micrograph also shows the size relationship of these structures to the flagellum.

When membrane preparations were washed in a high ionic strength buffer containing 2-mercaptoethanol as described previously and the membrane lipid was extracted with cholic acid (Salton, *et al.*, 1968) a fine hexagonal array was observed in the lipid depleted membranes (Figure 19c and d). This array possessed repeating units of approximately 3 nm in diameter and may represent the

insertion points of the stalks of the polar membrane structure in the plasma membrane.

These structures were not seen when the membrane preparations were fixed and embedded for thin sectioning. It is possible to maintain the polar membrane structure on the plasma membrane, in crude membrane preparations, by using the mordant tannic acid, however, the fine structure of stalks and knobs is obscured (Figure 20).

3.3. Minicell formation by *C. jejuni*

Electron microscopic examination of cells harvested at 27 through 33 hours post inoculation (stationary growth phase) showed the typical helical shape associated with *C. jejuni* and a normal constrictive form of cell division (as shown in Figures 4 and 9). However, among these normal cells were a few much smaller structures (0.1-0.3 μ m in diameter) bearing flagella (Figure 21). Thin sections of these structures showed that these minicells were bounded by outer membrane and by a plasma membrane (Figure 22). The interior contained few ribosomes and no obvious nucleoplasm. The membranes appeared to enclose these minicells entirely and on their inside was a structure identical to polar membrane.

Examination, by thin section, of a series of whole cells collected at 33, 37, and 48 hours after inoculation

provided an increased proportion of minicells to whole cells and allowed deduction of the events involved in forming the minicells. The apparent initial event in their formation involved increasing degrees of constriction of the plasma membrane, resulting in the segregation of the polar region of the cell (Figure 23). This constriction of the plasma membrane without constriction of the outer membrane (as observed in normal cell division) was also observed by negative stain (Figure 24). The profile of the peptidoglycan layer in the cell wall, expected to lie between the outer membrane and the plasma membrane, was not often seen on thin sections of young cells (Figure 4) and could not be located on thin sections of the segregating terminal regions of the minicells (Figure 24). The terminal events in the formation of these structures were not observed.

This phenomenon commonly observed in older cultures entering stationary phase, also occurred when younger cultures (18-24 hours) were subjected to adverse conditions, such as atmospheric levels of oxygen after growth on agar plated under a vacuum or resuspension of cells grown in broth in glass distilled water.

These minicells could be partially separated from the whole cells by differential centrifugation in a discontinuous sucrose gradient (30% (w/w) and 60% (w/w)).

However, the quantity of the minicells harvested in reasonably pure form was too small to allow biochemical analysis and the preparations were usually contaminated with whole cells, flagella and membrane fragments (Figure 25). This result was probably due to the sticky nature of *C. jejuni* cells and minicells, which prevented a clean isolation by gradient centrifugation, even in the presence of DNase and RNase. Figure 26 is a high magnification of a negatively stained minicell. This micrograph shows the presence of a hexagonal array within the structure similar to that observed in lipid depleted membranes.

3.4. Localization of ATPase Activity in Whole Cells by Histochemical Staining

Thin sections of *C. jejuni* cells prefixed with 4% (v/v) glutaraldehyde and incubated with ATP and Mg^{2+} cations were found to have heavy deposits of lead phosphate in the polar regions and light deposits elsewhere in the cell (Figure 27a-d). Spheroplasts of *C. jejuni* also showed localized deposits of lead phosphate along the inner surface of the plasma membrane (Figure 28a and b). Cells incubated with ATP and Ca^{2+} cations were found to have light lead phosphate deposits around the plasma membrane and the inclusion granule seen in the polar region (Figure 29a and b).

This histochemical staining method was also performed on five other bacterial species for comparison of the staining patterns. E. coli, when incubated with ATP- Ca^{2+} was found to have lead deposits randomly distributed throughout the cell and around the plasma membrane (Figure 30a and b). when incubated with ATP- Mg^{2+} no internal lead deposits were seen (Figure 30c). B. cereus, when incubated with ATP- Mg^{2+} , was found to have discrete deposits also randomly located throughout the cell (Figure 31). M. luteus, when incubated with ATP- Ca^{2+} showed lead deposits around the cell lining the plasma membrane (Figure 32a), with ATP- Mg^{2+} the lead deposits were localized in patches internal to the plasma membrane (Figure 32b). D. radiodurans showed localized patches of lead deposits when incubated with ATP- Mg^{2+} (Figure 33a and b) and a random distribution of lead phosphate throughout the cytoplasm when ATP- Ca^{2+} was used (Figure 33c). A. serpens when incubated with ATP- Mg^{2+} showed deposits randomly distributed throughout the cell (Figure 34a), however, when ATP- Ca^{2+} was used these deposits of lead phosphate were heavier at the poles and patchily localized around the plasma membrane (Figure 34b).

2

Micro

3.5. Isolation and Characterization of an F₁-like ATPase from C. jejuni

3.5.1. Analysis of ATP hydrolysing proteins in shock-washed membrane supernatants

The protein in the initial shock-wash supernatant from washed membrane preparations was examined after precipitation with polyethylene glycol as described by Vogel and Sternhart (1976). This protein fraction when analysed on a 5% non-denaturing polyacrylamide gel contained one major protein and three to four minor protein bands (Figure 35 lanes a and e). The major protein species was able to degrade ATP releasing P_i when incubated with 5mM ATP plus 10 mM MgCl₂ (Figure 35 lane c). ATP was not degraded by this protein when incubated with ATP alone (ie. no cations) or with ATP plus 10 mM CaCl₂ (Figure 35 lanes b and d). This protein could not degrade GTP in the presence of either Mg²⁺ or Ca²⁺ ions (Figure 35 lanes f and g). Two of the minor protein species hydrolysed ATP releasing P_i with Ca²⁺ ions being a potent activator and Mg²⁺ activating to a lesser degree (Figure 35 lanes c and d). One of the minor protein species degraded GTP releasing P_i in the presence of Ca²⁺ ions (Figure 35 lane g).

78

These minor ATP hydrolysing protein species were not present in the second shock-wash supernatant, whereas the major ATP hydrolysing protein was present in relatively large amounts. The specificity, relatively high molecular weight (as judged by the position of the protein in a 5% non-denaturing polyacrylamide gel), and isolation by shock wash suggested that this protein was a candidate for an F₁-like ATPase and was therefore selected to be purified and characterized.

3.5.2. Purification of the F₁-like ATPase

The protein fraction isolated from the second shock-wash supernatants of cell membranes by successive cuts with decreasing molecular weights of polyethylene glycol was applied to a column of Sepharose 6B. One protein peak was eluted. The concentrated protein sample obtained from the chromatography was applied to a 5% non-denaturing polyacrylamide gel and migrated as a single band (Figure 36a) demonstrating ATPase activity (Figure 36b). Table 1 shows that a four fold increase in specific activity of the enzyme was obtained after purification.

The specific activity of this protein fraction, when incubated with 5 mM ATP and 10 mM MgCl₂, was 0.2-0.3 units/mg of protein.

Table 1: Purification of Soluble ATPase

	Specific Activity
	(units activity/ mg protein)

Normal Membranes ^a	0.07 ± 0.02 ^d
-------------------------------	--------------------------

PEG concentrated

Soluble ATPase ^b	0.16 ± 0.04 ^d
-----------------------------	--------------------------

Sepharose 6B

Purified ATPase ^c	0.30 ± 0.03 ^d
------------------------------	--------------------------

a. Membranes prior to enzyme extraction. 400 ug protein/ml.

b. Polyethylene glycol 1000 (Fisher Sci.) 400 ug protein/ml.

c. Protein from single peak eluted from sepharose 6B column 400 ug protein/ml.

d. Standard error on the means of three experiments.

3.5.3. Electron Microscopy of ATPase

Negatively stained preparations of the purified ATPase, obtained after column chromatography, revealed the presence of dough-nut shaped particles 5-6 nm in diameter (Figure 37).

3.5.4. Reconstitution of ATPase Activity on Depleted Membranes

ATPase activity could be reconstituted on depleted membranes by addition of 200 ug/ml of the purified ATPase to 400 ug of protein per ml of shock-washed membranes (Table 2).

3.5.5. Subunit Composition

SDS and urea-PAGE revealed a single subunit with a molecular weight of approximately 75,000 Daltons (Figure 38). Silver staining of the gels failed to reveal other subunits. Furthermore, the single subunit band could not be resolved into 2 bands by under loading the gel or by running the gel for twice the amount of time required for the dye front to reach the bottom of the gel. When the F₁-ATPase was extracted and partially purified from E. coli K12 using the method of Vogel and Steinhart (1976), five subunits were obtained (data not shown).

Table 2: Reconstitution of ATPase Activity in Depleted Membranes

	Specific Activity ^b
Normal Membranes ^c	0.08 ± 0.03 ^f
Depleted Membranes ^d	---
Soluble ATPase	0.30 ± 0.05 ^f
Reconstituted Membranes	0.19 ± 0.04 ^f
Reconstituted Soluble ATPase Supernatant	0.10 ± 0.03 ^f

- a. Assay system contained 200 ug soluble ATPase protein/ml and 400 ug depleted membrane protein/ml.
- b. Units of activity/mg of protein.
- c. Membranes prior to enzyme extraction. 400 ug protein/ml.
- d. 400 ug protein /ml
- e. ATPase activity was undetectable.
- f. Standard error on the mean of three experiments.

Extraction of the *C. jejuni* ATPase with n-butanol as described by Salton and Schor (1974) also resulted in an enzyme with only one subunit, and a specific activity of 0.21 units per mg of protein.

3.5.6. Inhibition of membrane-bound and soluble F₁-ATPase Activity

The ATPase was examined for sensitivity to DCCD, oligomycin, and trypsin. The membrane-bound ATPase was sensitive to 5 μ M DCCD, but not to oligomycin, whereas the soluble purified enzyme was not inhibited by either chemical (Table 3). Trypsin (0.1-0.5 mg trypsin/mg protein for 5, 10, and 30 min. at 37°C) completely inhibited the activity of soluble ATPase (Table 4).

3.5.7. Enzyme activation and specificity

The purified ATPase required Mg^{2+} ions (10 mM) for activity (Figure 39). Ca^{2+} , Co^{2+} , Fe^{3+} , Zn^{2+} , Mn^{2+} , Na^{+} , and K^{+} did not individually activate the enzyme. As well, when Ca^{2+} and Mg^{2+} were combined at equal concentrations (10 mM) the enzyme activity was totally inhibited.

The ATPase was able to degrade ATP, releasing inorganic phosphate (P_i), but did not degrade ADP, GTP, or

Table 3: Effects of inhibitors on the membrane bound and purified ATPase from C. jejuni.

Inhibitors ^a	Concn (uM)	Activity ^b	
		Membrane Bound ATPase	Soluble ATPase
DCCD	1	84	91
	5	52	90
Oligomycin	10	89	92
	20	91	94

a. A 100 ug sample of either washed membranes or purified (soluble) ATPase was incubated in the regular assay mixture.

b. The control enzyme activity in the absence of inhibitors. DCCD was dissolved in methanol, the control contained an equal volume of methanol.

Table 4: Effect of Trypsin on Soluble ATPase Activity.

Concn		Specific Activity ^a			
Trypsin					
(mg/mg	Time (min.)	0	5	10	30
ATPase					
protein)					
0.0		---b	0.04	0.09	0.25
0.1		---b	---c	---c	---c
0.5		---b	---c	---c	---c

a. Units of activity/mg soluble ATPase protein.

b. Zero time controls.

c. Activity not detectable.

glycerol-6-phosphate (Figure 40). AMP, CTP, ITP, and TTP were also not hydrolysed by the enzyme.

The soluble enzyme lost approximately 50% of its activity upon storage at -20°C for 24 h, and was stable at 4°C for one week when stored in 10% (v/v) glycerol (Table 5).

3.6. Immunocytochemical Localization of ATPase from *C. jejuni*

Antibodies prepared against the purified Mg^{2+} activated ATPase as described in the methods chapter were used to localize this enzyme in whole cells using the post-embedding technique of Roth (1982). The bound antibodies were visualized by secondary labeling with protein A coated colloidal gold. The labeling pattern seen in thin sections of whole cells fixed with acrolein-glutaraldehyde (Burdett and Murray, 1974), post-fixed with osmium tetroxide, and embedded in Spurr's resin, are shown in Figures 41 and 42. These micrographs show definite gold label at the poles of the cells, but label also appears throughout the cytoplasm. Figure 41b shows the gold binding to a thin section of a minicell. Figures 41c and 41d show localized binding of the gold to spheroplasts of *C. jejuni* much like the pattern seen on histochemically stained spheroplasts (Figs. 28a and b). When the thin sections were etched with hydrogen peroxide, the pattern of gold label was more defined. Figure 43a and b, shows the label to have bound to the polar regions of the cell, as well as throughout the cytoplasm. The

Table 5: ~~Effect of~~ Freezing (-20° C) on Soluble ATPase
Activity.

Specific Activity	
Time (days)	(units activity/mg ATPase protein)
0	0.23
1	0.11
2	0.10
4	0.08
6	0.07
10	0.07

gold label does, however, appear to have bound in clusters in the polar regions and as isolated particles elsewhere in the cell. Control sections incubated with either preimmune IgG or no antibody (1% ovalbumin) showed minimal to no background labeling.

The gold labeling patterns seen in thin sections of whole cells fixed with acrolein-gutaraldehyde (Burdett and Murray, 1974) and embedded in Lowicryl KM4 are shown in Figures 44 and 45. Figure 44a and b show clustering of the gold at the pole of a cell, as well as isolated particles bound throughout the cytoplasm. Figure 45 shows a tangential section of the polar region of a cell with a large cluster of bound protein A colloidal gold particles. Isolated particles are also visible randomly distributed throughout the cell.

3.7. Antigenic Relatedness of the ATPase from *C. jejuni* to ATPases of other Bacterial Species and Rat Liver Mitochondria (Immunoblotting)

Crude antisera raised against the ATPase isolated from *C. jejuni* reacted with the purified ATPase in the Ouchterlony double diffusion assay and by crossed immunoelectrophoresis (data not shown). The IgG was isolated from the crude antisera and preimmune sera and the Western blotting method was used to test for

cross-reactivity with whole cell proteins of C. jejuni, E. coli, V. cholera, S. volutans, M. voltae, A. serpens MW5, and rat liver mitochondria, separated on a 12% SDS polyacrylamide gel. The IgG raised against the ATPase from C. jejuni reacted with four proteins from the whole cell extract of C. jejuni (Figure 46, lane f), the major reactive protein being approximately 66,000 daltons in molecular weight and the three minor proteins running below this protein in the gel. This antibody cross reacted with two proteins from the whole cells of A. serpens MW5, one corresponding to the major reactive protein in C. jejuni and one of a higher molecular weight (Figure 46, lane d). Cross reactivity was also observed with three proteins from E. coli K12 (Figure 46, lane b). One corresponding to the high molecular weight protein seen in A. serpens MW5, one corresponding with the major reactive protein in C. jejuni, and one of a lower molecular weight (Figure 46, lane b). This antiserum did not cross react with any cell proteins from V. cholera, S. volutans, M. voltae or rat liver mitochondria (Figure 46, lanes c, e, g, and h). The preimmune IgG did not react with cell proteins from any of the organisms tested.

IgG isolated from crude antiserum raised against the F₁-ATPase of E. coli K12 reacted with six proteins in the whole cells of E. coli K12 (Figure 47, lane b). This IgG

cross-reacted with one protein species in each of the whole cell preparations of V. cholera, A. serpens MW5, S. volutans, and rat liver mitochondria (Figure 47, lanes c, d, e, and h), but did not cross react with any whole cell proteins from either C. jejuni or M. voltae (Figure 47, lanes f and g).

Chapter 4

Discussion

The existence of polar membrane as a structure has been known for the past twenty four years. However, previous studies of this interesting structural complex associated with cytoplasmic membrane have limited themselves to trying to define its ultrastructure and compiling lists of organisms possessing it. The fine ultrastructure at the macromolecular level, biochemical composition and function(s) of polar membrane have not been investigated to date. Its location at the poles of the cell surrounding the flagellar insertion point has led to the hypothesis that polar membrane functions in energy transductions (Murray and Birch-Andersen, 1963). It has also been suggested that this structure serves as an anchor to which the flagella are attached in organisms possessing a polar tuft of flagella (Cohen-Bazire, 1967; Remsen, et al., 1968), but little else is known about it even though it is found in a variety of lophotrichous Gram negative bacteria.

This study was undertaken to clarify the nature and suggest the function of polar membrane in *C. jejuni*. This

bacterium was chosen for this work for the following reasons: it possessed polar membrane, it had a single polar flagellum, its poles were slightly tapered allowing for high resolution on thin section, and it was found to produce minicells from its polar regions with cytoplasmic membrane enriched in the polar membrane structure.

Recognition that the component units of polar membrane, the stalks and knobs, were structurally similar to the ATPase of the inner mitochondrial membrane and the cytoplasmic membrane of various bacterial species (Racker, 1967; Harold, 1972) led to a study of the ATPase activity of this organism and its relation to the polar membrane structure.

4.1. Ultrastructure of Polar Membrane

The basic structure of polar membrane in *C. jejuni* was previously described by Ritchie et al. (1966). The results obtained from thin sections of whole cells in this study were in agreement with the former work. Comparison of the structure seen on longitudinal thin sections and cross sections through the polar region suggested that polar membrane lined the inner surface of the cytoplasmic membrane at the poles of the cell in the form of a cylindrical cap. The cytoplasm contained within this area was found to contain fewer ribosomes than the rest of the

cytoplasm; an observation previously noted in *A. serpens* by Murray and Birch-Andersen (1963). The spherical inclusion body seen within this region, which was densely stained with lead citrate, may represent a phosphate storage granule and may be related to the function of polar membrane.

Visualization of polar membrane on the cytoplasmic membrane close to the site of septum formation, although not a consistent result, suggests it is laid down prior to daughter cell separation and may point to its requirement in the normal function of a young cell.

The association of polar membrane with the flagellar basal complex was further defined and found to encroach on the lateral margin of the basal complex to varying degrees from cell to cell. This structure was never seen to cover the area in the basal complex where the M ring was located indicating there was no visible contact or linkage with this organelle, despite the earlier speculations that there might be such a linkage. It should also be noted that polar membrane was sometimes observed at random sites along the cytoplasmic membrane in *C. jejuni* and not associated with the flagellar insertion site.

The component units of polar membrane in *C. jejuni* were found to be smaller in dimensions than those seen in *A. serpens* MW11 and those described for polar membrane in

E. mobilis (Remsen, et al., 1968). The stalks were 2-3 nm in diameter, 5 nm in length and the knobs were 5-6 nm in diameter in C. jejuni, whereas, in these other organisms the stalks were 3-4 nm in diameter, 5 nm in length and the knobs were 8-9 nm in diameter. This observation may account for the difficulty in resolving the components of this structure on thin sections of C. jejuni and suggests that although polar membrane is found in many Gram negative bacteria, suggesting a similar function, the size or form of component macromolecules may vary across the genera.

The impression that the cytoplasmic membrane in the region of the polar membrane is of a more rigid character than the rest of the lateral membrane, as seen both on thin sections of whole cells and in fractionated cell envelope preparations, suggests this portion of membrane is different from the rest of the lateral cytoplasmic membrane and may simply be a result of an increased protein to lipid ratio reflecting the insertion site of the polar membrane stalks.

The observation of arrays of dough-nut shaped particles on membrane fragments near the flagellar insertion point and in intact polar vesicles suggested that these structures were components of the polar

membrane complex, because they corresponded in size (5-6 nm), form, and location to the innermost part of the structure seen in thin sections. These complex particles appeared to be arranged in an hexagonal array on the inner surface of the cytoplasmic membrane. Ritchie, et al., (1968) suggested that the polar membrane structure was arranged in a honey comb lattice and these results support this hypothesis.

The fine (3 nm repeat) hexagonal array seen on lipid depleted membrane sheets may represent the insertion sites in the cytoplasmic membrane of the stalk portion of the polar membrane. This structure had almost a crystalline appearance and was similar to the two dimensional crystals of ATPase seen in sarcoplasmic reticulum membrane after induction with vanadate or inorganic phosphate (Skriver, et al., 1981; Maunsbach, et al., 1982; Herbert, et al., 1982; Dux and Martonosi, 1983a and b). This suggested that polar membrane might be a naturally occurring two dimensional membrane crystal and this formation might result in an increase or decrease in its functional activity. The ATPase activity is inhibited in the artificially induced sarcoplasmic membrane crystals (Herbert, et al., 1982). This could be due to inactivation by the vanadate and the Pi or a conformational change in the active sites of the enzyme in the crystalline array.

Alternatively, the close proximity of the globular head units (knobs) could result in ~~in~~ protein-protein interactions which allow for positive cooperativity in ATP hydrolysis thereby allowing for an increase in energy production (Munoz, 1982).

The apparent lability of the polar membrane structure when isolation was attempted and the increased frequency of observation of these structures when membrane preparations included a reducing agent, suggested that polar membrane may be sensitive to oxygen. This conclusion is plausible considering the fact that *C. jejuni* is a microaerophile.

4.2. Minicell formation by *C. jejuni*

The observation of small flagellated structures in late log and stationary phase cultures as well as whole cell preparations of *C. jejuni* led to a study of their nature and formation because they might have provided an enrichment of the polar membrane. It was found that these structures were minicells produced from the poles of the cell through a process involving constriction of the plasma membrane and segregation of the polar regions of the cells including the polar membrane structure. The profile of the peptidoglycan layer in the cell wall, expected to lie between the outer membrane and the cytoplasmic

membrane, was not often seen on thin section of young cells and could not be located on thin sections of the segregating terminal regions of the minicells. This may be due to the collapse of the periplasmic peptidoglycan gel against the outer membrane during dehydration as described by Hobot et al. (1984). It is assumed that the integrity of the peptidoglycan must be disturbed in the region of minicell formation, but remains intact along the main length of the cell because the rods did not spheroplast.

The experience of this laboratory with portions of cytoplasmic membrane bearing polar membrane has been that it becomes sharply curved with the concave side on the inside. This suggested that the turgor change likely in these cells during the initial event of minicell formation would allow this curvature and make a central contact and annealing of membranes the more likely. The prompt closure of the cell portions would then provide a cell-like environment and obviate the dissolution of the polar membrane structure that was usually experienced when attempts were made to fractionate it. The survival of the rod portions of the cells were not studied, but must be assumed to be more problematical.

The terminal events in the formation of these structures were not observed, but must be assumed to

involve a breakage of the outer membrane component of the cell wall allowing the release of these segregated polar regions, followed by fusion of the ruptured outer membrane. The formation of these minicells did not appear to occur by normal septation or constriction (Burdett and Murray, 1974; Adler et al., 1967). It is interesting to note that another polar flagellate, A. serpens, produces similar structures (Terasaki, 1970). This discard of the flagellated poles of the cell might be a survival feature used to rid the cell of its energy-requiring flagellar apparatus or it could be merely another degenerative feature related to coccoid body formation as described by Ogg (1962).

As stated earlier, one of the reasons for studying polar membrane in C. jejuni was the observation that this organism pinched off its poles producing minicells complete with cytoplasmic membrane and polar membrane. Attempts were made to use this phenomenon as a means to gain a membrane fraction enriched in polar membrane. This result was defeated by the sticky properties of both cells and minicells, which prevented a clean isolation of this membrane fraction.

4.3. Localization of ATPase Activity in whole cells by Histochemical staining

The localization of heavy internal lead deposits at the poles of the cells, after histochemical staining for ATPase activity, suggested that polar membrane is an assembly of ATPase rooted in an area of membrane. The presence of lighter lead deposits elsewhere in the cell may suggest the presence of other ATPases or simply the presence of the polar ATPase randomly distributed along the cytoplasmic membrane. Voelz (1964) showed, by histochemical staining (using the modified Wachstein-Meisel method of Otero-Vilardebo et al., 1963), the ATPase of E. coli to be randomly distributed on the internal surface of the cytoplasmic membrane, a result confirmed by this work. Vaituzis (1973) observed that Spirillum serpens (ATCC L1330), when histochemically stained in the presence of ATP and Mg^{2+} , did not reveal any activity preferentially associated with the poles of the cell. This organism appeared to have ATPase activity concentrated in the periplasmic space around the whole cell. The present study has shown that A. serpens MW5 shows random deposits of lead throughout the cytoplasm when incubated with ATP and Mg^{2+} and heavy lead deposition in the polar regions when incubated with ATP and Ca^{2+} . This result suggests that the polar ATPase of this

organism is activated by Ca^{2+} and not Mg^{2+} as in C. jejuni indicating that the function of polar membrane as an ATPase may cross generic boundaries but the biochemical properties of this structure may differ from organism to organism. Deposits in the periplasmic space were not observed. The discrepancy in the ATP- Mg^{2+} results may be due to the strain difference or to the prefixation with glutaraldehyde. Vaituzis (1973) used 6.5% glutaraldehyde prior to histochemical staining and this may have resulted in inactivation of internal ATPases in this organism.

Vaituzis (1973) also described the presence of lead deposition in the polar regions of Vibrio metchnikovii when incubated with ATP- Mg^{2+} . This organism, like other Vibrio sp. may possess polar membrane and as with C. jejuni this result may suggest a relationship between this structure and ATPase activity.

It has also been shown that the ATPase of Mycoplasma gallisepticum is localized at the poles of the cell (Munkres and Wachtel, 1967), however, polar membrane is not visible in thin sections of this organism.

The histochemical staining results found in B. cereus are similar to those found in B. licheniformis by Vaituzis (1973), suggesting a random distribution of ATPase activity within the cell. Polar membrane has not been described in this genus, but studies have been done

on the nature and location of its ATPase (Abram, 1965; Mirsky and Barlow, 1971)

M. luteus (formally M. lysodeikticus), a Gram positive coccus, appears to have either two different ATPases or one with different activities depending on its activation by Mg^{2+} or Ca^{2+} . ATPase activity was more localized when Mg^{2+} was used and randomly distributed when Ca^{2+} was the activator. The Gram negative coccus (but yet Gram positive in reaction) D. radiodurans showed the opposite results to M. luteus. Again, polar membrane has not been described in either of these genera, although a great deal of work has been focused on the ATPase of M. lysodeikticus (Chung and Salton, 1980; Ellar, et al., 1971; Huberman and Salton, 1979; Mollinedo, et al., 1983; Mollinedo, et al., 1980; Munoz, et al., 1968; Munoz, et al., 1969; Oppenheim and Salton, 1973; Salton, et al., 1968; Salton and Schor, 1972; Salton, et al., 1972; Urban and Salton, 1983a and b). This ATPase was found to be activated primarily by Ca^{2+} and to a lesser extent by Mg^{2+} and was found to be randomly distributed on the inner surface of the cytoplasmic membrane. It appears, therefore, that histochemical staining of M. luteus showed the presence of this ATPase and its differential activation by Mg^{2+} and Ca^{2+} .

4.4. Isolation and characterization of an F₁-like ATPase from *C. jejuni*.

The results obtained from analysis of the initial shock-wash supernatants of *C. jejuni* membranes for ATPase activity revealed the presence of three protein species (one major species and two minor ones) capable of degrading ATP and releasing Pi. The major protein species was chosen for study in that it specifically degraded ATP and was activated only by Mg²⁺ cations similar to the ATPase activity seen in histochemically stained cells. The other active protein species appeared to be general phosphatases in that they degraded a variety of nucleotide triphosphates and glycerol-6-phosphate. They were also primarily activated by Ca²⁺ thereby suggesting no relationship to the ATPase activity seen at the poles of the cell.

The major protein species possessing ATPase activity was purified by the method of Vogel and Steinhart (1976) and appeared to be homogenous as judged by polyacrylamide gel electrophoresis, even though the enzyme purification was only 4 fold. This relatively low purification factor may have resulted from the use of a purification method designed for extraction of the enzyme from *E. coli*, although, extraction with n-butanol, as done with the enzyme from *M. lysodeikticus* (Salton and Schor, 1974),

gave the same results. It has been noted that a great deal of variation exists in the procedures for ATPase release and purification from mitochondria, chloroplasts and bacterial membranes and a method that is successful for one organism may not work in another (Ishikawa, and Lehninger, 1962; Munoz, et al., 1968 and 1969; Knowles and Penefsky, 1972; McCarty and Racker, 1966; Strotmann, et al., 1973; Abrams, 1965; Davies and Bragg, 1972; Hanson and Kennedy, 1973; Futai, et al., 1974; Nelson, et al., 1974; Mirsky and Bariow, 1971; Ishida and Mizushima, 1969; Hachimori, et al., 1970; Adolfsen and Moudrianakis, 1971; Higashi, et al., 1975; Riebeling and Jungermann, 1976; Yoshida et al., 1975; Reed and Raveed, 1972; Johansson, et al., 1973). It is generally assumed that solubilization of F_1 -ATPases requires the removal of divalent cations, as these enzymes are associated with the membrane by a combination of ionic forces and hydrophobic interactions. The relative importance of these forces and the degree to which they act may explain the differences seen in solubilization regimes. In the case of the enzyme in *C. jejuni*, equal purification was obtained whether EDTA or an organic solvent was used, which suggested that some other factor or factors were involved in the low purification factor obtained. It is possible, considering the microaerophilic nature of *C. jejuni* and the fact that a

reducing agent added stability to the complex, that the ATPase was oxygen sensitive and this could account for the low purification factor and specific activity observed, as no precautions were taken to remove oxygen from the extraction and purification buffers. The possibility also exists that due to the crystalline nature of polar membrane and possibly this ATPase, the isolated enzyme may be rendered partially inactive after removal from its closely packed conditions on the membrane or it may be partially inactive and awaiting some final activation step. However, it should be noted that, despite the solubilization procedures used, there also exists a great deal of variation in the level of ATP hydrolytic activity (i.e. specific activity) of purified F₁-ATPases from different sources, the specific activity of the enzyme isolated from *M. lysodeikticus* is very low (6-8 $\mu\text{mol/min/mg}$ of protein) even though this enzyme exhibits latency (Ishikawa and Lenhninger, 1962; Munoz, et al., 1968; Munoz, et al., 1969). The specific activities of the bacterial F₁-ATPases also differ in that the enzymes from anaerobes and facultative anaerobes possess mid-range activities in the hydrolysis of ATP, those from aerobes (particularly those of *Bacillus* spp.) have very high specific activities (500-1000 $\mu\text{mol/min/mg}$ of protein), and

those from the strict aerobes (*M. lysodeikticus*, *A. faecalis*, *M. phlei*) have very low specific activities (3-8 $\mu\text{mol}/\text{min}/\text{mg}$ of protein) (Abrams and Smith, 1974; Abrams, et al., 1976; Futai, et al., 1974; Nelson, et al., 1974; Serrahina-Zieger and Monteil, 1978; Monteil and Serrahina-Zieger, 1978; Munoz, et al., 1969; Adolfsen and Moudrianakis, 1976; Ritz-Gold and Brodie, 1979; Ritz-Gold, et al., 1979; Bogin, et al., 1970; Ritz and Brodie, 1977).

All bacterial F_1 -ATPases share the property of appearing as spherical structures on the membrane and in purified form (Munoz, 1982). The stalk portion of the enzyme is not frequently observed (Munoz, 1982). The purified ATPase from *C. jejuni* also possesses this characteristic, although its dimensions (5-6 nm) are slightly smaller than the 9 nm described for many other bacterial F_1 -ATPases.

The ATPase of *C. jejuni* could be reconstituted on enzyme depleted membranes restoring ATP hydrolytic activity. This result suggested that the enzyme was not altered or damaged during the isolation and purification procedures used. It does not, however, ensure that the entire F_1 portion of the enzyme was removed from the membranes and it is possible that parts of the F_1 structure (eg. other subunits) were still associated with

the membrane after extraction.

The purified active part of the ATPase of C. jejuni appeared to be composed of one subunit, which is a situation observed in Lactobacillus casei (Biketov, et al., 1982; Tikhonova, et al., 1983). Other subunits might be lost during isolation and purification or may be lost due to oxygen lability. It is also possible as stated earlier that not all of the subunits were extracted from the membrane, but the methods were effective at extracting multiple subunits from E. coli K12.

It has been generally assumed that F₁-ATPases are composed of five subunit types (Munoz, 1982), although some variability has been observed. The delta subunit of E. coli F₁-ATPase disappears during certain purification procedures (Bragg, et al., 1973; Azocar and Munoz, 1977). The subunit composition of the F₁-ATPase of M. lysodeikticus shows great variation depending on the purification method, analytical techniques used, enzyme stability and the use of different substrains (Munoz, et al., 1969; Mollinedo, et al., 1980; Andreu, et al., 1973; Risi, et al., 1977; Carreira, et al., 1976 and 1977; Andreu and Munoz, 1975). In Strep. faecalis F₁-ATPase three subunits were seen using urea gel electrophoresis and four subunits were noted using SDS-gel electrophoresis (Schnebli, et al., 1970), a fifth subunit was located when

the enzyme was purified in the presence of Mg^{2+} (Abrams, *et al.*, 1976). A similar situation was found in the analysis of *M. lysodeikticus* F_1 -ATPase, the subunit composition was less complex under urea gel analysis than SDS-gel electrophoresis and variations in the staining of this protein with urea were observed (Schnebli, *et al.*, 1970). The F_1 -ATPases from *B. megaterium* and *B. subtilis* have a simpler subunit composition, only the two major subunits (alpha and beta) have been consistently isolated (Mirsky and Barlow, 1971; Ishida and Mizushima, 1969; Monteil and Serrahima-Zieger, 1978), and as mentioned earlier the F_1 -ATPase isolated from *Lactobacillus casei* possesses only one subunit (Biketov, *et al.*, 1982; Tikhonova, *et al.*, 1983).

Therefore, considering the variation which exists in the subunit composition of a variety of bacterial F_1 -ATPases it is plausible that the F_1 -ATPase isolated from *C. jejuni* possesses a single subunit type, although further work in this area is needed to state this conclusively.

The effects of the inhibitors DCCD and oligomycin are in agreement with those found for other procaryotic F_1 -ATPases (Haddock and Jones, 1977; Harold, 1972) and further suggested this enzyme belonged in the procaryotic H^+ -ATPase class. The inability of trypsin to activate this

enzyme suggested that it was not in a latent form as was the case for the enzyme from *M. lysodeikticus* (Ishikawa and Lehninger, 1962; Munoz, *et al.*, 1968 and 1969)

The enzyme isolated from *C. jejuni* was specifically activated by Mg^{2+} cations, a characteristic also pointing to its classification as a F_1 -ATPase, however, its inhibition by Ca^{2+} is unusual in that most Mg^{2+} dependent ATPases can be activated in a minor way by Ca^{2+} (Downie, *et al.*, 1979). The activation of F_1 -ATPases by cations is a another property which shows clear differences depending on the origin of the enzyme. F_1 -ATPases of animal origin (i.e. from mitochondria) require Mg^{2+} as an activator for the degradation of ATP giving specific activities of 100 μ mol ATP hydrolysed/min/mg of protein (Penerfsky, 1974 and 1979; Pedersen, 1975; Pedersen, *et al.*, 1978). The F_1 -ATPases of chloroplasts are activated by Ca^{2+} , however their specific activities are very low unless activated by heat treatment, trypsin digestion or dithiothreitol (Nelson, 1976; McCarty, 1979; Baird and Hammes, 1979; Vambutas and Racker, 1965; Matsubara, *et al.*, 1981). Alternately, bacterial F_1 -ATPases show a more complex pattern of activators and specific activities. Generally the activators of the soluble enzymes are Mg^{2+} and Ca^{2+} , although other metal cations have been found to be activators, and it appears that enzymes from anaerobes and

facultative anaerobes are Mg^{2+} -dependent ATPases while those from aerobes are Ca^{2+} -dependent ATPases (Munoz, 1982). However, these enzymes all appear to behave as Mg^{2+} ATPases when bound to membranes (Munoz, 1982). The observations reported here on the enzyme from *C. jejuni* show clear differences to other F_1 -ATPases and these may stem from the microaerophilic nature of this organism.

The F_1 -ATPase from *C. jejuni* specifically degraded ATP, but at a very low rate. There might be hydrolytic activity against other substrates but such an activity was not detectable by the methods used in this study. It is known that bacterial F_1 -ATPases generally possess hydrolytic activity for nucleotide substrates other than ATP, usually with lower specific activities (Mirsky and Barlow, 1971; Davies and Bragg, 1972; Hanson and Kennedy, 1973; Abrams and Smith, 1974).

The ATPase from *C. jejuni* also exhibited cold inactivation, a property common to F_1 -ATPases from mitochondria, chloroplasts and other bacteria (Penevsky and Warner, 1965; Bennum and Racker, 1969; Livne and Racker, 1969; Monteil and Serrahima-Zieger, 1978). The F_1 -ATPase of *B. megaterium* is very cold labile (Mirsky and Barlow, 1972) and the cold sensitivity of *E. coli* ATPase depends on the conditions of enzyme extraction and on the strain (Azocar and Munoz, 1977; Hanson and Kennedy, 1973; Davies and Bragg, 1972; Futai, et al., 1974).

4.5. Immunocytochemical localization of ATPase in C.
jejuni

The immunocytochemical localization of this ATPase by immunoelectron microscopy suggested that it was associated with the polar membrane structure because the colloidal gold label was clustered at the poles of the cell. The location of the enzyme (using colloidal gold-antibody detection methods) elsewhere in the cell suggested the presence of the ATPase generally distributed at sites other than the pole. It should be noted that often the polar membrane structure was visible on thin sections but the gold label was not seen bound to the structure. This would be expected due to the inherent problems experienced in the post-embedding labeling technique because of the orientation in the resin embedding. When the structure is visible it is cut across by the plane of the section (i.e. in the resin) and few if any of the target molecules are available as an antibody binding site (Roth, 1982). One would hope for a tangential section to the polar membrane with revelation of a number of antibody binding sites.

The use of Spurr's resin for this technique and osmium fixation of the cells gave acceptable results. However, etching of the sections increased the gold labeling and enhanced the localization of the enzyme. This

is an indication that the enzyme's antigenicity was not greatly affected by the fixation and embedding procedures. The use of Lowicryl KM4 for embedding *C. jejuni* provided a better method for antigen preservation because samples were not subjected to high temperatures for polymerization of the resin. It was found that larger clusters of gold particles at the poles of the cells were seen on Lowicryl KM4 sections and again the best result was seen in sections tangential to the polar regions. However, the cell wall and membrane contrast was very poor and polar membrane could not be seen.

The gold labelling pattern was consistent with the lead deposition pattern seen in histochemically stained cells. These results and the structural similarities suggest that polar membrane and the purified Mg^{2+} -dependent ATPase from *C. jejuni* are directly related.

The location and concentration of this ATPase at the poles of the cells, near the flagellar insertion point, suggests it may function in energy transduction or proton translocation for flagellar rotation. The production of minicells may indeed be a mechanism useful to the cell for ridding itself of this concentrated ATP hydrolysing region.

4.6. Antigenic relatedness of the ATPase from *C. jejuni* to ATPases of other bacterial species and rat liver mitochondria

The antibody prepared against the ATPase from *C. jejuni* recognized four protein species in whole cell extracts of this organism. This result may suggest the presence of other subunits of this ATPase which were not extractable using the methods of Vogel and Steinhart (1976) or of Salton and Schor (1974) but, more likely, may simply be cross reaction with other cell proteins sharing some specificity.

The cross reactivity of this antibody with protein species from whole cell extracts of *E. coli* K12 and *A. serpens* MW5 suggested some conservation of sequences in these proteins, but the relationship to ATPase is uncertain. There are broader differences because *A. serpens* MW5 also possesses polar membrane, has an enzyme that appears to be activated by Ca^{2+} and inhibited by Mg^{2+} . The other three bacterial species tested also possess polar membrane, but the antibody to the *C. jejuni* ATPase did not cross react with any cell proteins in these organism, nor did it cross react with mitochondrial proteins. These results suggest that the composition of polar membrane may differ depending on its origin, although its function may be similar, and this ATPase may

have evolved along different lines than those for other bacterial species and mitochondria.

The antibody prepared against the F₁-ATPase of E. coli K12 showed cross reactivity with A. serpens MW5, V. cholera, S. volutans and rat liver mitochondria. It did not cross react with any cell proteins from either C. jejuni or Methanococcus voltae. As well, a monoclonal antibody to the beta subunit of E. coli K12 did not cross react with C. jejuni (S. Dunn, Dept. Biochem., Univ. Western Ont., personal communication). It has been shown that monoclonal antibodies prepared against the beta subunit of the E. coli ATPase show cross reactivity with a variety of bacterial genera, and beta subunits of mitochondrial and chloroplast F₁ ATPases (Dunn, 1985). This suggests that this subunit has been highly conserved throughout evolution. However, it appears that the ATPases of C. jejuni and the phylogenetically separate archaeobacterium M. voltae (Woese, 1987) may have evolved along different pathways. The cross reactivity of the C. jejuni antibody with E. coli K12 may represent a domain recognized in an otherwise unique protein.

4.7. Conclusions

This study examined the ultrastructure and nature of polar membrane in C. jejuni and showed that the structure

was composed of dough-nut shaped particles (5-6 nm in diameter) arranged on short stalks in an hexagonal array on the membrane. The cytoplasmic membrane, when depleted of its lipids, possessed areas with a fine hexagonal array (3 nm repeat) and this may represent the stalks themselves or the insertion sites of the stalks of the polar membrane structure into the membrane. Polar membrane was found to be a tightly packed array suggesting it to be a naturally occurring two dimensional crystal similar to the vanadium induced ATPase crystals seen on sarcoplasmic membranes (Dux and Martonosi, 1983a).

It was also found that visualization of this structure on the membrane was enhanced under conditions of high ionic strength and in the presence of a reducing agent, suggesting that polar membrane was sensitive to oxidation. This result emphasized the need for increased study into the oxygen sensitivity of proteins isolated from *C. jejuni*.

This study also described the presence and formation of minicells, possessing polar membrane, from the poles of whole cells of *C. jejuni*. This process was found to occur by means other than the normal septation described by Burdett and Murray, (1974) and Adler, et al. (1967) and was thought to be enhanced by the rigidity of the cytoplasmic membrane in the region of the polar membrane

such that the tugor change likely in formation of these structures would allow an inward curvature of the membranes making the central contact and annealing more likely.

The nature of polar membrane was found, by histochemical staining, to be expressed as Mg^{2+} -dependent ATPase activity in appropriate locations, i.e., at the poles of the cells and in localized patches along the inner surface of the cytoplasmic membrane in spheroplasts of this organism. A study of ATPase activity in A. serpens MW5, by histochemical staining, showed a Ca^{2+} -dependent ATPase activity associated with the polar membrane regions. This suggested that while polar membrane is structurally similar in many lophotrichous Gram negative bacteria its biochemical properties may differ.

The isolation of a Mg^{2+} -dependent ATPase from C. jejuni was accomplished and, it had the biochemical characteristics of an F_1 -ATPase. Further work is required in this area. Antibodies prepared against this enzyme were shown to localize (by immunoelctron microscopy) in the polar regions of the cells near to and sometimes on the polar membrane structure. This study suggested that polar membrane was an array of a Mg^{2+} -dependent ATPase. It was hypothesised that minicell formation was

facilitated by the presence of the polar membrane structure and provided a method by which the cell could, by fortunate accident, rid itself of an energy requiring apparatus.

The study of the antigenic relationship of this ATPase to those from other bacterial genera and rat liver mitochondria showed, again, that biochemical differences may exist between the components of polar membrane in different organisms and that the ATPase of *C. jejuni* and *M. voltae* appear to have evolved along different pathways to those of *E. coli* and mitochondria.

The presence of polar membrane in a variety of bacterial genera, as well as the seeming conservation of different antigenic specificities, pose some interesting phylogenetic questions. It is possible that a more detailed study of this structure in other organisms may lead to a greater understanding of the phylogeny of the organisms possessing polar membrane. A study of the ATPase activity of polar membrane in different organisms may also provide valuable insight into the conservation of the active units of this metabolically important enzyme and begin a more productive look at the differences of F₁-ATPases rather than the persistent search for homologies within this enzyme group.

8

Figure 1: Diagram of the polar membrane in
Ectothiorhodospira mobilis. (From
Remsen, etal., 1968).

Figure 2: Conceptual model of FoF1 of Escherichia coli.

This model was drawn based on reconstitution experiments, subunit stoichiometry, predicted secondary structures, and subunit interactions. This model is not drawn to scale. Lettering of subunits is described in the text.

027

Figure 3: Growth curve for C. jejuni grown in 500 ml

- volumes of Brucella broth (Difco) without shaking at 40o C.

Figure 4: Thin section of C. jejuni, fixed with 5% acrolein and 0.25% glutaraldehyde in 0.05 M cacodylate buffer (pH 7.5) and stained with 1% Uranyl acetate (aqueous) and Reynolds lead citrate, showing the cytoplasmic membrane (CM), polar membrane (PM), outer membrane (OM), polar flagellum (f), ribosome free area (RF), and spherical inclusion bodies (SI). Bar = 100 nm.

Figure 5: Thin section of A. serpens MW5 fixed with 5% acrolein and 0.25% glutaraldehyde in 0.05 M cacodylate buffer (pH 7.5) and stained with 1% uranyl acetate (aqueous) and Reynolds lead citrate, showing cytoplasmic membrane (CM), polar membrane (PM), outer membrane (OM), peptidoglycan (PG), and ribosome free area (RF). Bar = 100 nm.

Figure 6: Cross section of C. jejuni fixed and stained as in Fig. 4, showing polar membrane (PM) lining the inner surface of the cytoplasmic membrane (CM). , OM, outer membrane. Bar = 100 nm.

Figure 7: Cross section of A. serpens MWS fixed and stained as in Fig. 5, showing patches of polar membrane (PM) on the inner surface of the plasma membrane. OM, outer membrane; PG, peptidoglycan; RS, regular structured array. Bar = 100 nm.

63

Figure 8: Thin section of C. jejuni fixed and stained as in Fig. 4, showing septum (S) between two daughter cells and polar membrane (PM) lining the cytoplasmic membrane (CM). Bar = 100 nm.

2

Figure 9: Thin section of a dividing C. jejuni cell, showing a normal septum (S) and no visible polar membrane on the cytoplasmic membrane at this site. Polar membrane (PM) is visible at the opposing end of one daughter cell. Bar = 100 nm.

Figure 10: Thin section of a polar region of C. jejuni, showing association of polar membrane (PM) with the flagellar basal complex and the M ring (M), and fibrous material (FM) of the basal complex. OM, outer membrane; CM, cytoplasmic membrane. Bar = 100 nm.

Figure 11: Thin section of the polar region of C. jejuni showing the association of polar membrane (PM) to the flagellar basal complex and the concentric membrane rings (CMR), P ring (P), and the M ring (M) of the basal complex. F, flagellum. Bar = 100 nm.

83?

Figure 12: Thin section of the polar region of C. jejuni showing association of polar membrane (PM) to the flagellar basal complex and the M ring (M) of this complex. RF, ribosome free area; SI, spherical inclusion body; F, flagellum. Bar = 100 nm.

027

Figure 13: Thin section of the polar region of C. jejuni showing the association of polar membrane (PM) to the flagellar basal complex and the M ring (M), concentric membrane rings (CMR), and the P ring (P). Bar = 100 nm.

A thin section of the polar region of C. jejuni, showing the association of polar membrane (PM) to the flagellar basal complex (FB). The image displays a cross-section of the bacterial cell, with the polar membrane (PM) and flagellar basal complex (FB) visible. The ribosome free area (Rf) and spherical inclusion body (SI) are also present.

Figure 14: Thin section of the polar region of C. jejuni showing the association of polar membrane (PM) to the flagellar basal complex (FB). Rf, ribosome free area; SI, spherical inclusion body. Bar = 100 nm.

Figure 15: Thin section of the polar region of C. jejuni showing the association of polar membrane (PM) to the flagellar basal complex. CMR, concentric membrane rings; RF, ribosome free area. Bar = 100 nm.

Figure 16: Longitudinal and cross section of C. jejuni showing polar membrane (PM) and the rigid cytoplasmic membrane (RCM) associated with the structure. Bar = 100 nm.

Figure 17: French pressure cell disrupted membrane preparation of C. jejuni negatively stained with 1% phosphotungstic acid (pH 6.8). 17a, shows dough-nut shaped particles (arrow) associated with a membrane fragment from the polar region as denoted by the presence of a flagellum (f). 17b, shows an intact polar vesicle with dough-nut shaped particles (arrow) on the inner membrane. Bars = 100 nm.

Figure 18: French pressure cell disrupted membrane preparation of C. jejuni negatively stained with 1% phosphotungstic acid (pH 6.8) showing an enclosed membrane vesicle with dough-nut shaped particles (arrow) associated with the internal membrane of the vesicle; the outer enclosing membrane contains a concentric membrane ring (CMR) denoting it as outer membrane from the polar region of the cell. Bar = 100 nm.

Figure 19: Negatively stained membrane preparations of C. jejuni. 19a, membranes prepared in a high ionic strength buffer plus 2.5 mM 2-mercaptoethanol showing hexagonally arranged dough-nut shaped particles (arrow). 19b, membranes prepared by the modified monolayer technique of Nermut (1983) showing hexagonally arranged dough-nut shaped particles (arrow). 19c, and 19d, membranes delipidated with 1% cholic acid, showing fine hexagonal arrays (arrows) in flattened membrane sheets. Bars = 100 nm.

Figure 20: Thin section of membrane preparation from *C. jejuni* pre-fixed with 1% tannic acid and fixed with 5% acrolein and 0.25% glutaraldehyde showing poorly resolved polar membrane (PM) associated with the cytoplasmic membrane (CM). OM, outer membrane. Bar = 100 nm.

Figure 21: Negatively stained minicells from C. jejuni bearing flagella (arrow). Bar = 100 nm.

Figure 22: Thin sections of minicells from C. jejuni.

OM, outer membrane; CM, cytoplasmic membrane;

PM, polar membrane. Bar = 100 nm.

Figure 23: Thin sections of C. jejuni showing formation of minicells. 23 (1) to (4) shows increasing constriction of the cytoplasmic membrane resulting in segregation of the polar regions of the cells. Bar = 100 nm.

Figure 24: Negatively stained whole cell of C. jejuni, showing a minicell forming in the polar region (arrow). Bar = 100 nm.

491

Figure 25> Thin section of minicell preparation showing contamination with whole cells, membrane vesicles and flagellar debris. Bar = 100 nm.

Figure 26: Negative stain of a minicell from C. jejuni showing an hexagonal pattern (arrow) in the interior. Bar = 100 nm.

Figure 27) a to d, thin sections of *C. jejuni* fixed with 1% glutaraldehyde and histochemically stained for the presence of Mg^{2+} -ATPase activity. Sections were counterstained with saturated uranyl acetate in 50% (v/v) methanol. Arrows show areas of heavy lead phosphate deposition.

Figure 28: a and b, thin sections of spheroplasts of C. jejuni fixed with 1% glutaraldehyde and histochemically stained for Mg^{2+} -ATPase activity. Arrows show localized lead phosphate deposition.

Figure 29: a and b, thin sections of C. jejuni fixed with 1% glutaraldehyde and histochemically stained for Ca^{2+} -ATPase activity. Arrows show lead phosphate deposition.

3 of/de 3

MICRO

Figure 30: a and b, thin sections of E. coli K12 fixed with 1% glutaraldehyde and histochemically stained for Ca^{2+} -ATPase activity. 30c, thin section of E. coli K12 histochemically stained for Mg^{2+} -ATPase activity. Bar = 100 nm.

Figure 31: Thin section of B. cereus fixed with 1% glutaraldehyde and histochemically stained for Mg^{2+} -ATPase activity. Bar = 200 nm.

Figure 32: Thin sections of M. luteus fixed with 1% glutaraldehyde and histochemically stained for (a), Ca^{2+} -ATPase activity and (b), Mg^{2+} -ATPase activity. Bars = 200 nm.

B

Figure 33: Thin sections of D. radiodurans fixed with 1% glutaraldehyde and histochemically stained for (a and b), Mg^{2+} -ATPase activity, and (c) Ca^{2+} -ATPase activity. Bars = 200 nm.

Figure 34: Thin sections of A. serpens MW5 fixed with 1% glutaraldehyde and histochemically stained for (a), Mg^{2+} -ATPase activity, and (b) Ca_{2+} -ATPase activity. Bars = 200 nm.

Figure 35: Non-dissociating gel analysis of protein profile and ATPase activity in polyethylene glycol precipitated material from the initial shock-wash supernatant from membranes of *C. jejuni*. Lane a and e, protein profile stained with Coomassie brilliant blue. Lane b, incubated with 5 mM ATP. Lane c, incubated with 5 mM ATP plus 10 mM $MgCl_2$. Lane d, incubated with 5 mM ATP plus 10 mM $CaCl_2$. Lane f, incubated with 5 mM GTP plus 10 mM $MgCl_2$. Lane g, incubated with 5 mM GTP plus 10 mM $CaCl_2$. Lanes b to d, and f, and g were stained for the presence of inorganic phosphate by the method of Abrams, et al., (1976).

Figure 36: Non-dissociating-PAGE of purified ATPase (a) stained with Coomassie brilliant blue (arrow shows dye front) and (b) incubated with 5 mM ATP and 10 mM $MgCl_2$ and stained for the presence of Pi by the method of Abrams, et al., (1976).

Figure 37: Negatively stained solubilized ATPase from C.
jejuni. Bar = 100 nm.

024

Figure 38: SDS-PAGE of RNA polymerase (lane a) and C. jejuni ATPase (lane, b). Urea-PAGE of C. jejuni ATPase (lane c). Gels were stained for the presence of protein with Coomassie brilliant blue.

30k

30k

30k

30k

a

b

c

Figure 39: Non-dissociating PAGE analysis of the effect of cations on *C. jejuni* ATPase activity as revealed by release of Pi from ATP according to the method of Abrams, et al., (1976). Lane a, stained for protein with Coomassie brilliant blue. Lane b, gel slice was incubated with 5 mM ATP and 10 mM MgCl₂. Lane c, gel slice was incubated with 5 mM ATP and 10 mM CaCl₂. Lane d, gel slice was incubated with 5 mM ATP alone (no cations).

Figure 40: Non-dissociating-PAGE analysis of the substrate specificity of the ATPase of C. jejuni. Lane a, stained for protein with Coomassie brilliant blue. The other gel slices were incubated with the following substrates plus 10 mM MgCl₂: Lane b, 5 mM ATP; Lane c, 5 mM ADP; Lane d, 5 mM GTP; Lane e, 5 mM glycerol-6-phosphate. ATPase activity was shown by staining for Pi according to the method of Abrams, et al., (1976).

Figure 41: a to d, immunocytochemical localization of C.
jejuni ATPase on thin sections of Spurr
embedded whole cells. Arrow shows gold
particles bound to a minicell (Fig. 41b). Gold
particles = 10 to 15 nm.

Figure 42: a and b, immunocytochemical localization of C.
jejuni ATPase on thin sections of Spurr
embedded whole cells. Bar = 100 nm.

Figure 43: a and b, immunocytochemical localization of C.
jejuni ATPase on hydrogen peroxide etched
sections of Spurr embedded whole cells. Gold
particles = 10 to 15 nm.

Figure 44: a and b, immunocytochemical localization of C.
jejuni ATPase on thin sections of Lowicryl KM4
embedded whole cells. Arrows show clusters of
gold particles. Gold particles = 5 nm.

Figure 45: Immunocytochemical localization of C. jejuni ATPase on thin sections of Lowicryl KM4 embedded whole cells. Arrow shows large cluster of gold particles at the pole of a cell. Gold particles = 5 nm.

Figure 46: Western blots of whole cell proteins reacted with antibody prepared against the ATPase isolated from C. jejuni. Lane a, molecular weight markers. Lane b, E. coli K12. Lane c, V. cholera. Lane d, A. serpens MW5. Lane e, S. volutans. Lane f, C. jejuni. Lane g, M. voltae. Lane h, rat liver mitochondria. Arrows on lanes b, d, and f, mark the protein species which bound antibodies.

Figure 47: Western blots of whole cell proteins reacted with antibody prepared against the F1-ATPase from E. coli K12. Lane a, molecular weight markers. Lane b, E. coli K12. Lane c, V. cholera. Lane d, A. serpens MW5. Lane e, S. volutans. Lane f, C. jejuni. Lane g, M. voltae. Lane h, rat liver mitochondria. Arrows in lanes b, c, d, e, and h, mark protein species which bound antibodies.

References

Abram, D. 1965. Electromicroscope observation on intact cells, protoplasts, and cytoplasmic membrane of Bacillus stearothermophilus. J. Bacteriol. 89: 855-873.

Abrams, A. 1965. The release of bound adenosine triphosphatase from isolated bacterial membranes and the properties of the solubilised enzyme. J. Biol. Chem. 240: 3675-3681.

Abrams, A. 1976. Bacterial ATPases. In The Enzymes of Biological Membranes (Martosi, A., ed.), vol 3, pp. 57-73. Plenum Press, N.Y.

Abrams, A., and Baron, C. 1970. Inhibitory action of carbodiimides on bacterial membrane ATPase. Biochem. Biophys. Res. Commun. 41: 858-863.

Abrams, A. and Smith, J.B. 1974. Bacterial Membrane ATPase. in The Enzymes (Boyer, P.D., ed.), vol. 10, pp. 395-429, Academic Press, N.Y.

Abrams, A., McNamara, P., and Johnson, F.B. 1960. Adenosine triphosphatase in isolated bacterial cell membranes. J. Biol Chem. 235: 3659.

Abrams, A., Smith, J.B., and Baron, C. 1972.

Carbodiimide-resistant membrane adenosine triphosphatase in mutants of Streptococcus faecalis. I. Studies on the mechanism of resistance. J. Biol. Chem 247: 1484-1488.

Abrams, A., Morris, D., and Jensen, C. 1976. Chymotryptic conversion of bacterial membrane ATPase to an active form with modified alpha chains and defective membrane binding properties. Biochem. 15: 5560-5565.

Ackrell, B.A.C., and Jones, C.W. 1971. The respiratory system of Azotobacter vinelandii. I. Properties of phosphorylating respiratory membranes. Eur. J. Biochem. 20: 22-28.

Addison, R., and Scarborough, G.A. 1981. Solubilization and purification of the Neurospora plasma membrane H⁺-ATPase. J. Biol. Chem. 256: 13165-13171.

Adler, H.I., Fisher, W.D., Cohen, A., and Hardigee, A.A. 1967. Miniature Escherichia coli cells deficient in DNA. Proc. Natl. Acad. Sci. U.S.A. 57: 321-326.

Adolfsen, R., McClung, J.A., and Moudrianakis, E.N. 1975. Electrophoretic microheterogeneity and subunit composition of the 13S coupling factors of oxidative photosynthetic phosphorylation. Biochem. 14: 1727-1735.

Adolfson, R., and Moudrianakis, E.N. 1971. Purification and properties of two soluble coupling factors of oxidative phosphorylation from Alcaligenes faecalis. Biochem. 10: 2247-2253.

Adolfson, R., and Moudrianakis, E.N. 1976. Molecular polymorphism and mechanisms of activation and deactivation of the hydrolytic functions of the coupling factor of oxidative phosphorylation. Biochem. 15: 4163-4170.

Amzel, L.M., and Pedersen, P.L. 1983. Proton ATPases: Structure and mechanism. Annu. Rev. Biochem. 52: 801-824

Andreu, J.M., Albendea, J.A., and Munoz, E. 1973. Membrane adenosine triphosphatase of Micrococcus lysodeikticus; molecular properties of the purified enzyme unstimulated by trypsin. Eur. J. Biochem. 37: 505-515.

Andreu, J.M., and Munoz, E. 1975. Micrococcus lysodeikticus ATPase; purification by preparative gel electrophoresis and subunit structure studied by urea and sodium dodecyl-sulphate gel electrophoresis. Biochim. Biophys. Acta 387: 228-233.

Archer, D.B., Rodwell, A.W., and Rodwell, E.S. 1978. The nature and location of Acholeplasma laidlawii membrane proteins investigated by two-dimensional gel electrophoresis. *Biochim. Biophys. Acta* 513: 268-283.

Asano, A., Cohen, N.S., Baker, R.F., and Brodie, A.F. 1973. Orientation of the cell membrane in ghosts and electron transport particles of Mycobacterium phlei. *J. Biol. Chem.* 248: 3386-3397.

Azocar, O., and Munoz, E. 1977. Extrinsic and intrinsic factors that influence inactivation and purification of the unstable adenosine triphosphatase solubilized from membranes of an escherichia coli K12 strain. *Biochim. Biophys. Acta* 482: 438-452.

Baccarini-Melandri, A., Gest, H., and San Pretro, A. 1970. A coupling factor in bacterial photophosphorylation. *J. Biol. Chem.* 245: 1224-1226.

Baird, B.A., Hammes, G.G. 1979. Structure of oxidative- and photo-phosphorylation coupling factor complexes. *Biochim. Biophys. Acta* 549: 31-54.

Bayer, M.E. 1968. Areas of adhesion between wall and membrane of Escherichia coli. J. Gen. Microbiol. 53: 395-404.

Balch, W.E., and Wolfe, R.S. 1979. Specificity and biological distribution of Co enzyme M (2-mercaptoethanesulfonic acid). J. Bacteriol. 137: 256-263.

Bell, R.M., Mavis, R.D., Osborn, M.J., and Vagelos, P.R. 1971. Enzymes of phospholipid metabolism: Localization in the cytoplasmic and outer membrane of the cell envelope of Escherichia coli and Salmonella typhimurium. Biochim. Biophys. Acta 249: 628-635.

Bengis-Garber, C., and Gromet-Elhanan, Z. 1979. Purification of the energy-transducing adenosine triphosphatase complex from Rhodospirillum rubrum. Biochem. 18: 3577-3581.

Bennum, A., and Racker, E. 1969. Partial resolution of the enzymes catalyzing photophosphorylation. IV. Interaction of coupling factor 1 from chloroplasts with components of the chloroplast membrane. J. Biol. Chem. 244: 1325-1331.

Berg, H.C. 1974. Dynamic properties of bacterial flagellar motors. *Nature* (London) 249: 77-79.

Berg, H.C. 1975. Bacterial behaviour. *Nature* (London) 254: 389-392.

Beveridge, T.J. 1981. Ultrastructure, chemistry, and function of the bacterial wall. *Int. Rev. Cytol.* 72: 229-317.

Beveridge, T.J., and Davies, J.A. 1983. Cellular responses of Bacillus subtilis and Escherichia coli to the Gram stain. *J. Bacteriol.* 156: 846-858.

Biketov, S.F., Kasho, V.N., Kozlov, I.A., Mileykovskaya, Y.I., Ostrovsky, D.N., Skulachev, V.P., Tikhonova, G.V., and Tsuprun, V.L. 1982. F₁-like ATPase from anaerobic bacterium Lactobacillus casei contains six similar subunits. *Eur. J. Biochem.* 129: 241-250.

Birch-Andersen, A., Maaloe, O., and Sjostrand. 1953. High resolution electron micrographs of sections of Escherichia coli. *Biochim. Biophys. Acta* 12: 395-400.

Bogin, E., Higashi, T., and Brodie, A.F. 1970. Oxidative phosphorylation in fractionated bacterial systems XLIII. Coupling factors associated with the NAD^+ -linked electron transport pathway. Arch. Biochem. Biophys. 136: 337-351.

Bowman, E.J. 1983. Comparison of the vacuolar membrane ATPase of Neurospora crassa with the mitochondrial and plasma membrane ATPase. J. Biol. Chem. 258: 15238-15244.

Bowman, E.J., Bowman, B.J., and Slayman, C.W. 1981. Isolation and characterization of plasma membranes from wild type Neurospora crassa. J. Biol. Chem. 256: 12336-12342.

Bowman, B.J., Mainzer, S.E., Allen, K.E., and Slayman, C.W. 1978. Effects of inhibitors on the plasma membrane and mitochondrial adenosine triphosphatase of Neurospora crassa. Biochim. Biophys. Acta 512: 13-28.

Bragg, P.D., Davies, P.L., and Hou, C. 1973. Effect of removal or modification of subunit polypeptides on the coupling factor and hydrolytic activities of the Ca^{2+} and Mg^{2+} -activated adenosine triphosphatase of Escherichia coli. Arch. Biochem. Biophys. 159: 664-670.

Bragg, P.D., and Hou, C. 1972. Organization of proteins in the native and reformed outer membrane of Escherichia coli. Biochim. Biophys. Acta 274: 478-488

Bragg, P.D., and Hou, C. 1975. Subunit composition function and spacial arrangement in the Ca^{2+} - and Mg^{2+} -activated adenosine triphosphatases of Escherichia coli and Salmonella typhimurium. Arch. Biochem. Biophys. 167: 311-321.

Braun, V., and Rehn, K. 1969. Chemical characterization, spatial distribution, and function of a lipoprotein (murein-lipoprotein) of Escherichia coli cell wall: The specific effect of trypsin on the membrane structure. Eur. J. Biochem. 10: 426-438.

Braun, V., and Sieglin, U. 1970. The covalent murein-lipoprotein structure of the Escherichia coli cell wall. The attachment site of the lipoprotein on the murein. Eur. J. Biochem. 13: 336-346.

Braun, V., and Wolff, H. 1970. The murein-lipoprotein linkage in the cell wall of Escherichia coli. Eur. J. Biochem. 14: 387-391.

Bronk, J.R., and Lesse, H.J. 1974. Accumulation of amino-acids and glucose by the mammalian small intestine. Symp. Soc. exp. Biol. 28: 283-304.

Burdett, I.D., and Murray, R.G.E. 1974. Septum formation in Escherichia coli: Characterization of septal structure and the effects of antibiotics on cell division. J. Bacteriol. 119: 303-324.

Carreira, J., Andreu, J.M., and Munoz, E. 1977. Differential sensitivity to trypsin digestion of purified forms of Micrococcus lysodeikticus ATPase. A study of their structural and conformational differences and mechanism of conversion. Biochim. Biophys. Acta 492: 387-398.

Carreira, J., Munoz, E., Andreu, J.M., and Nieto, M. 1976. Micrococcus lysodeikticus membrane ATPase: Effect of trypsin on stimulation of a purified form of the enzyme and identification of its natural inhibitor. Biochim. Biophys. Acta 436: 183-189.

Catterall, W.A., and Pedersen, P.L. 1971. ATPase from rat liver mitochondria. I. Purification, homogeneity, and physical properties. J. biol. Chem. 246: 4987-4994.

Chalcroft, J.P., Bullivant, S., and Howard, B.H. 1973. Ultrastructural studies on Selenomonas ruminantium from sheep rumen. J. Gen. Microbiol. 79: 135-146.

Chapman, G.B., and Hillier, J. 1953. Electron microscopy of ultra-thin sections of bacteria. I. Cellular division in Bacillus cereus. J. Bacteriol. 66: 362-373.

Cheng, K.-J., and Costerton, J.W. 1977. Alkaline phosphatase activity of rumen bacteria. Appl. Environ. Microbiol. 34: 586-590.

Chung, Y-S., and Salton, M.R.J. 1980. Inhibition of Micrococcus lysodeikticus membrane-bound and purified F₁-ATPase by N-ethoxycarbonyl-2-ethoxy-1, 2-dihydroquinoline. FEMS Letters 8: 183-186.

Clarke, D.J., and Morris, J.G. 1976. Partial purification of a dicyclohexylcarbodiimide-sensitive membrane adenosine triphosphatase complex from the obligately anaerobic bacterium Clostridium pasteurianum. Biochem. J. 154: 725-729.

Cohen-Bazire, B., and Kunisawa, R. 1963. The fine structure of Rhodospirillum rubrum. J. Cell Biol. 16: 401-419.

Costerton, J.W., Ingram, J.M., and Cheng, K.J. 1974.
Structure and function of the cell envelope of
Gram-negative bacteria. *Bacteriol. Rev.* 38: 87-110.

Costerton, J.W., Murray, R.G.E., and Robinow, C.F. 1961.
Observations on the motility and structure of
Vitreoscilla. *Can. J. Microbiol.* 7: 329-339.

Coulton, J.W., and Murray, R.G.E. 1977.
Membrane-associated components of the bacterial flagellar
apparatus. *Biochim. Biophys. Acta* 465: 290-310.

Coulton, J.W., and Murray, R.G.E. 1978. Cell envelope
association in Aquaspirillum serpens flagella. *J.*
Bacteriol. 136: 1037-1049.

Cox, G.B., Downie, J.A., Fayle, D.R.H., Gibson, F., and
Radik, J. 1978. Inhibition, by a protease inhibitor, of the
solubilization of the F₁-portion of the Mg²⁺-stimulated
adenosine triphosphatase of Escherichia coli. *J.*
Bacteriol. 133: 287-292.

Curtis, N.A.C., Richmond, M.H., and Sykes, R.B. 1972.
"Periplasmic" location of a beta-lactamase specified
either by a plasmid or a chromosomal gene in Escherichia
coli. *J. Bacteriol.* 112: 1433-1434.

Datta, D.B., Arden, B., and Henning, U. 1977. Major proteins of the Escherichia coli outer cell envelope membrane as bacteriophage receptors. J. Bacteriol. 131: 821-829.

Davies, P.L., and Bragg, P.D. 1972. Properties of a soluble Ca^{2+} - and Mg^{2+} -activated ATPase released from Escherichia coli membranes. Biochim. Biophys. Acta 266: 273-284.

Deguchi, N., Jorgensen, P.L., and Maunsbach, A.B. 1977. Ultrastructure of the sodium pump. Comparison of thin sectioning, negative staining and freeze-fracture of purified, membrane-bound (Na^+ , K^+)-ATPase. J. Cell Biol. 75: 619-634.

Delhez, J., Dufour, J.P., Thines, D., and Goffeau, A. 1977. Comparison of the properties of plasma membrane-bound ATPase in the yeast Schizosaccharomyces pombe. Eur. J. Biochem. 79: 319-328.

De Meis, L. and Vianna, A. 1979. Energy interconversion by the Ca^{2+} -dependent ATPase of sarcoplasmic reticulum. Annu. Rev. Biochem. 48: 275-292.

De Pamphilis, M.L., and Adler, J. 1971. Purification of intact flagella from Escherichia coli and Bacillus subtilis. J. Bacteriol. 105: 376-383.

De Petris, S. 1965. Ultrastructure of the cell wall of Escherichia coli. J. Ultrastruct. Res. 12: 247-262.

De Petris, S. 1967. Ultrastructure of the cell wall of Escherichia coli and chemical nature of its constituent layers. J. Ultrastruct. Res. 19: 45-83.

Dimmitt, K., and Simon, M. 1971. Purification and thermal stability of intact Bacillus subtilis flagella. J. Bacteriol. 105: 369-375.

Di Rienzo, J.M., Nakamura, K., and Inouye, M. 1978. The outer membrane proteins of Gram-negative bacteria: Biosynthesis, assembly and functions. Annu. Rev. Biochem. 47: 481-532.

Doetsch, R.N., and Sjöblad, R.D. 1980. Flagellar structure and function in eubacteria. Annu. Rev. Microbiol. 34: 69-108.

Downie, J.A., Gibsons, F., and Cox, G.B. 1979. Membrane ATPases of prokaryotic cells. Annu. Rev. Biochem. 48: 103-131.

Dufour, J.P., and Goffeau, A. 1978. Solubilization by lysolecithin and purification of the plasma membrane ATPase of the yeast Schizosaccharomyces pombe. J. Biol. Chem. 253: 7026-7032.

Dunn, S.D., Tozer, R.G., Antczak, D.F., and Heppel, L.A. 1985. Monoclonal antibodies to Escherichia coli F₁-ATPase. Correlation of binding site location with interspecies cross reactivity and effects on enzyme activity. J. Biol. Chem. 260: 10418-10425.

Dux, L., and Martonosi, A. 1983a. Two dimensional arrays of proteins in sarcoplasmic reticulum and purified Ca²⁺-ATPase vesicles treated with vanadate. J. Biol. Chem. 258: 2599-2603.

Dux, L., and Martonosi, A. 1983b. Ca²⁺-ATPase membrane crystals in sarcoplasmic reticulum. J. Biol. Chem. 258: 10111-10115.

Ellar, D.J., Munoz, E., and Salton, M.R.J. 1971. The effect of low concentrations of glutaraldehyde on Micrococcus lysodeikticus membranes: Changes in the release of membrane-associated enzymes and membrane structure. Biochim. Biophys. Acta, 225: 140-150.

Epstein, F. 1969. The role of sodium and potassium ATPase in renal sodium reabsorption. In Renal transport and diuretics (Thurau, K., and Jahrmarker, H., eds.), pp. 67-76. Springer, N.Y.

Epstein, W., Whitelaw, V., and Hesse, J. 1978. A K^+ transport ATPase in Escherichia coli. J. Biol. Chem. 253: 6666-6668.

Ernster, L., Carlsson, C., Hundal, T., and Norbenbrand, K. 1979. Mitochondrial ATPase Inhibitor: Properties and Applications. Methods in Enzymol. 55: 399-407.

Essmann, M., Christiansen, C., Karlsson, K.A., Hansson, G.C., and Skou, J.C. 1980. Hydrodynamic properties of solubilized $(Na^+ + K^+)$ -ATPase from rectal glands of Squalus acanthias. Biochim. Biophys. Acta 603: 1-12.

Evans, M.C.W. 1969. Progress in Photosynthetic Research. Vol. III, (Metzner, H., ed.), pp. 1474-1475. Tubingen. International Biol. Union.

Ferris, F.G., and Beveridge, T.J. 1983. Structure and cell envelope associations of flagellar basal complexes of Vibrio cholerae and Campylobacter fetus. Can. J. Microbiol. 30: 322-333.

Forbush, B., Kaplan, J.H., and Hoffman, J.F. 1978. characterization of a new photoaffinity derivative of ouabain: Labeling of the large polypeptide and a proteolipid component of the Na, K-ATPase. Biochem. 17: 3667-3676.

Forgacs, M.D., Cantely, L., Weidenna, B., Alstiel, L., and Branton, D. 1978. Clathrin-coated vesicles and ATP-dependent proton pump. Proc. Natl. Acad. Sci. U.S.A. 80: 1300-1303.

Foster, D.L., and Fillingame, R.H. 1978. Purification of a dicyclohexylcarbodiimide sensitive, energy transducing adenosine triphosphatase complex from Escherichia coli. Fed. Proc. 37: 1520, abst. 1379.

Foster, D.L., and Fillingame, R.H. 1979. Energy-transducing H^+ -ATPase of Escherichia coli: Purification, reconstitution and subunit composition. J. Biol. Chem. 254: 8230-8236.

Funahara, Y., and Nikaido, H. 1980. Asymmetric localization of lipopolysaccharides on the outer membrane of Salmonella typhimurium. J. Bacteriol. 141: 1463-1465.

Futai, M., and Kanazawa, H. 1980. Role of subunits in proton translocating ATPase (F_0F_1). Curr. Topics Bioenerg. 10: 181-215.

Futai, M., and Kanazawa, H. 1983. Structure and function of proton-translocating adenosine triphosphatase (F_0F_1): Biochemical and molecular biological approaches. Microbiol. Rev. 47: 285-312.

Futai, M., Sternweis, P.C., and Heppel, L.A. 1974. Purification and properties of reconstitutively active and inactive ATPase from Escherichia coli. Proc. Natl. Acad. Sci. U.S.A. 71: 2725-2729.

Gallagher, S.R., and Leonard, R.T. 1982. Effect of vanadate, molybdate and azide on membrane associated ATPase and soluble phosphatase activities of corn roots. Plant Physiol. 70: 1335-1340.

Galloway, C.J.; Dean, G.E., Marsh, M., Rudnick, g., and Mellman, I. 1983. Acidification of macrophage and fibroblast endocytic vesicles in vitro. Proc. Natl. Acad. Sci. U.S.A. 80: 3334-3338.

Garber, M.P., and Steponkus, P.L. 1974. Identification of chloroplast coupling factor by freeze-etching and negative staining techniques. J. Cell Biol. 63: 24-34.

Gay, N.J., and Walker, J.E. . 1981. The atp operon: nucleotide sequence of the region encoding the alpha subunit of Escherichia coli ATP-synthase. Nucleic Acids Res. 9: 2187-2194.

Ghuysen, J.M. 1968. Use of bacteriolytic enzymes in determination of cell wall structure and their role in cell metabolism. Bacteriol Rev. 32: 425-464.

Glauert, A.M., and Thornley, M.J. 1969. The topography of the bacterial cell wall. Annu. Rev. Microbiol. 23: 159-198.

Glickman, J., Croen, K., Kelly, S., and Al-Awqati, Q. 1983. Golgi membranes contain an electrogenic H⁺ pump in parallel to a chloride conductance. J. Cell. Biol. 97: 1303-1308.

Glynn, I.M. 1957. The action of cardiac glycosides on sodium and potassium movement in human red cells. J. Physiol. (London) 136: 148-173.

Goffeau, A., and Slayman, C.W. 1982. The proton-translocating ATPase of the fungal plasma membrane. Biochim. Biophys. Acta 639: 197-223.

Gram, C. 1884. Über die isolirte Färbung der Schizomyceten in Schnitt-und Trockenpräparaten. Fortschr. Med. 2: 185-189.

Hachimori, A., Maramatsu, N., and Nosch, Y. 1970. Studies on an ATPase of thermophilic bacteria I. Purification and properties. Biochim. Biophys. Acta. 206: 426-437.

Haddock, B.A., and Jones, C.W. 1977. Bacterial Respiration. Bacteriol. Rev. 41: 47-99.

Hansen, F.G., Nielson, J., Riise, E., and von Meyenburg, K. 1981. The genes for the eight subunits of the membrane bound ATP synthase of Escherichia coli. Mol. Gen. Genet. 183: 463-472.

Hanson, R.L., and Kennedy, E.P. 1973. Energy-transducing adenosine triphosphatase from Escherichia coli: purification, properties and inhibition by antibody. J. Bacteriol. 114: 772-781.

Hare, J.E. 1975. Purification and characterization of an ATPase complex from membranes of Escherichia coli which is sensitive to DCCD. Biochem. Biophys. Res. Commun. 66: 1329-1337.

Harold, F.M. 1972. Conservation and transformation of energy by bacterial membranes. *Bacteriol. Rev.* 36: 172-230.

Harold, F.M., Baarda, J.R., Baron, C., and Abrams, A. 1969. Inhibition of membrane-bound adenosine triphosphatase and cation transport in *Streptococcus faecalis* by N,N'-dicyclohexyl-carbodiimide. *J. Biol. Chem.* 244: 2261-2268.

Hasselbach, W., and Suko, J. 1974. Calcium and phosphate turnover in the sarcoplasmic membranes. *Biochem. Soc. Spec. Publ.* 4: 159-173.

Hastings, D.F., and Reynolds, J.A. 1979. Molecular weight of (Na⁺,K⁺)-ATPase from shark rectal gland. *Biochem.* 18: 817-821.

Herbert, H., Jorgensen, P.L., Skriver, E., and Maunsbach, A.B. 1982. Crystalline patterns of membrane-bound (Na⁺K⁺)-ATPase. *Biochim. Biophys. Acta* 689: 571-574.

Hess, H.H., Lees, M.B., and Derr, J.E. 1978. A linear Lowry-Folin assay for both water-soluble and sodium dodecyl sulfate-solubilized proteins. *Anal. Biochem.* 85: 295-300.

Hickman, D.D., and Frenkel, A.W. 1965. Observations on the structure of Rhodospirillum rubrum. J. Cell Biol. 25: 279-291.

Higashi, T., Kalra, V.K., Lee, S.H., Brogin, E., and Brodie, A.F. 1975. Energy-transducing membrane-bound coupling factor ATPase from Mycobacterium phlei. I. Purification, homogeneity and properties. J. Biol. Chem. 250: 6541-6548.

Hinkle, P.C., and McCarty, R.E. 1978. How cells make ATP. Sci. Amer. 238: 104-123.

Hobot, J.A., Carlemalm, E., Villigerr, W., and Kellenberger, E. 1984. Periplasmic gel: new concept resulting from the reinvestigation of bacterial cell envelope ultrastructure by new methods. J. Bacteriol. 160: 143-152.

Hodges, T.K. 1976. ATPases associated with membranes of plant cells. In Encyclopedia of Plant Physiology, New Series, vol. 2, Parts A, B. Berlin, Heidelberg, N.Y. Springer Verlag. 2A: 260-283.

Hodgkin, A.L., and Keynes, R.D. 1955. Active transport of cations in giant axons from Sepia and Loligo. J. Physiol. (London) 128: 28-60.

Hokin, L.E., Dahl, J.L., Deupree, J.D., Dixon, J.F., Hackney, J.F., and Perdue, F. 1973. Studies on the characterization of sodium-potassium transport adenosine triphosphatase. X. Purification of the enzyme from the rectal gland of Squalus acanthias. J. Biol. Chem. 248: 2593-2605.

Huberman, M., and Salton, M.R.J. 1979. Purification and properties of the latent F₁-ATPase of Micrococcus lysodeikticus. Biochim. Biophys. Acta 547: 230-240.

Hylemon, P. B., Wells, J.S., Kreig, N.R., and Jannasch, H.W. 1973. The genus Spirillum: A taxonomic study. Int. J. Syst. Bacteriol. 23: 340-380.

Inouye, M. 1979. Bacterial outer membranes. John Wiley and Sons, Inc., N.Y.

Inouye, M., and Yee, M.L. 1972. Specific removal of proteins from the envelope of Escherichia coli by protease treatments. J. Bacteriol. 112: 585-592.

Ishida, M., and Mizushima, S. 1969. Membrane ATPase of Bacillus megaterium. I. Properties of membrane ATPase and its solubilized form. J. Biochem. (Tokyo) 66: 33-43.

Ishikawa, S. 1970. Properties of an oxidative phosphorylation system reconstituted from coupling factors in Micrococcus lysodeikticus. J. Biochem. (Tokyo) 67: 297-312.

Ishikawa, S., and Lehninger, A.L. 1962. Reconstitution of oxidative phosphorylation in preparations from Micrococcus lysodeikticus. J. Biol. Chem. 237: 2401-2408.

Jilka, R.L., Martonosi, A.N., and Tillack, T.W. 1975. Effect of the purified $[Mg^{2+} + Ca^{2+}]$ -activated ATPase of sarcoplasmic reticulum upon the passive Ca^{2+} permeability and ultrastructure of phospholipid vesicles. J. Biol. Chem. 250: 7511-7524.

Josephson, L., and Cantely, L.C. 1977. Isolation of a potent (Na-K)ATPase inhibitor from striated muscle. Biochem. 16: 4572-4578.

Johansson, B.C., Baltscheffsky, M., Baltscheffsky, H.,
Baccarini-Melandri, A., and Melandri, M. 1973.

Purification and properties of a coupling factor
(Ca⁺-dependent adenosine triphosphatase) from
Rhodospirillum rubrum. Eur. J. Biochem. 40: 109-117.

Johansson, B.C., and Baltscheffsky, M. 1975. On subunit
composition of the coupling factor (ATPase) from
Rhodospirillum rubrum. FEBS Lett. 53: 221-224.

Johansson, K.-E., and Hjerten, S. 1974. Localization of
the Tween20-soluble membrane proteins of Acholeplasma
laidlawii by crossed immunoelectrophoresis. J. Mol. Biol.
86: 341-348.

Johansson, K.-E., and Wroblewski, H. 1978. Crossed
immunoelectrophoresis in the presence of Tween 20 or
sodium deoxycholate of purified membrane proteins from
Acholeplasma laidlawii. J. Bacteriol. 136: 324-330.

Johnson, R.C., Walsh, M.P., Ely, B., and Shapiro, L. 1979.
Flagellar hook and basal complex of Caulobacter
crescentus. J. Bacteriol. 138: 984-989.

Jones, C.W., and Redfern, E.R. 1967. The cytochrome system
of Azotobacter vinelandii. Biochim. Biophys. Acta 143:
340-353.

Kagawa, Y., and Racker, E. 1966. Partial resolution of the enzymes catalysing oxidative phosphorylation. IX.

Reconstruction of oligomycin-sensitive adenosine triphosphatase. J. Biol. Chem. 241: 2467-2474.

Kagawa, Y., Sone, N., Yoshida, M., Hirita, H., and Okamoto, H. 1976. Proton translocating ATPase of a thermophilic bacterium. Morphology, subunits and chemical composition. J. Biochem. (Tokyo) 80: 141-151.

Kalra, V.K., Lee, S., Ritz, C.J., and Brodie, A.F. 1975. Purification and properties of membrane-bound coupling factor-latent ATPase from Mycobacterium phlei. J. Supramolecular. Struct. 3: 231-241.

Kaneda, T. 1977. Fatty acids of the genus Bacillus. An example of branched-chain preference. Bacteriol. Rev. 41: 391-418.

Kanazawa, H., Mabuchi, K., Kayano, T., Noumi, T., Sekiya, T., and Futai, M. 1981. Nucleotide sequence of the genes for F_o components of the proton-translocating ATPase from Escherichia coli: prediction of the primary structure of F_o subunits. Biochem. Biophys. Res. Commun. 103: 613-620.

Kandler, O. 1982. Cell wall structures and their phylogenetic implication. Zentralblatt für Bakteriologie, Mikrobiologie und Hygiene, I Abteilung, Originale C-Allgemeine, Angewandte und Ökologische Mikrobiologie 3: 149-160.

Karmali, M.A., and Skirrow, M.B. 1984. Taxonomy of the genus Campylobacter. In Campylobacter Infection in Man and Animals (Butzler, J.P., ed.). CRC Press Inc., Boca Raton, Fl. pp. 1-20.

Keeler, R.F., Ritchie, A.E., Bryner, J.H., and Elmore, J. 1966. The preparation and characterization of cell walls and the preparation of flagella in Vibrio fetus. J. Gen. Microbiol. 43: 439-454.

Kellenberger, E., and Ryter, A. 1958. Cell wall and cytoplasmic membrane of Escherichia coli. J. Biophys. Biochem. 4: 323-325.

Kellermann, O., and Szmecman, S. 1980. Active transport of maltose in Escherichia coli K12. Involvement of a "periplasmic" binding protein. Eur. J. Biochem. 47: 139-149.

Kinne Saffran, E., and Kinne, R. 1974. Presence of a bicarbonate stimulated ATPase in brush border microvillus membranes of proximal tubules. Proc. Soc. Exp. Med. 146: 751-753.

Knowles, A.F., and Penesky, H.S. 1972. The subunit structure of beef heart mitochondrial atpase: physical and chemical properties of isolated subunits. J. Biol. Chem. 247: 6624-6630.

Kobayashi, H., and Ankaku, Y. 1972. Membrane-bound adenosine triphosphatase of Escherichia coli. I. Partial purification and properties. J. Biochem. (Tokyo) 71: 387-399.

Koval, S.F., and Jarrel, K.F. 1986. Ultrastructure and biochemistry of the cell wall of Methanococcus voltae. J. Bacteriol. 169: 1298-1306.

Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. Nature (London) 227: 680-685.

Lechevalier, M.P. 1977. Lipids in bacterial taxonomy -- a taxonomists view. CRC Critic. Rev. Microbiol. 5: 109-210.

Lee, S., Cohen, N.S., and Brodie, A.F. 1976. Restoration of oxidative phosphorylation by purified N,N'-dicyclohexylcarbodiimide-sensitive latent adenosine triphosphatase from Mycobacterium phlei. Proc. Natl. Acad. Sci. U.S.A. 73: 3050-3053.

Leonard, E.P., and Provenza, D.V. 1972. Comparative study of the various methods for the ultrastructural localization of alkaline phosphatase activity. Histochemie. 30: 1-12.

Lien, S., and Racker, E. 1971. Preparation and assay of chloroplast coupling factor CF₁. Meth. Enzymol. 23: 547-555.

Lim, S-H., and Salton, M.R.J. 1981. F₁-ATPase of Micrococcus lysodeikticus is not a glycoprotein. Biochem. Biophys. Acta 638: 275-281.

Livne, A., and Racker, E. 1969. Partial resolution of the enzymes catalysing photophosphorylation. V. Interaction of coupling factor 1 from chloroplasts with ribonucleic acid and lipids. J. Biol. Chem. 244: 1332-1338.

Lo, T.C.Y. 1979. The molecular mechanisms of substrate transport in Gram-negative bacteria. Can. J. Biochem. 57: 289-330.

Lucke, F.K., and Klemme, J.H. 1976. Coupling factor adenosine-5' triphosphatase from Rhodospirillum rubrum. Simple and rapid procedure for its purification. Z. Naturforsch. 31: 272-279.

Luderitz, O., Westphal, O., Staub, A.M., and Nikaido, H. 1971. Isolation and chemical and immunological characterization of bacterial lipopolysaccharides. In Chapt. 4. Microbiol Toxins, vol. 4. Bacterial Endotoxins (Weinbaum, G., Kadis, S., and Ajl, S.J., eds.) Acad. Press, N.Y., and London. pp.145-233.

MacFarland, B.H., and Inesi, G. 1971. Solubilization of sarcoplasmic reticulum with triton x-100. Arch. Biochem. Biophys. 145: 456-464.

MacGregor, C.H., Bishop, C.W., and Blech, J.E. 1979. Localization of proteolytic activity in the outer membrane of Escherichia coli. J. Bacteriol. 137: 574-583.

Mack Ivey, D., and Ljungdahl, L.G. 1986. Purification and characterization of the F₁-ATPase from Clostridium thermoaceticum. J. Bacteriol. 165: 252-257.

- MacLennan, D.H., Seeman, P., Iles, G.H., and Yop, C.C.
1971. Membrane formation by the adenosine triphosphatase
of sarcoplasmic reticulum. J. Biol. Chem. 246: 2702-2710.
- Macnab, R.M. 1979. How do flagella propel bacteria. Trends
Biochem. Sci. (Pers. ed) 4: N10-N13.
- Malpartida, F., and Serrano, R. 1980. Purification of the
yeast plasma membrane ATPase solubilized with a novel
zwitterionic detergent. FEBS Lett. 111: 69-72.
- Malpartida, F., and Serrano, R. 1981. Phosphorylated
intermediate of the ATPase form the Plasma membrane of
yeast. Eur. J. Biochem. 116: 413-417.
- Manning, P.A., and Reeves, P. 1976. Outer membrane of
Escherichia coli K12: Differentiation of protein 3A and
3B on acrylamide gels and further characterization of
con(tolG) mutants. J. Bacteriol. 127: 1070-1079.
- Manning, P.A., Puspurs, A., and Reeves, P. 1976. Outer
membrane of Escherichia coli K12. Isolation of mutants
with altered protein 3A by using host range mutants of
bacteriophage K3. J. Bacteriol. 127: 1080-1084.

Matsubara, H., Hase, T., Hashimoto, T., and Tagawa, K.
1981. Amino acid sequences of an intrinsic inhibitor of
mitochondrial ATPase from yeast. J. Biochem. 90:
1159-1165.

Maunsbach, A.B., Skriver, E., Herbert, H., and Jorgensen,
P.L. 1983. Electron Microscope analysis of two dimensional
crystals of membrane bound Na,K-ATPase. Curr. topics Memb.
Trans. 19: 123-126.

McCarty, R.E. 1979. Interaction between nucleotides and
coupling factor 1 in chloroplasts. Trends Biochem. Sci. 4:
28-30.

McCarty, R.E., and Racker, E. 1966. Effect of a coupling
factor and its antiserum on photophosphorylation and
hydrogen ion transport. Brookhaven Symp. biol. 19:
202-214.

McCoy, E.C., Wiltberger, H.A., and Winter, A.J. 1976.
Major outer membrane protein of Campylobacter fetus:
Physical and immunological characterization. Infect.
Immun. 13: 1258-1265.

Meissner, G., Conner, G.E., and Fleischer, D. 1973. Isolation of sarcoplasmic reticulum by zonal centrifugation and purification of Ca^{2+} -pump and Ca^{2+} -binding proteins. *Biochim. Biophys. Acta* 298: 246-269.

Melandri, B.A., Baccarini-Melandri, A., Crofts, A.R., and Codgell, R. 1972. Energy transduction in photosynthetic bacteria. V. Role of coupling factor ATPase in energy conversion as revealed by light of ATP-induced quenching of atebvine fluorescence. *FEBS Lett.* 24: 141-145.

Merrell, B.R., Walker, R.I., and Coolbaugh, J.C. 1981. Campylobacter fetus s.s. jejuni, a newly recognized enteric pathogen: morphology and intestinal colonization. *Scanning Electron. Microsc.* 4: 125-131.

Mirsky, R., and Barlow, V. 1971. Purification and properties of ATPase from the cytoplasmic membrane of Bacillus megaterium KM. *Biochim. Biophys. Acta* 241: 835-845.

Mitchel, P. 1966. Chemiosmotic coupling in oxidative and photosynthetic phosphorylation. *Biol. Rev. Cambridge Philos. Soc.* 41: 445-502.

Mitchell, P. 1981. Biochemical mechanism of proton-motivated phosphorylation in F_0F_1 adenosine triphosphatase molecules. In *Mitochondria and micorsomes* (Lee, C.P., and schatz, G., eds.). Addison-Wesley Pub. co., Inc., Reading, Mass. pp. 427-457.

Moller, J.V., Andersen, J.P., and le Maire, M. 1982. The sarcoplasmic retivulum Ca^{2+} -ATPase. *Molecular Cellular Biochem.* 42: 83-107.

Mollinedo, F., Larraga, v., Coll, F.J., and Munoz, E. 1980. Role of the subunits of the energy transducing adenosine triphosphatase from Micrococcus lysodeikticus membranes, studied by proteolytic digestion and immunological approaches. *Biochem. J.* 186: 713-723.

Mollinedo, F., Larraga, V., and Munoz, E. 1983. Comparative aspects of the attachment of F_1 -ATPase to Micrococcus lysodeikticus membranes: Role of ions and subunits. *J. Gen. Microbiol.* 129: 3465-3472.

Monteil, H., and Serrahina-Zieger, M. 1978., Les ATPase Bacteriennes: Proprietes moleculaires et fonctions. *Bull. Inst. Pasteur* 76: 207-246.

Munkres, M., and Wachtel, A. 1967. Histochemical localization of phosphatases in Mycoplasma gallisepticum. J. Bacteriol. 93: 1096-1103.

Muñoz, E. 1982. Polymorphism and conformational dynamics of F₁-ATPase from bacterial membranes. A model for regulation of these enzymes on the basis of molecular plasticity. Biochim. Biophys. Acta 650: 233-265.

Muñoz, E., Freer, J.H., Ellar, D.J., and Salton, M.R.J. 1968. Membrane associated ATPase activity from Micrococcus lysodeikticus. Biochim. Biophys. Acta 150: 531-533.

Muñoz, E., Salton, M.R.J., Ng, M.H., and Schor, M.T. 1969. Membrane adenosine triphosphatase of Micrococcus lysodeikticus. Purification, properties of the "soluble" enzyme and properties of the membrane-bound enzyme. Eur. J. Biochem. 7: 490-501.

Murray, R.G.E. 1957. Direct evidence for a cytoplasmic membrane in sectioned bacteria. Can. J. Microbiol. 3: 531-532.

Murray, R.G.E., and Birch-Andersen, A. 1963. Specialized structure in the region of the flagella tuft in Spirillum serpens. Can. J. Microbiol. 9: 393-401.

Murray, R.G.E., Steed, P., and Elson, H.E. 1965. The location of the mucopeptide in sections of the cell wall of Escherichia coli and other Gram-negative bacteria. Can. J. Microbiol. 11: 547-560.

Nakae, T. 1976a. Identification of the outer membrane protein of Escherichia coli that produces transmembrane channels in reconstituted vesicle membranes. Biochem. Biophys. Res. Commun. 71: 877-884.

Nakae, T. 1976b. Outer membrane of Salmonella: Isolation of protein complex that produces transmembrane channels. J. Biol. Chem. 251: 2176-2178.

Nelson, N. 1976. Structure and function of chloroplas ATPase. Biochim. Biophys. Acta 456: 314-338.

Nelson, N., Kanner, B.I., and Gutnick, D.L. 1974. Purification and properties of $Mg^{2+}Ca^{2+}$ adenosine triphosphatase from Escherichia coli. Proc. Natl. Acad. Sci. U.S.A. 71: 2720-2724.

Nieuwenhuis, F.J.R.M., van der Drift, J.A.M., Voet, A.B., and Van Dam, K. 1974. Evidence for a naturally occurring ATPase-inhibitor in Escherichia coli. Biochim. Biophys. Acta 368: 461-463.

Ng, L.K., Sherburne, R., Taylor, D.E., and Stiles, M.E.
1985. Morphological forms and viability of Campylobacter
species studied by electron microscopy. J. Bacteriol. 164:
338-343.

O'Farrell, P.H. 1975 High resolution two-dimensional
electrophoresis of proteins. J. Biol. Chem. 250:
4007-4021.

Ogg, J.E. 1962. Studies on the coccoid form of ovine
Vibrio fetus. I. Cultural and serologic investigations.
Amer. J. Vet. Res. 23: 354-358.

Ohkuma, S., Moriyam, Y., and Takano, T. 1982.
Identification and characterization of a proton pump on
lysosomes by fluorescein isothiocyanate-dextran
fluorescence. Proc. Natl. Acad. Sci. U.S.A. 79: 2758-2762.

O'Neal, S.G., Rhoads, D.B., and Racker, E. 1979. Vanadate
inhibition of sarcoplasmic reticulum Ca^{2+} -ATPase and other
ATPases. Biochem., Biophys. Res. Commun. 89: 845-850.

Oppenheim, J.D., and Salton, M.R.J. 1973. Localization and
distribution of Micrococcus lysodeikticus membrane ATPase
determined by ferritin labeling. Biochim. Biophys. Acta
298: 297-322.

Osborn, M.J. 1968. Structure and biosynthesis of the bacterial cell wall. *Annu. Rev. Biochem.* 38: 501-538.

Osborn, M.J., Gander, J.E., Parisi, E., and Carson, J. 1972. Mechanism of assembly of the outer membrane of Salmonella typhimurium. Isolation and characterization of cytoplasmic and outer membrane. *J. Biol. Chem.* 247: 3962-3972.

Osborn, M.J., and Wu, H.C.P. 1980. Proteins of the outer membrane of Gram-negative bacteria. *Annu. Rev. Microbiol.* 34: 369-442.

Otero-Vilardebo, L.R., Lane, N., and Godman, G.C. 1963. Demonstration of mitochondrial ATPase activity in formalin-fixed colonic epithelial cells. *J. Cell Biol.* 19: 647-652.

Owen, P., and Salton, M.R.J. 1975. Antigenic and enzymatic architecture of Micrococcus lysodeikticus membranes established by crossed immunoelectrophoresis. *Proc. Natl. Acad. Sci. U.S.A.* 72: 3711-3715.

Owen, P., and Salton, M.R.J. 1977. Membrane asymmetry and expression of cell surface antigens of Micrococcus lysodeikticus established by crossed immunoelectrophoresis. *J. Bacteriol.* 132: 974-985.

Owen, P., and Smyth, C.J. 1977. Enzyme analysis by quantitative immunoelectrophoresis. In *Immunochemistry of Enzymes and Their Antibodies* (Salton, M.R.J., ed.), John Wiley, N.Y. pp. 147-202.

Pandit-Hovenkamp, H. G. 1967. Microbial phosphorylating preparations: Azotobacter. *Meth. Enzymol.* 10: 152-157.

Pedersen, P.L. 1975. Interaction of homogeneous mitochondrial ATPase from rat liver with adenine nucleotides and inorganic phosphate. *J. Supramolecul. Struct.* 3: 222-230.

Pedersen, P.L., Amzel, L.M., Soper, J.W., Cirtron, N., and Hüllihen J. 1978. Structure, function, and regulation of the mitochondrial adenosine-triphosphatase complex of rat-liver. In *Energy Conservation in Biological Membranes* (Schafer, G., and Klingenberg, M., eds.), Springer-Verlag, Berlin. pp. 159-194.

Pedersen, P.L., Hüllihen, J., and Wehrle, J.P. 1981. Proton adenosine triphosphatase complex of rat liver. The effect of trypsin on the F₁ and F₀ moieties of the enzyme. *J. Biol. Chem.* 256: 1362-1369.

Penefsky, H.S. 1974. Mitochondrial and chloroplasts ATPases. In The Enzymes (Boyer, P.D., ed.), Acad. Press, N.Y. and London. Vol. 10, pp. 375-394.

Penefsky, H.S. 1979. Mitochondrial ATPase. Adv. Enzymol. 49: 223-280.

Penefsky, H.S., and Warner, R. 1965. Partial resolution of the enzymes catalysing oxidative phosphorylation. VI Studies on the mechanism of cold-inactivation of mitochondrial adenosine triphosphatase. J. Biol. Chem. 240: 4694-4702.

Phelps, D.C., Nordenbrand, K., Nelson, B.D., and Ernster, L. 1975. Inhibition of purified mitochondrial ATPase (F₁) by bathophenanthroline and relief of the inhibition by uncouplers. Biochem. Biophys. Res. Commun. 63: 1005-1012.

Pinchot, G.B. 1953. Phosphorylation coupled to electron transport in cell-free extracts of Alcaligenes faecalis. J. Biol. Chem. 205: 65-74.

Pinchot, G.B. 1957. A polynucleotide coenzyme of oxidative phosphorylation. J. biol. Chem. 229: 1-9.

Pinchot, G.B., and Racker, E. 1951. Phosphorus Metabolism (McElroy, W.D., and Glass, B., eds.), Baltimore, John Hopkins Univ. Press. Vol. 1. p. 366.

Racker, E. 1967. Resolution and reconstitution of the inner mitochondrial membrane. Red. Proc. 26: 1335-1340.

Reed, D.W., and Raveed, D. 1972. Some properties of the ATPase from chromatophores of Rhodopseudomonas spheroides and its structural relationship to the bacteriochlorophyll proteins. Biochim. Biophys. Acta 283: 79-91.

Remsen, C.C., Watson, S.W., Waterbury, J.B., and Truper, H.B. 1968. Fine structure of Ectothiorhodospira mobilis Pelsh. J. Bacteriol. 95: 2374-2392.

Reynolds, E.S. 1963. The use of lead citrate at high pH as an electron opaque stain in electron microscopy. J. Cell Biol. 17: 208-213.

Riebeling, V., and Jungermann, K. 1976. Properties and function of Clostridial membrane ATPase. Biochim. Biophys. Acta 430: 434-444.

Risi, S., Hockel, M., Hulla, F.W., and Dose, K. 1977. F₁-ATPase from Micrococcus sp. ATCC 398. Purification by

ion-exchange chromatography and further characterization.
(auto)proteolysis and dissociative effects. Eur. J.
Biochem. 81: 103-109.

Ritchie, A.E., Keller, R.F., and Bryner, J.J. 1966.
Anatomical features of Vibrio fetus: electron microscopy
survey. J. Gen. Microbiol. 43: 427-438.

Ritz, C., and Brodie, A.F. 1977. Effects of trypsin
treatment on the structure and function of solubilized
coupling factor-latent ATPase of Mycobacterium phlei.
Biochem. Biophys. Res. Commun. 75: 933-939.

Ritz-Gold, C., and Brodie, A.F. 1979. Limited proteolysis
of coupling factor-latent ATPase from Mycobacterium phlei.
Effects of different enzymes and modifying agents.
Biochim. Biophys. Acta 547: 18-26.

Ritz-Gold, C.J., Gold, C.M., and Brodie, A.F. 1979.
Tryptic proteolysis of coupling factor-latent ATPase from
Mycobacterium phlei. Theoretical modeling of
structure-function relationships. Biochim. Biophys. ~~Acta~~
547: 1-17.

Rogers, H.J. 1983. Bacterial Cell Structure. In Aspects

of Microbiology, vol. 6 (Cole, J.A., Knowles, C.J., and Schiessinger, D., eds.), Amer. Soc. Microbiol. Publ., Washington, D.C.

Rogers, H.J., Perkins, H.R., and Ward, J.B. 1980. Microbial cell walls and membranes. Chapman and Hall, London, N.Y.

Rosenbusch, J.P. 1974. Characterization of the major envelope protein from Escherichia coli: Regular arrangement of the peptidoglycan and unusual dodecyl sulphate binding. J. Biol. Chem. 249: 8019-8029.

Roth, J. 1982. The protein A-gold (pAG) technique. Qualitative and quantitative approach for antigen localization on thin sections. In Techniques in Immunocytochemistry, Vol. 1 (Bullock, G.R., and Petrusz, P., eds.), Acad. Press, N.Y., and London. pp. 107-133.

Roth, J., Bendayn, M., Carlamalm, E., Billiger, W., and Garavito, R.M. 1981. Enhancement of structural preservation and immunocytochemical staining in low temperature embedded pancreatic tissue. J. Histochem. Cytochem. 29: 663-671.

Rottem, S., and Razin, S. 1967. Electrophoretic patterns of membrane proteins of Mycoplasma. J. Bacteriol. 94: 359-363.

Ryter, A., and Piechaud, M. 1963. Etude au microscope électronique de quelques souches de Moraxella. Ann. Inst. Past. 105: 1071.

Sabatine, D., Golman, D., Sabin, E., Sherman, J., Morimoto, T., Kreibich, G., and Adesnik, M. 1981. Mechanisms for the incorporation of proteins in membranes and organelles. J. Cell Biol. 92: 1-22.

Salton, M.R.J. 1953. Studies of the bacterial cell wall. IV. The composition of the walls of some Gram-positive and Gram-negative bacteria. Biochim. Biophys. Acta 10: 512-523.

Salton, M.R.J. 1971. The bacterial membrane. In Biomembranes I (Manson, L.A., ed.), Plenum, N.Y. pp.1-65.

Salton, M.R.J. 1974. Membrane associated enzymes in bacteria. In Advances in Microbial Physiology II. (Rose, A.H., and Tempest, D.W. eds.), pp.213-283.

Salton, M.R.J. 1978. Structure and function of bacterial plasma membranes. Symp. Soc. Gen. Microbiol. 28: 201-223.

Salton, M.R.J., and Owen, P. 1976. Bacterial membrane structure. Annu. Rev. Microbiol. 30: 451-482.

Salton, M.R.J., and Schor, M.T. 1972. Subunit structures and properties of two forms of ATPase released from Micrococcus lysodeikticus ATPase from membranes extracted with n-butanol. Biochem. Biophys. Res. Commun. 49: 350-357.

Salton, M.R.J., and Schor, M.T. 1974. Release and purification of Micrococcus lysodeikticus ATPase from membranes extracted with n-butanol. Biochim. biophys. Acta. 345: 74-82.

Salton, M.R.J., Freer, J.H., and Ellar, D.J. 1968. Electron transport components localized in a lipid-depleted sheet isolated from Micrococcus lysodeikticus membranes by deoxylate extraction. Biochem. Biophys. Res. Commun. 33: 909-915.

Salton, M.R.J., Schor, M.T., and Ng, M.H. 1972. Internal localization of Micrococcus lysodeikticus membrane ATPase by iodination with ^{125}I . Biochim. Biophys. Acta 290: 408-413.

Sarkadi, B. 1980. Active calcium transport in human red cells. Biochim. Biophys. Acta 604: 159-190.

Schleifer, K.H., and Kandler, O. 1972. Peptidoglycan types of bacterial cell walls and their taxonomic implications. Bacteriol. Rev. 36: 407-477.

Schnebli, H.P., and Abrams, A. 1970. Membrane adenosine triphosphate from Streptococcus faecalis. Preparation and homogeneity. J. Biol. Chem. 245: 1115-1121.

Schnebli, H.P., Vatter, A.E., and Abrams, A. 1970. Membrane adenosine triphosphate from Streptococcus faecalis. Molecular weight, subunit structure and amino acid composition. J. Biol. Chem. 245: 1122-1127.

Senior, A.E., Fayle, D.R.H., Downie, J.A., Gibson, F., and Cox, G.B. 1979. Properties of membranes from mutant strains of Escherichia coli in which the beta subunit of the adenosine triphosphatase is abnormal. Biochem. J. 180: 111-118.

Serrahina-Zieger, M., and Monteil, H. 1978. Membrane ATPase of Bacillus subtilis. I. Purification and properties. Biochim. Biophys. Acta 502: 445-457.

Serrano, R. 1984. Plasma membrane ATPase of fungi and plants as a novel type of proton pump. Curr. Topics Cell Regul. 23: 87-126.

Silverman, M.R., and Simon, M.I. 1972. Flagellar assembly mutants in Escherichia coli. J. Bacteriol. 112: 986-993.

Silverman, M.R., and Simon, M.I. 1977. Bacterial flagella. Annu. Rev. Microbiol. 31: 397-419.

Skriver, E., Maunsbach, A.B., and Jorgensen, P.L. 1981. Formation of two-dimensional crystals in pure membrane-bound Na^+ , K^+ -ATPase. FEBS Lett. 131: 219-222.

Smibert, R.M. 1965. Vibrio fetus var. intestinalis isolated from fecal and intestinal contents of clinically normal sheep: biochemical and cultural characteristics of microaerophilic vibrios isolated from the intestinal contents of sheep. Amer. J. Vet. Res. 26: 320-327.

Smibert, R.M. 1978. The genus Campylobacter. Annu. Rev. Microbiol. 32: 673-709.

Smibert, R.M., 1984. Genus Campylobacter Sebald and Veron. 1963, 907. In Bergy's Manual of Systematic Bacteriology. Vol. 1, (Krieg, N.R., and Holt, J.G., eds.), Williams and Wilkins, Baltimore and London. pp.111-118.

Smyth, C.J., Friedman-Kien, A.E., and Salton, M.R.J. 1976. Antigenic analysis of Neisseria gonorrhoeae by crossed immunoelectrophoresis. Infect. Immun. 13: 1273-1288.

Smyth, C.J., Seigel, J., Salton, M.R.J., and Owen, P.
1978. Immunochemical analysis of inner and outer membranes
of Escherichia coli by crossed immunoelectrophoresis. J.
Bacteriol. 133: 306-319.

Spinning, J.G., Strych, A., Price, A.H., Helander, H.F.,
and Sachs, G. 1973. Properties of ATPase of gastric
mucosa. V. Preparation of membranes and mitochondria by
zonal centrifugation. Biochim. Biophys. Acta 311: 545-564.

Steck, T.L., and Fox, C.F. 1972. Membrane proteins. In
Membrane Molecular Biology, (Fox, C.F., and Keith, A.D.,
eds.), Sinauer Assoc., Inc., Stamford, Calif. pp. 27-75.

Steven, A.C., tenHeggeler, B., Muller, R., Kistler, J.,
and Rosenbusch, J.P. 1977. Ultrastructure of a periodic
protein layer on the outer membrane of Escherichia coli.
J. Cell Biol. 72: 292-301.

Stoeckenium, W., and Rowen, R. 1967. A morphological study
of Halobacterium halobium and its lysis in media of low
salt concentration. J. Cell Biol. 34: 365-393.

Strötmann, H., Hesse, H., and Edelman, K. 1973.
Quantitative determination of coupling factor CF₁ of
chloroplasts. Biochim. Biophys. Acta 314: 202-210.

Sone, N., Yoshida, M., Hirata, H., and Kagawa, Y. 1975. Purification and properties of a dicyclohexylcarbodiimide sensitive ATPase from a thermophilic bacterium. J. Biol. Chem. 250: 7917-7923.

Stone, D.K., Xiao-Song, X., and Racker, E. 1983. An ATP-driven proton pump in clathrin-coated vesicles. J. Biol. Chem. 258: 4059-4062.

Sun, I.L., Phelps, D.C., and Crape, F.L. 1975. Lipophilic chelator inhibition of Escherichia coli membrane-bound ATPase activity and prevention of inhibition by uncouplers. FEBS Lett. 54: 253-258.

Terasaki, Y. 1970. Some observations on the life history of Spirillum serpens. Bull. Suzugamine Women's Coll. Nat. Sci. 15: 9-17.

Thompson, B.G., and Murray, R.G.E. 1982. The fenestrated peptidoglycan layer of Deinococcus radiodurans. Can. J. Microbiol. 28: 522-525.

Tikhonova, G.V., Biketov, S.F., Kasho, V.N., Kozolv, I.A., Mileihovshaya, E.I., Ostrovskii, D.N., Simakova, I.M., and Skulachev, V.P. 1983. Peculiarities of the subunit composition of H⁺-ATPase from the anaerobic bacterium Lactobacillus casei. Biokhimiya 48: 1348-1353.

Tonomura, Y., and Inoue, A. 1974. The substructure of myosin and the reaction mechanism of its adenosine triphosphatase. *Molecul. Cell Biochem.* 5: 127-143.

Towbin, H., Staehelin, T., and Gordon, J. 1979. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets. *Proc. Natl. Acad. Sci. U.S.A.* 76: 5340-5354.

Uchida, E., Ohsumi, Y., and Anraku, Y. 1985. Purification and properties of H⁺-translocating, Mg²⁺-adenosine triphosphatase from vacuolar membranes of Saccharomyces cerevisiae. *J. Biol. Chem.* 260: 1090-1095.

Urban, C., and Salton, M.R.J. 1983a. Antigenic expression of the Micrococcus lysodeikticus (luteus) F₁-ATPase as determined by immunoelectrophoretic analysis. *FEMS Microbiol. Lett.* 19: 97-101.

Urban, C., and Salton, M.R.J. 1983b. Immunochemical analysis of Micrococcus lysodeikticus (luteus) F₁-ATPase and its subunits. *Biochim. Biophys. Acta* 724: 230-240.

Vaituzis, Z. 1973. Localization of adenosine triphosphatase activity in motile bacteria. *Can. J. Microbiol.* 19: 1265-1267.

Vambutas, V.K., and Racker, E. 1965. Partial resolution of the enzymes catalyzing photophosphorylation. I.

Stimulation of photophosphorylation by a preparation of a latent Ca^{++} -dependent adenosine triphosphatase from chloroplasts. J. Biol. Chem. 240: 2660-2667.

Van de Stadt, R.J., Van Dam, K., and Slater, E.C. 1974.

Interaction of aurovertin with sub-mitochondrial particles, deficient in ATPase inhibitor. Biochim. Biophys. Acta 347: 224-239.

Van Iterson, W. 1961. Some features of a remarkable organelle in Bacillus subtilis. J. Biophys. Biochem. Cytol. 9: 183-192.

Voelz, H. 1964. Sites of adenosine triphosphatase activity in bacteria. J. Bacteriol. 88: 1196-1198.

Vogel, G., Schairer, H.U., and Steinhart, R. 1978.

Complementation in vitro of mutant and wild-type ATPase of Escherichia coli using isolated subunits. Eur. J. Biochem. 87: 155-160.

Vogel, G., and Steinhart, R. 1976. ATPase of Escherichia coli: Purification, dissociation, and reconstitution of the active complex from the isolated subunits. Biochem. 15: 208-216.

Wakabayashi, T., Kubota, M., Yoshida, M., and Kagawa, Y.
1977. Structure of ATPase (coupling factor TF₁) from a
thermophilic bacterium. J. Mol. Biol. 117: 515-519.

Ward, J.B. 1981. Teichoic and teichuronic acids:
Biosynthesis, assembly, and location. Microbiol. Rev. 45:
211-243.

Weidel, W. 1950. In Viruses (Delbruck, M., ed.),
California Inst. Tech. pp. 119-121.

Weiss, R.L. 1974. Subunit cell wall of Sulfolobus
acidocaldarius. J. Bacteriol. 118: 275-284.

Wetzel, B.K., Spicer, S.S., Dvorak, H.F., and Heppel, L.A.
1970. Cytochemical localization of certain phosphatases in
Escherichia coli. J. Bacteriol. 104: 529-542.

White, D.A., Lennerz, W.J., and Schnaitman, C.A. 1972.
Distribution of lipids in the wall and cytoplasmic
membrane subfraction of the cell envelope of escherichia
coli. J. Bacteriol. 109: 686-690.

Woese, L.R. 1987. Bacterial Evolution. Microbiol. Rev. 51
221-271.

Yoshida, M., Sone, N., Hirata, H., and Kagawa, Y. 1975. A highly stable adenosine triphosphatase from a thermophilic bacterium; purification, properties and reconstitution. J. Biol. Chem. 250: 7910-7916.