

Western University

Scholarship@Western

---

Paediatrics Publications

Paediatrics Department

---

12-1-2020

**Erratum: Correction to: Predicting Dropout from Children's Mental Health Services: Using a Need-Based Definition of Dropout (Child psychiatry and human development (2020) 51 1 (13-26))**

Kimberly W. Dossett  
*Western University*

Graham J. Reid  
*Western University, greid@uwo.ca*

Follow this and additional works at: <https://ir.lib.uwo.ca/paedpub>

---

**Citation of this paper:**

Dossett, Kimberly W. and Reid, Graham J., "Erratum: Correction to: Predicting Dropout from Children's Mental Health Services: Using a Need-Based Definition of Dropout (Child psychiatry and human development (2020) 51 1 (13-26))" (2020). *Paediatrics Publications*. 1879.  
<https://ir.lib.uwo.ca/paedpub/1879>



## Correction to: Predicting Dropout from Children's Mental Health Services: Using a Need-Based Definition of Dropout

Kimberly W. Dossett<sup>1</sup> · Graham J. Reid<sup>1,2,3,4</sup>

Published online: 22 April 2020  
© Springer Science+Business Media, LLC, part of Springer Nature 2020

### Correction to:

**Child Psychiatry & Human Development**, 51(1):13–26.  
<https://doi.org/10.1007/s10578-019-00906-4>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original version of this article was unfortunately published with an error in Table 2. The confidence intervals listed in Table 2 were published incorrectly. The correct version of confidence intervals in Table 2 should read as below.

**Table 2** Description and prevalence of dropout according to each definition

Dropout definition	Description of dropout	Prevalence of dropout % (95% CI)
Dose	Attending < 12 sessions within 16-weeks	93.5% (90.9, 95.2)
Clinician judgement	A coding at discharge indicated the child/family has dropped out, or refused additional treatment	53.3% (49.1, 57.6)
Need-based	Low need: attending < 8 sessions High need & married parents: attending < 16 sessions High need & single parent: attending < 24 sessions	63.0% (58.7, 66.9)

*Note:*  $N = 521$   
95% CI 95% confidence intervals.

The original article can be found online at <https://doi.org/10.1007/s10578-019-00906-4>.

✉ Kimberly W. Dossett  
kdossett@uwo.ca

<sup>1</sup> Department of Psychology, Western University, London, Canada

<sup>2</sup> Department of Family Medicine, Western University, London, Canada

<sup>3</sup> Department of Paediatrics, Western University, London, Canada

<sup>4</sup> Children's Health Research Institute, London, Canada